

NATIONAL QUALITY FORUM—Measure Testing (subcriteria 2a2, 2b1-2b6)

Measure Number (if previously endorsed): Click here to enter NQF number

Measure Title: Click here to enter measure title

Date of Submission: Click here to enter a date

Type of Measure:

<input type="checkbox"/> Outcome (including PRO-PM)	<input type="checkbox"/> Composite – STOP – use composite testing form
<input type="checkbox"/> Intermediate Clinical Outcome	<input type="checkbox"/> Cost/resource
<input type="checkbox"/> Process (including Appropriate Use)	<input type="checkbox"/> Efficiency
<input type="checkbox"/> Structure	

Instructions

- Measures must be tested for all the data sources and levels of analyses that are specified. **If there is more than one set of data specifications or more than one level of analysis, contact NQF staff** about how to present all the testing information in one form.
- **For all measures, sections 1, 2a2, 2b1, 2b2, and 2b4 must be completed.**
- **For outcome and resource use measures**, section **2b3** also must be completed.
- If specified for **multiple data sources/sets of specifications** (e.g., claims and EHRs), section **2b5** also must be completed.
- Respond to all questions as instructed with answers immediately following the question. All information on testing to demonstrate meeting the subcriteria for reliability (2a2) and validity (2b1-2b6) must be in this form. An appendix for *supplemental* materials may be submitted, but there is no guarantee it will be reviewed.
- If you are unable to check a box, please highlight or shade the box for your response.
- Maximum of 25 pages (including questions/instructions; minimum font size 11 pt; do not change margins). **Contact NQF staff if more pages are needed.**
- Contact NQF staff regarding questions. Check for resources at [Submitting Standards webpage](#).
- For information on the most updated guidance on how to address social risk factors variables and testing in this form refer to the release notes for version 7.1 of the Measure Testing Attachment.

Note: The information provided in this form is intended to aid the Standing Committee and other stakeholders in understanding to what degree the testing results for this measure meet NQF’s evaluation criteria for testing.

2a2. Reliability testing ¹⁰ demonstrates the measure data elements are repeatable, producing the same results a high proportion of the time when assessed in the same population in the same time period and/or that the measure score is precise. For **instrument-based measures** (including PRO-PMs) **and composite performance measures**, reliability should be demonstrated for the computed performance score.

2b1. Validity testing ¹¹ demonstrates that the measure data elements are correct and/or the measure score correctly reflects the quality of care provided, adequately identifying differences in quality. For **instrument-based measures (including PRO-PMs) and composite performance measures**, validity should be demonstrated for the computed performance score.

2b2. Exclusions are supported by the clinical evidence and are of sufficient frequency to warrant inclusion in the specifications of the measure; ¹²

AND

If patient preference (e.g., informed decisionmaking) is a basis for exclusion, there must be evidence that the exclusion impacts performance on the measure; in such cases, the measure must be specified so that the information about patient preference and the effect on the measure is transparent (e.g., numerator category computed separately, denominator exclusion category computed separately). ¹³

2b3. For outcome measures and other measures when indicated (e.g., resource use):

- **an evidence-based risk-adjustment strategy** (e.g., risk models, risk stratification) is specified; is based on patient factors (including clinical and social risk factors) that influence the measured outcome and are present at start of care; ^{14,15} and has demonstrated adequate discrimination and calibration

OR

- rationale/data support no risk adjustment/ stratification.

2b4. Data analysis of computed measure scores demonstrates that methods for scoring and analysis of the specified measure allow for identification of statistically significant and practically/clinically meaningful ¹⁶ differences in performance;

OR

there is evidence of overall less-than-optimal performance.

2b5. If multiple data sources/methods are specified, there is demonstration they produce comparable results.

2b6. Analyses identify the extent and distribution of missing data (or nonresponse) and demonstrate that performance results are not biased due to systematic missing data (or differences between responders and nonresponders) and how the specified handling of missing data minimizes bias.

Notes

10. Reliability testing applies to both the data elements and computed measure score. Examples of reliability testing for data elements include, but are not limited to: inter-rater/abstractor or intra-rater/abstractor studies; internal consistency for multi-item scales; test-retest for survey items. Reliability testing of the measure score addresses precision of measurement (e.g., signal-to-noise).

11. Validity testing applies to both the data elements and computed measure score. Validity testing of data elements typically analyzes agreement with another authoritative source of the same information. Examples of validity testing of the measure score include, but are not limited to: testing hypotheses that the measures scores indicate quality of care, e.g., measure scores are different for groups known to have differences in quality assessed by another valid quality measure or method; correlation of measure scores with another valid indicator of quality for the specific topic; or relationship to conceptually related measures (e.g., scores on process measures to scores on outcome measures). Face validity of the measure score as a quality indicator may be adequate if accomplished through a systematic and transparent process, by identified experts, and explicitly addresses whether performance scores resulting from the measure as specified can be used to distinguish good from poor quality. The degree of consensus and any areas of disagreement must be provided/discussed.

12. Examples of evidence that an exclusion distorts measure results include, but are not limited to: frequency of occurrence, variability of exclusions across providers, and sensitivity analyses with and without the exclusion.

13. Patient preference is not a clinical exception to eligibility and can be influenced by provider interventions.

14. Risk factors that influence outcomes should not be specified as exclusions.

15. With large enough sample sizes, small differences that are statistically significant may or may not be practically or clinically meaningful. The substantive question may be, for example, whether a statistically significant difference of one percentage point in the percentage of patients who received smoking cessation counseling (e.g., 74 percent v. 75 percent) is clinically meaningful; or whether a statistically significant difference of \$25 in cost for an episode of care (e.g., \$5,000 v. \$5,025) is practically meaningful. Measures with overall less-than-optimal performance may not demonstrate much variability across providers.

1. DATA/SAMPLE USED FOR ALL TESTING OF THIS MEASURE

Often the same data are used for all aspects of measure testing. In an effort to eliminate duplication, the first five questions apply to all measure testing. If there are differences by aspect of testing, (e.g., reliability vs. validity) be sure to indicate the specific differences in question 1.7.

1.1. What type of data was used for testing? (Check all the sources of data identified in the measure specifications and data used for testing the measure. Testing must be provided for all the sources of data specified and intended for measure implementation. **If different data sources are used for the numerator and denominator, indicate N [numerator] or D [denominator] after the checkbox.**)

Measure Specified to Use Data From: (must be consistent with data sources entered in S.17)	Measure Tested with Data From:
<input type="checkbox"/> abstracted from paper record	<input type="checkbox"/> abstracted from paper record
<input type="checkbox"/> claims	<input type="checkbox"/> claims
<input type="checkbox"/> registry	<input type="checkbox"/> registry
<input type="checkbox"/> abstracted from electronic health record	<input type="checkbox"/> abstracted from electronic health record
<input type="checkbox"/> eMeasure (HQMF) implemented in EHRs	<input type="checkbox"/> eMeasure (HQMF) implemented in EHRs
<input type="checkbox"/> other: Click here to describe	<input type="checkbox"/> other: Click here to describe

1.2. If an existing dataset was used, identify the specific dataset (the dataset used for testing must be consistent with the measure specifications for target population and healthcare entities being measured; e.g., Medicare Part A claims, Medicaid claims, other commercial insurance, nursing home MDS, home health OASIS, clinical registry).

1.3. What are the dates of the data used in testing? [Click here to enter date range](#)

1.4. What levels of analysis were tested? (testing must be provided for all the levels specified and intended for measure implementation, e.g., individual clinician, hospital, health plan)

Measure Specified to Measure Performance of: (must be consistent with levels entered in item S.20)	Measure Tested at Level of:
<input type="checkbox"/> individual clinician	<input type="checkbox"/> individual clinician
<input type="checkbox"/> group/practice	<input type="checkbox"/> group/practice
<input type="checkbox"/> hospital/facility/agency	<input type="checkbox"/> hospital/facility/agency
<input type="checkbox"/> health plan	<input type="checkbox"/> health plan
<input type="checkbox"/> other: Click here to describe	<input type="checkbox"/> other: Click here to describe

1.5. How many and which measured entities were included in the testing and analysis (by level of analysis and data source)? (identify the number and descriptive characteristics of measured entities included in the analysis (e.g., size, location, type); if a sample was used, describe how entities were selected for inclusion in the sample)

1.6. How many and which patients were included in the testing and analysis (by level of analysis and data source)? (identify the number and descriptive characteristics of patients included in the analysis (e.g., age, sex, race, diagnosis); if a sample was used, describe how patients were selected for inclusion in the sample)

1.7. If there are differences in the data or sample used for different aspects of testing (e.g., reliability, validity, exclusions, risk adjustment), identify how the data or sample are different for each aspect of testing reported below.

1.8 What were the social risk factors that were available and analyzed? For example, patient-reported data (e.g., income, education, language), proxy variables when social risk data are not collected from each patient (e.g. census tract), or patient community characteristics (e.g. percent vacant housing, crime rate) which do not have to be a proxy for patient-level data.

2a2. RELIABILITY TESTING

Note: *If accuracy/correctness (validity) of data elements was empirically tested, separate reliability testing of data elements is not required – in 2a2.1 check critical data elements; in 2a2.2 enter “see section 2b2 for validity testing of data elements”; and skip 2a2.3 and 2a2.4.*

2a2.1. What level of reliability testing was conducted? (may be one or both levels)

- Critical data elements used in the measure** (e.g., inter-abstractor reliability; data element reliability must address ALL critical data elements)
- Performance measure score** (e.g., signal-to-noise analysis)

2a2.2. For each level checked above, describe the method of reliability testing and what it tests (describe the steps—do not just name a method; what type of error does it test; what statistical analysis was used)

2a2.3. For each level of testing checked above, what were the statistical results from reliability testing? (e.g., percent agreement and kappa for the critical data elements; distribution of reliability statistics from a signal-to-noise analysis)

2a2.4 What is your interpretation of the results in terms of demonstrating reliability? (i.e., what do the results mean and what are the norms for the test conducted?)

2b1. VALIDITY TESTING

2b1.1. What level of validity testing was conducted? (may be one or both levels)

- Critical data elements** (data element validity must address ALL critical data elements)
- Performance measure score**
 - Empirical validity testing**
 - Systematic assessment of face validity of performance measure score as an indicator** of quality or resource use (i.e., is an accurate reflection of performance on quality or resource use and can distinguish good from poor performance) **NOTE:** Empirical validity testing is expected at time of maintenance review; if not possible, justification is required.

2b1.2. For each level of testing checked above, describe the method of validity testing and what it tests (describe the steps—do not just name a method; what was tested, e.g., accuracy of data elements compared to authoritative source, relationship to another measure as expected; what statistical analysis was used)

2b1.3. What were the statistical results from validity testing? (e.g., correlation; t-test)

2b1.4. What is your interpretation of the results in terms of demonstrating validity? (i.e., what do the results mean and what are the norms for the test conducted?)

2b2. EXCLUSIONS ANALYSIS

NA no exclusions — skip to section [2b3](#)

2b2.1. Describe the method of testing exclusions and what it tests (describe the steps—do not just name a method; what was tested, e.g., whether exclusions affect overall performance scores; what statistical analysis was used)

2b2.2. What were the statistical results from testing exclusions? (include overall number and percentage of individuals excluded, frequency distribution of exclusions across measured entities, and impact on performance measure scores)

2b2.3. What is your interpretation of the results in terms of demonstrating that exclusions are needed to prevent unfair distortion of performance results? (i.e., the value outweighs the burden of increased data collection and analysis. *Note: If patient preference is an exclusion, the measure must be specified so that the effect on the performance score is transparent, e.g., scores with and without exclusion*)

2b3. RISK ADJUSTMENT/STRATIFICATION FOR OUTCOME OR RESOURCE USE MEASURES

If not an intermediate or health outcome, or PRO-PM, or resource use measure, skip to section [2b4](#).

2b3.1. What method of controlling for differences in case mix is used?

- No risk adjustment or stratification
- Statistical risk model with [Click here to enter number of factors](#) risk factors
- Stratification by [Click here to enter number of categories](#) risk categories
- Other, [Click here to enter description](#)

2b3.1.1 If using a statistical risk model, provide detailed risk model specifications, including the risk model method, risk factors, coefficients, equations, codes with descriptors, and definitions.

2b3.2. If an outcome or resource use component measure is not risk adjusted or stratified, provide rationale and analyses to demonstrate that controlling for differences in patient characteristics (case mix) is not needed to achieve fair comparisons across measured entities.

2b3.3a. Describe the conceptual/clinical and statistical methods and criteria used to select patient factors (clinical factors or social risk factors) used in the statistical risk model or for stratification by risk (e.g., potential factors identified in the literature and/or expert panel; regression analysis; statistical significance of $p < 0.10$; correlation of x or higher; patient factors should be present at the start of care)

Also discuss any “ordering” of risk factor inclusion; for example, are social risk factors added after all clinical factors?

2b3.3b. How was the conceptual model of how social risk impacts this outcome developed? Please check all that apply:

- Published literature
- Internal data analysis
- Other (please describe)

2b3.4a. What were the statistical results of the analyses used to select risk factors?

2b3.4b. Describe the analyses and interpretation resulting in the decision to select social risk factors (e.g. prevalence of the factor across measured entities, empirical association with the outcome, contribution of unique variation in the outcome, assessment of between-unit effects and within-unit effects.) Also describe the impact of adjusting for social risk (or not) on providers at high or low extremes of risk.

2b3.5. Describe the method of testing/analysis used to develop and validate the adequacy of the statistical model or stratification approach (describe the steps—do not just name a method; what statistical analysis was used)

Provide the statistical results from testing the approach to controlling for differences in patient characteristics (case mix) below.

If stratified, skip to [2b3.9](#)

2b3.6. Statistical Risk Model Discrimination Statistics (e.g., c-statistic, R-squared):

2b3.7. Statistical Risk Model Calibration Statistics (e.g., Hosmer-Lemeshow statistic):

2b3.8. Statistical Risk Model Calibration – Risk decile plots or calibration curves:

2b3.9. Results of Risk Stratification Analysis:

2b3.10. What is your interpretation of the results in terms of demonstrating adequacy of controlling for differences in patient characteristics (case mix)? (i.e., what do the results mean and what are the norms for the test conducted)

2b3.11. Optional Additional Testing for Risk Adjustment (*not required, but would provide additional support of adequacy of risk model, e.g., testing of risk model in another data set; sensitivity analysis for missing data; other methods that were assessed*)

2b4. IDENTIFICATION OF STATISTICALLY SIGNIFICANT & MEANINGFUL DIFFERENCES IN PERFORMANCE

2b4.1. Describe the method for determining if statistically significant and clinically/practically meaningful differences in performance measure scores among the measured entities can be identified

(describe the steps—do not just name a method; what statistical analysis was used? Do not just repeat the information provided related to performance gap in 1b)

2b4.2. What were the statistical results from testing the ability to identify statistically significant and/or clinically/practically meaningful differences in performance measure scores across measured entities? *(e.g., number and percentage of entities with scores that were statistically significantly different from mean or some benchmark, different from expected; how was meaningful difference defined)*

2b4.3. What is your interpretation of the results in terms of demonstrating the ability to identify statistically significant and/or clinically/practically meaningful differences in performance across measured entities? *(i.e., what do the results mean in terms of statistical and meaningful differences?)*

2b5. COMPARABILITY OF PERFORMANCE SCORES WHEN MORE THAN ONE SET OF SPECIFICATIONS
If only one set of specifications, this section can be skipped.

Note: *This item is directed to measures that are risk-adjusted (with or without social risk factors) OR to measures with more than one set of specifications/instructions (e.g., one set of specifications for how to identify and compute the measure from medical record abstraction and a different set of specifications for claims or eMeasures). It does not apply to measures that use more than one source of data in one set of specifications/instructions (e.g., claims data to identify the denominator and medical record abstraction for the numerator). **Comparability is not required when comparing performance scores with and without social risk factors in the risk adjustment model. However, if comparability is not demonstrated for measures with more than one set of specifications/instructions, the different specifications (e.g., for medical records vs. claims) should be submitted as separate measures.***

2b5.1. Describe the method of testing conducted to compare performance scores for the same entities across the different data sources/specifications *(describe the steps—do not just name a method; what statistical analysis was used)*

2b5.2. What were the statistical results from testing comparability of performance scores for the same entities when using different data sources/specifications? *(e.g., correlation, rank order)*

2b5.3. What is your interpretation of the results in terms of the differences in performance measure scores for the same entities across the different data sources/specifications? *(i.e., what do the results mean and what are the norms for the test conducted)*

2b6. MISSING DATA ANALYSIS AND MINIMIZING BIAS

2b6.1. Describe the method of testing conducted to identify the extent and distribution of missing data (or nonresponse) and demonstrate that performance results are not biased due to systematic missing data (or differences between responders and nonresponders) and how the specified handling of

missing data minimizes bias (*describe the steps—do not just name a method; what statistical analysis was used*)

2b6.2. What is the overall frequency of missing data, the distribution of missing data across providers, and the results from testing related to missing data? (*e.g., results of sensitivity analysis of the effect of various rules for missing data/nonresponse; if no empirical sensitivity analysis, identify the approaches for handling missing data that were considered and pros and cons of each*)

2b6.3. What is your interpretation of the results in terms of demonstrating that performance results are not biased due to systematic missing data (or differences between responders and nonresponders) and how the specified handling of missing data minimizes bias? (*i.e., what do the results mean in terms of supporting the selected approach for missing data and what are the norms for the test conducted; if no empirical analysis, provide rationale for the selected approach for missing data*)