

NQF 2009 Measure Maintenance Schedule

This is the list of measures due for maintenance review in 2009; categorized by topic area. Some of these measures may be subject to undergo maintenance review in 2010 dependent upon inclusion within CDP projects.

Measure Number	Title	Description	Steward	Potential Project Assignment
CARDIOVASCULAR				
0013	Blood pressure measurement	Percentage of patient visits with blood pressure measurement recorded among all patient visits for patients aged > 18 years with diagnosed hypertension.	American Medical Association - Physician Consortium for Performance Improvement	
0070	CAD: Beta-Blocker Therapy-Prior myocardial infarction (MI)	Percentage of patients with prior MI at any time who were prescribed beta-blocker therapy.	American Medical Association - Physician Consortium for Performance Improvement	
0065	Coronary Artery Disease (CAD): Symptom and Activity Assessment	Percentage of patients with CAD who were evaluated for both level of activity and anginal symptoms during one or more office visits.	American Medical Association - Physician Consortium for Performance Improvement	
0066	CAD: ACE inhibitor/angiotensin receptor blocker (ARB) Therapy	Percentage of patients with CAD who also have diabetes and/or LSVD who were prescribed ACE inhibitor or ARB therapy.	American Medical Association - Physician Consortium for Performance Improvement	
0067	CAD: Antiplatelet Therapy	Percentage of patients with CAD who were prescribed antiplatelet therapy.	American Medical Association - Physician Consortium for Performance Improvement	

Measure Number	Title	Description	Steward	Potential Project Assignment
0068	Ischemic Vascular Disease (IVD): Use of Aspirin or another Antithrombotic	Percentage of patients who have documentation of use of aspirin or another antithrombotic during the 12-month measurement period.	National Committee for Quality Assurance	
0071	Acute Myocardial Infarction (AMI): Persistence of Beta-Blocker Treatment After a Heart Attack	Percentage of patients whose days' supply of beta blockers dispensed is ≥ 135 days in the 180 days following discharge.	National Committee for Quality Assurance	
0072	CAD: Beta-Blocker Treatment after a Heart Attack	Percentage of patients who have a claim indicating beta blocker therapy or who received an ambulatory prescription for beta-blockers rendered within 7 days after discharge.	National Committee for Quality Assurance	
0073	IVD: Blood Pressure Management	Percentage of patients who, at their most recent blood pressure reading during the 12-month measurement period, had a blood pressure result of $< 140/90$ mm HG.	National Committee for Quality Assurance	Patient Outcomes
0074	CAD: Drug Therapy for Lowering LDL-Cholesterol	Percentage of patients with CAD who were prescribed a lipid – lowering therapy (based on current ACC/AHA guidelines)	American Medical Association - Physician Consortium for Performance Improvement	
0075	IVD: Complete Lipid Profile and LDL Control < 100	Percentage of patients with a full lipid profile completed during the 12-month measurement period with date of each component of the profile documented; LDL-C < 100 .	National Committee for Quality Assurance	Patient Outcomes
0076	CAD: optimally managed modifiable risk	Percentage of members who have optimally managed modifiable risk factors (LDL, tobacco non-use, blood pressure control, aspirin usage).	Minnesota Community Measurement	Patient Outcomes

Measure Number	Title	Description	Steward	Potential Project Assignment
0077	Heart Failure (HF): Assessment of Activity Level	Percentage of patient visits or patients with HF with assessment of activity level.	American Medical Association - Physician Consortium for Performance Improvement	
0078	Heart Failure (HF) : Assessment of Clinical Symptoms of Volume Overload (Excess)	Percentage of patient visits or patients with HF with assessment of clinical symptoms of volume overload (excess).	American Medical Association - Physician Consortium for Performance Improvement	
0079	Heart Failure (HF) : Left Ventricular Function Assessment	Percentage of patients with HF with quantitative or qualitative results of left ventricular function (LVF) assessment recorded.	American Medical Association - Physician Consortium for Performance Improvement	
0081	Heart Failure (HF) : ACEI/ ARB Therapy	Percentage of patients with HF who also have left ventricular systolic dysfunction (LVSD) who were prescribed ACE inhibitor or ARB therapy.	American Medical Association - Physician Consortium for Performance Improvement	
0083	Heart Failure (HF) : Beta-blocker therapy	Percentage of patients with HF who also have LVSD who were prescribed beta-blocker therapy.	American Medical Association - Physician Consortium for Performance Improvement	
0084	Heart Failure (HF) : Warfarin Therapy Patients with Atrial Fibrillation	Percentage of patients with HF who also have paroxysmal or chronic atrial fibrillation who were prescribed warfarin therapy.	American Medical Association - Physician Consortium for Performance Improvement	

Measure Number	Title	Description	Steward	Potential Project Assignment
0085	Heart Failure (HF) : Weight Measurement	Percentage of patient visits for patients with HF with weight measurement recorded.	American Medical Association - Physician Consortium for Performance Improvement	
0018	Controlling High Blood Pressure	Percentage of patients with last BP < 140/80 mm Hg.	National Committee for Quality Assurance	
0017	Hypertension Plan of Care	Percentage of patient visits during which either systolic blood pressure \geq 140 mm Hg or diastolic blood pressure \geq 90 mm Hg, with documented plan of care for hypertension.	American Medical Association - Physician Consortium for Performance Improvement	
0217	Surgery Patients with Recommended Venous Thromboembolism (VTE) Prophylaxis Ordered	Percentage of surgery patients with recommended Venous Thromboembolism (VTE) Prophylaxis ordered during admission	Centers for Medicare & Medicaid Services	
0218	Surgery Patients Who Received Appropriate Venous Thromboembolism (VTE) Prophylaxis Within 24 Hours Prior to Surgery to 24 Hours After Surgery End Time	Percentage of surgery patients who received appropriate Venous Thromboembolism (VTE) Prophylaxis within 24 hours prior to surgery to 24 hours after surgery end time	Centers for Medicare & Medicaid Services	
0162	Heart Failure: Angiotensin converting enzyme inhibitor (ACEI) for left ventricular systolic dysfunction (LVSD)	Percentage of AMI patients with LVSD and without ACEI contraindications who were prescribed an angiotensin converting enzyme inhibitor (ACEI) for left ventricular systolic dysfunction (LVSD)	Centers for Medicare & Medicaid Services	

Measure Number	Title	Description	Steward	Potential Project Assignment
0082	Heart Failure (HF) : Patient Education	Percentage of patients who were provided with patient education on disease management and health behavior changes during one or more visit(s).	American Medical Association - Physician Consortium for Performance Improvement	
DIABETES/METABOLIC/RENAL				
0055	Diabetes: Eye exam	Percentage of adult patients with diabetes aged 18-75 years who received a dilated eye exam or seven standard field stereoscopic photos with interpretation by an ophthalmologist or optometrist or imaging validated to match diagnosis from these photos during the reporting year, or during the prior year, if patient is at low risk** for retinopathy **Patient is considered low risk if the following criterion is met: has no evidence of retinopathy in the prior year	National Committee for Quality Assurance	
0056	Diabetes: Foot exam	Percentage of adult patients with diabetes aged 18-75 years who received a foot exam (visual inspection, sensory exam with monofilament, or pulse exam)	National Committee for Quality Assurance	
0057	Hemoglobin A1c testing	Percentage of adult patients with diabetes aged 18-75 years receiving one or more A1c test(s) per year	National Committee for Quality Assurance	
0059	Hemoglobin A1c management	Percentage of adult patients with diabetes aged 18-75 years with most recent A1c level greater than 9.0% (poor control)	National Committee for Quality Assurance	Patient Outcomes
0060	Hemoglobin A1c test for pediatric patients	Percentage of pediatric patients with diabetes with a HBA1c test in a 12-month measurement period	National Committee for Quality Assurance	
0061	Diabetes: Blood Pressure Management	Percentage of patient visits with blood pressure measurement recorded among all patient visits for patients aged > 18 years with diagnosed hypertension.	National Committee for Quality Assurance	Patient Outcomes

Measure Number	Title	Description	Steward	Potential Project Assignment
0062	Diabetes: Urine protein screening	Percentage of adult diabetes patients aged 18-75 years with at least one test for microalbumin during the measurement year or who had evidence of medical attention for existing nephropathy (diagnosis of nephropathy or documentation of microalbuminuria or albuminuria)	National Committee for Quality Assurance	
0063	Diabetes: Lipid profile	Percentage of adult patients with diabetes aged 18-75 years receiving at least one lipid profile (or ALL component tests)	National Committee for Quality Assurance	
0064	Diabetes Measure Pair: A Lipid management: low density lipoprotein cholesterol (LDL-C) <130, B Lipid management: LDL-C <100	Percentage of adult patients with diabetes aged 18-75 years with most recent (LDL-C) <130 mg/dL B: Percentage of patients 18-75 years of age with diabetes whose most recent LDL-C test result during the measurement year was <100 mg/dL	National Committee for Quality Assurance	Patient Outcomes
0003	Bipolar Disorder: Assessment for diabetes	Percentage of patients treated for bipolar disorder who are assessed for diabetes within 16 weeks after initiating treatment with an atypical antipsychotic agent	Center for Quality Assessment and Improvement in Mental Health	
PULMONARY				
0001	Asthma assessment	Percentage of patients who were evaluated during at least one office visit for the frequency (numeric) of daytime and nocturnal asthma symptoms	American Medical Association - Physician Consortium for Performance Improvement	
0025	Management plan for people with asthma	Percentage of patients for whom there is documentation that a written asthma management plan was provided either to the patient or the patient's caregiver OR, at a minimum, specific written instructions on under what conditions the patient's doctor should be contacted or the patient should go to the emergency room	IPRO	

Measure Number	Title	Description	Steward	Potential Project Assignment
0027	Smoking Cessation, Medical assistance: a. Advising Smokers to Quit, b. Discussing Smoking Cessation Medications, c. Discussing Smoking Cessation Strategies	Percentage of patients who received advice to quit smoking Percentage of patients whose practitioner recommended or discussed smoking cessation medications Percentage of patients whose practitioner recommended or discussed smoking cessation methods or strategies	National Committee for Quality Assurance	
0028	Measure pair: a. Tobacco Use Assessment, b. Tobacco Cessation Intervention	Percentage of patients who were queried about tobacco use one or more times during the two-year measurement period Percentage of patients identified as tobacco users who received cessation intervention during the two-year measurement period	American Medical Association	
0036	Use of appropriate medications for people with asthma	Percentage of patients who were identified as having persistent asthma during the measurement year and the year prior to the measurement year and who were dispensed a prescription for either an inhaled corticosteroid or acceptable alternative medication during the measurement year	National Committee for Quality Assurance	
0047	Asthma: pharmacologic therapy	Percentage of all patients with mild, moderate, or severe persistent asthma who were prescribed either the preferred long-term control medication (inhaled corticosteroid) or an acceptable alternative treatment	American Medical Association	
0058	Inappropriate antibiotic treatment for adults with acute bronchitis	Percentage of patients who were diagnosed with bronchitis and were dispensed an antibiotic on or within three days after the episode date	National Committee for Quality Assurance	
0080	Chronic Obstructive Pulmonary Disease (COPD): assessment of oxygen saturation	Percentage of patients with COPD with oxygen saturation assessed at least annually	American Medical Association - Physician Consortium for Performance Improvement	

Measure Number	Title	Description	Steward	Potential Project Assignment
0091	COPD: spirometry evaluation	Percentage of patients with COPD who had a spirometry evaluation documented	American Medical Association - Physician Consortium for Performance Improvement	
0102	COPD: inhaled bronchodilator therapy	Percentage of symptomatic patients with COPD who were prescribed an inhaled bronchodilator	American Medical Association - Physician Consortium for Performance Improvement	
CANCER				
0031	Breast Cancer Screening	Percentage of eligible women 50-69 who receive a mammogram in a two year period	National Committee for Quality Assurance	
0032	Cervical Cancer Screening	Percentage of women 18-64 years of age, who received one or more Pap tests during the measurement year or the two years prior to the measurement year	National Committee for Quality Assurance	
0034	Colorectal Cancer Screening	Percentage of adults 50-80 years of age who had appropriate screening for colorectal cancer (CRC) including fecal occult blood test during the measurement year or, flexible sigmoidoscopy during the measurement year or the four years prior to the measurement year or, double contrast barium enema during the measurement year or the four years prior to the measurement year or, colonoscopy during the measurement year or the nine years prior to the measurement year	National Committee for Quality Assurance	
0208	Family Evaluation of Hospice Care	Percentage of family members of all patients enrolled in a hospice program who give satisfactory answers to the survey instrument	National Cancer Institute	Palliative Care

Measure Number	Title	Description	Steward	Potential Project Assignment
0209	Comfortable Dying	Percentage of patients who were uncomfortable because of pain on admission to hospice whose pain was brought under control within 48 hours	National Cancer Institute	Palliative Care
0210	Proportion receiving chemotherapy in the last 14 days of life	Percentage of patients who died from cancer receiving chemotherapy in the last 14 days of life	National Cancer Institute	Palliative Care
0211	Proportion with more than one emergency room visit in the last days of life	Percentage of patients who died from cancer with more than one emergency room visit in the last days of life	National Cancer Institute	Palliative Care
0212	Proportion with more than one hospitalization in the last 30 days of life	Percentage of patients who died from cancer with more than one hospitalization in the last 30 days of life	National Cancer Institute	Palliative Care
0213	Proportion admitted to the ICU in the last 30 days of life	Percentage of patients who died from cancer admitted to the ICU in the last 30 days of life	Institute for Clinical and Evaluative Sciences	Palliative Care
0214	Proportion dying from Cancer in an acute care setting	Percentage of patients who died from cancer dying in an acute care setting	Institute for Clinical and Evaluative Sciences	Palliative Care
0215	Proportion not admitted to hospice	Percentage of patients who died from cancer not admitted to hospice	National Cancer Institute	Palliative Care
BONE/JOINT				
0037	Osteoporosis testing in older women	Percentage of female patients aged 65 and older who reported receiving a bone density test (BMD) to check for osteoporosis	National Committee for Quality Assurance	
0050	Osteoarthritis: functional and pain assessment	Percentage of patients with osteoarthritis who were assessed for function and pain	American Medical Association - Physician Consortium for Performance Improvement	

Measure Number	Title	Description	Steward	Potential Project Assignment
0051	Osteoarthritis: assessment for use of anti-inflammatory or analgesic over-the-counter (OTC) medications	Percentage of patient visits with assessment for use of anti-inflammatory or analgesic OTC medications	American Medical Association - Physician Consortium for Performance Improvement	
0052	Low back pain: use of imaging studies	Percentage of patients with new low back pain who received an imaging study (plain x-ray, MRI, CT scan) conducted on the episode start date or in the 28 days following the episode start date	National Committee for Quality Assurance	
0053	Osteoporosis management in women who had a fracture	Percentage of women 65 years and older who suffered a fracture and who had either a bone mineral density (BMD) test or prescription for a drug to treat or prevent osteoporosis in the six months after the date of fracture	National Committee for Quality Assurance	
0054	Arthritis: disease modifying antirheumatic drug (DMARD) therapy in rheumatoid arthritis	Percentage of patients 18 years and older, diagnosed with rheumatoid arthritis who have had at least one ambulatory prescription dispensed for a DMARD	National Committee for Quality Assurance	
GERIATRIC HEALTH/NURSING HOME				
0035	Fall risk management in older adults: a. Discussing fall risk, b. Managing fall risk	Percentage of patients aged 75 and older who reported that their doctor or other health provider talked with them about falling or problems with balance or walking Percentage of patients aged 75 and older who reported that their doctor or other health provider had done anything to help prevent falls or treat problems with balance or walking	National Committee for Quality Assurance	Nursing Homes
0195	Residents with a decline in their ability to move about in their room and the adjacent corridor.	Percentage of residents with a valid target assessment and a valid prior assessment, with a decline in their ability to move about in their room and the adjacent corridor.	Centers for Medicare & Medicaid Services	Nursing Homes

Measure Number	Title	Description	Steward	Potential Project Assignment
0196	Residents with a urinary tract infection	Percentage of residents on most recent assessment with a urinary tract infection	Centers for Medicare & Medicaid Services	Nursing Homes
0030	Urinary Incontinence Management in Older Adults - a. Discussing urinary incontinence, b. Receiving urinary incontinence treatment	Percentage of patients 65 years of age and older who reported having a urine leakage problem in the last six months and who discussed their urinary leakage problem with their current practitioner The percentage of patients 65 years of age and older who reported having a urine leakage problem in the last six months and who received treatment for their current urine leakage problem	National Committee for Quality Assurance	Nursing Homes
0029	Counseling on physical activity in older adults - a. Discussing Physical Activity, b. Advising Physical Activity	Percentage patients 65 years of age and older who reported: discussing their level of exercise or physical activity with a doctor or other health provider in the last 12 months Percentage patients 65 years of age and older who reported receiving advice to start, increase, or maintain their level of exercise or physical activity from a doctor or other health provider in the last 12 months	National Committee for Quality Assurance	Nursing Homes
0192	Residents who experience moderate to severe pain during the 7-day assessment period (risk-adjusted)	Percentage of residents who experience moderate to severe pain during the 7-day assessment period	Centers for Medicare & Medicaid Services	Nursing Homes
0186	Recently hospitalized residents who experienced moderate to severe pain at any time during the 7-day assessment period	Percentage of recently hospitalized patients with valid SNF PPS 14-day assessment, who experienced moderate to severe pain at any time during the 7-day assessment period	Centers for Medicare & Medicaid Services	Nursing Homes

Measure Number	Title	Description	Steward	Potential Project Assignment
0022	Drugs to be avoided in the elderly: a. Patients who receive at least one drug to be avoided, b. Patients who receive at least two different drugs to be avoided.	Percentage of patients ages 65 years and older who received at least one drug to be avoided in the elderly in the measurement year. Percentage of patients 65 years of age and older who received at least two different drugs to be avoided in the elderly in the measurement year.	National Committee for Quality Assurance	
0187	Recently hospitalized residents with pressure ulcers (risk adjusted)	Recently hospitalized residents with pressure ulcers	Centers for Medicare & Medicaid Services	Nursing Homes
0190	Nurse staffing hours - 4 parts	Percentage of daily work in hours by the entire group of nurses or nursing assistants spent tending to residents	Centers for Medicare & Medicaid Services	Nursing Homes
0191	Residents who lose too much weight	Percentage of residents on most recent assessment who lose too much weight	Centers for Medicare & Medicaid Services	Nursing Homes
0182	Residents whose need for more help with daily activities has increased	Percentage of residents with a valid target and a valid prior assessment whose need for more help with daily activities has increased	Centers for Medicare & Medicaid Services	Nursing Homes
0183	Low-risk residents who frequently lose control of their bowel or bladder	Percentage of residents with a valid target assessment and not qualifying as high risk who frequently lose control of their bowel or bladder	Centers for Medicare & Medicaid Services	Nursing Homes
0184	Residents who have a catheter in the bladder at any time during the 14-day assessment period. (risk adjusted)	Percentage of residents with a valid target assessment who have a catheter in the bladder at any time during the 14-day assessment period.	Centers for Medicare & Medicaid Services	Nursing Homes
0193	Residents who were physically restrained daily during the 7-day assessment period	Percentage of residents on most recent assessments who were physically restrained daily during the 7-day assessment period	Centers for Medicare & Medicaid Services	Nursing Homes

Measure Number	Title	Description	Steward	Potential Project Assignment
0194	Residents who spent most of their time in bed or in a chair in their room during the 7-day assessment period	Percentage of residents on most recent assessment. who spent most of their time in bed or in a chair in their room during the 7-day assessment period	Centers for Medicare & Medicaid Services	Nursing Homes
0197	Residents with worsening of a depressed or anxious mood.	Percentage of residents with a valid target assessment and a valid prior assessment with worsening of a depressed or anxious mood.	Centers for Medicare & Medicaid Services	Nursing Homes
0198	High-risk residents with pressure ulcers	Percentage of residents with a valid target assessment and one of the following inclusion criteria: 1. Impaired in mobility or transfer on the target assessment 2. Comatose on the target assessment 3. Suffer malnutrition on the target assessment who have pressure ulcers	Centers for Medicare & Medicaid Services	Nursing Homes
0199	Average-risk residents with pressure ulcers	Percentage of residents with a valid target assessment and not qualifying as high risk with pressure ulcers	Centers for Medicare & Medicaid Services	Nursing Homes
INFECTIOUS DISEASES				
0039	Flu shots for Adults Ages 50-64	Percentage of patients age 50-64 who report having received an influenza vaccination during the past influenza vaccination season	National Committee for Quality Assurance	
0040	Flu Shot for Older Adults	Percentage of patients age 65 and over who received an influenza vaccination from September through December of the year	National Committee for Quality Assurance	
0041	Influenza Vaccination	Percentage of patients who received an influenza vaccination	American Medical Association	
0043	Pneumonia vaccination status for older adults	Percentage of patients 65 years of age and older who ever received a pneumococcal vaccination	National Committee for Quality Assurance	
0044	Pneumonia Vaccination	Percentage of patients who ever received a pneumococcal vaccination	National Committee for Quality Assurance	

Measure Number	Title	Description	Steward	Potential Project Assignment
0151	Initial antibiotic received within 6 hours of hospital arrival	Percentage of pneumonia patients 18 years of age and older who receive their first dose of antibiotics within 6 hours after arrival at the hospital	Centers for Medicare & Medicaid Services, The Joint Commission	
MENTAL HEALTH				
0004	Initiation and Engagement of Alcohol and Other Drug Dependence Treatment: a. Initiation, b. Engagement	Percentage of adults aged 18 and over diagnosed with AOD abuse or dependence and receiving a related service who initiate treatment Assessment of the degree to which members engage in treatment with two additional AOD treatments within 30 days after initiating treatment.	National Committee for Quality Assurance	
0110	Bipolar Disorder and Major Depression: Appraisal for alcohol or chemical substance use	Percentage of patients with depression or bipolar disorder with evidence of an initial assessment that includes an appraisal for alcohol or chemical substance use	Center for Quality Assessment and Improvement in Mental Health	
0111	Bipolar Disorder: Appraisal for risk of suicide	Percentage of patients with bipolar disorder with evidence of an initial assessment that includes an appraisal for risk of suicide.	Center for Quality Assessment and Improvement in Mental Health	
0112	Bipolar Disorder: Level-of-function evaluation	Percentage of patients treated for bipolar disorder with evidence of level-of-function evaluation at the time of the initial assessment and again within 12 weeks of initiating treatment	Center for Quality Assessment and Improvement in Mental Health	
0185	Recently hospitalized residents with symptoms of delirium (risk-adjusted)	Percentage of recently hospitalized patients, with a valid SNF PPS 14-day assessment with symptoms of delirium	Centers for Medicare & Medicaid Services	Nursing Homes

Measure Number	Title	Description	Steward	Potential Project Assignment
0103	Major Depressive Disorder: Diagnostic Evaluation	Percentage of patients with a diagnosis of major depressive disorder who met the DSM-IV™ criteria during the visit in which the new diagnosis or recurrent episode was identified	American Medical Association - Physician Consortium for Performance Improvement	
0104	Major Depressive Disorder: Suicide Risk Assessment	Percentage of patients who had a suicide risk assessment completed at each visit	American Medical Association - Physician Consortium for Performance Improvement	
0105	New Episode of Depression: (a) Optimal Practitioner Contacts for Medication Management, (b) Effective Acute Phase Treatment, (c) Effective Continuation Phase Treatment	<p>Percentage of patients who were diagnosed with a new episode of depression and treated with antidepressant medication, and who had at least three follow-up contacts with a practitioner during the 84-day (12-week) Acute Treatment Phase.</p> <p>b. Percentage of patients who were diagnosed with a new episode of depression, were treated with antidepressant medication and remained on an antidepressant drug during the entire 84-day Acute Treatment Phase.</p> <p>c. Percentage of patients who were diagnosed with a new episode of depression and treated with antidepressant medication and who remained on an antidepressant drug for at least 180 days.</p>	National Committee for Quality Assurance	
0109	Bipolar Disorder and Major Depression: Assessment for Manic or hypomanic behaviors	Percentage of patients treated for depression who were assessed, prior to treatment, for the presence of current and/or prior manic or hypomanic behaviors.	Center for Quality Assessment and Improvement in Mental Health	

Measure Number	Title	Description	Steward	Potential Project Assignment
CHILD HEALTH				
0024	Body Mass Index (BMI) 2 through 18 years of age	Percentage children, 2 through 18 years of age, whose weight is classified based on BMI percentile for age and gender	National Initiative for Children's Healthcare Quality	
0038	Childhood Immunization Status	Percentage of children 2 years of age who had four DtaP/DT, three IPV, one MMR, three H influenza type B, three hepatitis B, one chicken pox vaccine (VZV) and four pneumococcal conjugate vaccines by their second birthday. The measure calculates a rate for each vaccine and two separate combination rates	National Committee for Quality Assurance	
0002	Appropriate testing for children with pharyngitis	Percentage of patients who were diagnosed with pharyngitis, prescribed an antibiotic, and who received a group A streptococcus test for the episode	National Committee for Quality Assurance	
0069	Appropriate treatment for children with upper respiratory infection (URI)	Percentage of children who were given a diagnosis of URI and were not dispensed an antibiotic prescription on or three days after the episode date	National Committee for Quality Assurance	
0026	Measure pair - a. Tobacco use prevention for infants, children and adolescents, b. Tobacco use cessation for infants, children and adolescents	Percentage of patients' charts showing either that there is no tobacco use/exposure or (if a user) that the current use was documented at the most recent clinic visit Percentage of patients with documented tobacco use or exposure at the latest visit who also have documentation that their cessation interest was assessed or that they received advice to quit	Institute for Clinical Systems Improvement	

Measure Number	Title	Description	Steward	Potential Project Assignment
0106	Diagnosis of attention deficit hyperactivity disorder (ADHD) in primary care for school age children and adolescents	Percentage of patients newly diagnosed with attention deficit hyperactivity disorder (ADHD) whose medical record contains documentation of Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) or Diagnostic and Statistical Manual for Primary Care (DSM-PC) criteria being addressed.	Institute for Clinical Systems Improvement	
0107	Management of attention deficit hyperactivity disorder (ADHD) in primary care for school age children and adolescents	Percentage of patients diagnosed with attention deficit hyperactivity disorder (ADHD) and on first-line medication whose medical record contains documentation of a follow-up visit twice a year.	Institute for Clinical Systems Improvement	
0108	ADHD: Follow-Up Care for Children Prescribed Attention-Deficit/Hyperactivity Disorder (ADHD) Medication.	<p>a. Initiation Phase: Percentage of children 6 – 12 years of age as of the Index Prescription Episode Start Date with an ambulatory prescription dispensed for and ADHD medication and who had one follow-up visit with a practitioner with prescribing authority during the 30-Day Initiation Phase.</p> <p>b. Continuation and Maintenance (C&M) Phase: Percentage of children 6 – 12 years of age as of the Index Prescription Episode Start Date with an ambulatory prescription dispensed for ADHD medication who remained on the medication for at least 210 days and who in addition to the visit in the Initiation Phase had at least two additional follow-up visits with a practitioner within 270 days (9 months) after the Initiation Phase ends.</p>	National Committee for Quality Assurance	

Measure Number	Title	Description	Steward	Potential Project Assignment
PERINATAL				
0012	Prenatal Screening for Human Immunodeficiency Virus (HIV)	Percentage of patients who gave birth during a 12-month period who were screened for HIV infection during the first or second prenatal care visit.	American Medical Association - Physician Consortium for Performance Improvement	
0015	Prenatal Blood Groups (ABO), D (Rh) Type	Percentage of patients who gave birth during a 12-month period who had a determination of blood group (ABO) and D (Rh) type by the second prenatal care visit.	American Medical Association - Physician Consortium for Performance Improvement	
0016	Prenatal Blood Group Antibody Testing	Percentage of patients who gave birth during a 12-month period who were screened for blood group antibodies during the first or second prenatal care visit.	American Medical Association - Physician Consortium for Performance Improvement	
0014	Prenatal Anti-D Immune Globulin	Percentage of D-negative, unsensitized patients who gave birth during a 12-month period who received anti-D immune globulin at 26-30 weeks gestation.	American Medical Association - Physician Consortium for Performance Improvement	
OTHER				
0019	Documentation of medication list in the outpatient record	Percentage of patients having a medication list in the medical record.	National Committee for Quality Assurance	
0020	Documentation of allergies and adverse reactions in the outpatient record	Percentage of patients having documentation of allergies and adverse reactions in the medical record.	National Association for Healthcare Quality	
0021	Therapeutic monitoring: Annual monitoring for patients on persistent medications	Percentage of patients 18 years and older who received at least 180-day supply of medication therapy for the selected therapeutic agent and who received annual monitoring for the therapeutic agent.	National Committee for Quality Assurance	

Measure Number	Title	Description	Steward	Potential Project Assignment
0023	Body Mass Index (BMI) in adults > 18 years of age	Percentage of adults with BMI documentation in the past 24 month.	City of New York Department of Health and Mental Hygiene	
0033	Chlamydia screening in women	Percentage of eligible women who were identified as sexually active who had at least one test for Chlamydia during the measurement year	National Committee for Quality Assurance	
0527	Prophylactic antibiotic received within 1 hour prior to surgical incision SCIP-Inf-2		Centers for Medicare & Medicaid Services	
0528	Prophylactic antibiotic selection for surgical patients		Centers for Medicare & Medicaid Services	
0529	Prophylactic antibiotics discontinued within 24 hours after surgery end time		Centers for Medicare & Medicaid Services	
0138	Urinary catheter-associated urinary tract infection for intensive care unit (ICU) patients	Percentage of intensive care unit patients with urinary catheter-associated urinary tract infections	Centers for Disease Control and Prevention	
0139	Central line catheter-associated blood stream infection rate for ICU and high-risk nursery (HRN) patients	Percentage of ICU and high-risk nursery patients, who over a certain amount of days acquired a central line catheter-associated blood stream infections over a specified amount of line-days	Centers for Disease Control and Prevention	
0140	Ventilator-associated pneumonia for ICU and high-risk nursery (HRN) patients	Percentage of ICU and HRN patients who over a certain amount of days have ventilator-associated pneumonia	Centers for Disease Control and Prevention	