NQF 2010 Time-Limited Review Schedule

Measure Number	Title	Description	Steward	Anticipated Review
CARDIO	ASCULAR/STRO	KE		
0466	Use of Patch during Conventional Carotid Endarterectomy	Percentage of patients undergoing conventional (non-eversion) carotid endarterectomy who have patch closure of the arteriotomy.	Society for Vascular Surgery	Quarter 3
0456	Participation in a Systematic National Database for General Thoracic Surgery	Participation in at least one multi- center, standardized data collection and feedback program that provides benchmarking of the physician's data relative to national and regional programs and uses process and outcome measures.	Society of Thoracic Surgeons	Quarter 3
0442	Functional Communication Measure: Writing	This measure describes the change in functional communication status subsequent to speech-language pathology treatment related to writing.	American Speech- Language- Hearing Association	Quarter 3
0443	Functional Communication Measure: Swallowing	This measure describes the change in functional communication status subsequent to speech-language pathology treatment of patients who exhibit difficulty in swallowing.	American Speech- Language- Hearing Association	Quarter 3
0444	Functional Communication Measure: Spoken Language Expression	This measure describes the change in functional communication status subsequent to speech-language pathology treatment related to spoken language expression.	American Speech- Language- Hearing Association	Quarter 3
0445	Functional Communication Measure: Spoken Language Comprehension	This measure describes the change in functional communication status subsequent to speech-language pathology treatment related to spoken language comprehension.	American Speech- Language- Hearing Association	Quarter 3
0446	Functional Communication Measure: Reading	This measure describes the change in functional communication status subsequent to speech-language pathology treatment of patients with reading disorders.	American Speech- Language- Hearing Association	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0447	Functional Communication Measure: Motor Speech	This measure describes the change in functional communication status subsequent to speech-language pathology treatment of patients who exhibit deficits in speech-production.	American Speech- Language- Hearing Association	Quarter 3
0448	Functional Communication Measure: Memory	This measure describes the change in functional communication status subsequent to speech-language pathology treatment of patients with memory deficits.	American Speech- Language- Hearing Association	Quarter 3
0449	Functional Communication Measure: Attention	This measure describes the change in functional communication status subsequent to speech-language pathology treatment of patients who have attention deficits	American Speech- Language- Hearing Association	Quarter 3
0465	Perioperative Anti-platelet Therapy for Patients undergoing Carotid Endarterectomy	Percentage of patients undergoing carotid endarterectomy (CEA) who are taking an anti-platelet agent (aspirin or clopidogrel) within 48 hours prior to surgery and are prescribed this medication at hospital discharge following surgery	Society for Vascular Surgery	Quarter 3
0458	Pulmonary Function Tests before major anatomic lung resection (pneumonectom y, lobectomy)	Percentage of thoracic surgical patients, >/= 18 years of age who underwent at least one pulmonary function test no more than 12 months prior to a major lung resection.	Society of Thoracic Surgeons	Quarter 3
0507	Stenosis measurement in carotid imaging studies	Percentage of final reports for carotid imaging studies (neck MR angiography [MRA], neck CT angiography [CTA], neck duplex ultrasound, carotid angiogram) performed that include direct or indirect reference to measurements of distal internal carotid diameter as the denominator for stenosis measurement	American Medical Association, American Medical Association - Physician Consortium for Performance Improvement	Quarter 4

Measure Number	Title	Description	Steward	Anticipated Review
PULMON	ARY			
0501	Confirmation of Endotracheal Tube Placement	Any time an endotracheal tube is placed into an airway in the Emergency Department or an endotracheal tube is placed by an outside provider and that patient arrives already intubated (EMS or hospital transfer) or when an airway is placed after patients arrives to the ED there should be some method attempted to confirm ETT placement	Cleveland Clinic	Quarter 4
DIABETE	S/METABOLIC/RI	ENAL		1
0370	Monitoring hemoglobin levels below target minimum	Percentage of all adult (>=18 years old) hemodialysis or peritoneal dialysis patients with ESRD >=3 months and who had Hb values reported for at least 2 of the 3 study months, who have a mean Hb <10.0 g/dL for a 3 month study period, irrespective of ESA use.	Centers for Medicare & Medicaid Services	Quarter 2
0417	Diabetic Foot & Ankle Care, Peripheral Neuropathy – Neurological Evaluation	Percentage of patients, 18 years or older, with diabetes aged 18 years and older with who had a lower extremity neurological exam with risk categorization performed and a treat plan established at least once within 12 months a diagnosis of diabetes mellitus who had a neurological examination of their lower extremities during one or more office visits within 12 months	American Podiatric Medical Association	Quarter 3
0416	Diabetic Foot & Ankle Care, Ulcer Prevention – Evaluation of Footwear	Percentage of patients aged 18 years and older with a diagnosis of diabetes mellitus who were evaluated for proper footwear and sizing during one or more office visits within 12 month	American Podiatric Medical Association	Quarter 3
0451	Call for a Measure of Glycemic Control with Intravenous Insulin Implementation	Intravenous insulin glycemic control protocol implemented for cardiac surgery patients with diabetes or hyperglycemia admitted into an intensive care unit	Society of Thoracic Surgeons, Lifescan, A Johnson & Johnson Company	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
CANCER	L			I
0377	Myelodysplastic Syndrome (MDS) and Acute Leukemias – Baseline Cytogenetic Testing Performed on Bone Marrow	Percentage of patients aged 18 years and older with a diagnosis of MDS or an acute leukemia who had baseline cytogenic testing performed on bone marrow.	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0379	Chronic Lymphocytic Leukemia (CLL) – Baseline Flow Cytometry	Percentage of patients aged 18 years and older with a diagnosis of CLL who had baseline flow cytometry studies performed	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0381	Oncology: Treatment Summary Documented and Communicated – Radiation Oncology	Percentage of patients with a diagnosis of cancer who have undergone brachytherapy or external beam radiation therapy who have a treatment summary report in the chart that was communicated to the physician(s) providing continuing care within one month of completing treatment	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0382	Oncology: Radiation Dose Limits to Normal Tissues	Percentage of patients with a diagnosis of cancer receiving 3D conformal radiation therapy with documentation in medical record that normal tissue dose constraints were established within five treatment days for a minimum of one tissue	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0384	Oncology: Pain Intensity Quantified – Medical Oncology and Radiation Oncology (paired with 0383	Percentage of visits for patients with a diagnosis of cancer currently receiving intravenous chemotherapy or radiation therapy in which pain intensity is quantified	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0383	Oncology: Plan	Percentage of visits for patients with a	American	Quarter 3
	of Care for Pain –	diagnosis of cancer currently receiving	Medical	
	Medical	intravenous chemotherapy or	Association -	
	Oncology and	radiation therapy who report having	Physician	
	Radiation	pain with a documented plan of care	Consortium for	
	Oncology (paired with 0384)	to address pain	Performance Improvement	
0388	Prostate Cancer:	Percentage of patients with prostate	American	Quarter 3
	Three-	cancer receiving external beam	Medical	
	Dimensional	radiotherapy to the prostate only who	Association -	
	Radiotherapy	receive 3D-CRT (three-dimensional	Physician	
		conformal radiotherapy) or IMRT	Consortium for	
		(intensity modulated radiation	Performance	
		therapy)	Improvement	
0389	Prostate Cancer:	Percentage of patients with a	American	Quarter 3
0303	Avoidance of	diagnosis of prostate cancer, at low	Medical	Quarter 5
	Overuse Measure	risk of recurrence, receiving	Association -	
	– Isotope Bone	interstitial prostate brachytherapy, OR	Physician	
	•		Consortium for	
	Scan for Staging	external beam radiotherapy to the		
	Low-Risk Patients	prostate, OR radical prostatectomy,	Performance	
		OR cryotherapy who did not have a	Improvement	
		bone scan performed at any time		
0200		since diagnosis of prostate cancer	A	0
0390	Prostate Cancer:	Percentage of patients with a	American	Quarter 3
	Adjuvant	diagnosis of prostate cancer, at high	Medical	
	Hormonal	risk of recurrence, receiving external	Association -	
	Therapy for High-	beam radiotherapy to the prostate	Physician	
	Risk Patients	who were prescribed adjuvant	Consortium for	
		hormonal therapy (GnRH agonist or	Performance	
		antagonist)	Improvement	
0385	Oncology:	Percentage of patients aged 18 years	American	Quarter 3
	Chemotherapy	and older with Stage IIIA through IIIC	Medical	
	for Stage IIIA	colon cancer who are prescribed or	Association -	
	through IIIC	who have received adjuvant	Physician	
	Colon Cancer	chemotherapy within the 12 month	Consortium for	
	Patients	reporting period	Performance	
			Improvement	
0378	Documentation	Percentage of patients aged 18 years	American	Quarter 3
	of Iron Stores in	and older with a diagnosis of MDS	Medical	
	Patients	who are receiving erythropoietin	Association -	
	Receiving	therapy with documentation of iron	Physician	
	Erythropoietin	stores prior to initiating erythropoietin	, Consortium for	
	Therapy	therapy	Performance	

Measure Number	Title	Description	Steward	Anticipated Review
0380	Multiple Myeloma – Treatment with Bisphosphonates	Percentage of patients aged 18 years and older with a diagnosis of multiple myeloma, not in remission, who were prescribed or received intravenous bisphosphonates within the 12 month reporting period	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0386	Oncology: Cancer Stage Documented Cancer stage documented	Percentage of patients with a diagnosis of breast, colon, or rectal cancer seen in the ambulatory setting who have a baseline AJCC cancer stage or documentation that the cancer is metastatic in the medical record at least once during the 12 month reporting period	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0391	Breast Cancer Resection Pathology Reporting- pT category (primary tumor) and pN category (regional lymph nodes) with histologic grade	Percentage of breast cancer resection pathology reports that include the pT category (primary tumor), the pN category (regional lymph nodes) and the histologic grade	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0392	Colorectal Cancer Resection Pathology Reporting- pT category (primary tumor) and pN category (regional lymph nodes) with histologic grade	Percentage of colon and rectum cancer resection pathology reports that include the pT category (primary tumor), the pN category (regional lymph nodes) and the histologic grade	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0455	Recording of Clinical Stage for Lung Cancer and Esophageal Cancer Resection	Percentage of all surgical patients undergoing treatment procedures for lung or esophageal cancer that have clinical TNM staging provided	Society of Thoracic Surgeons	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0457	Recording of Performance Status (Zubrod, Karnofsky, WHO or ECOG Performance Status) Prior to Lung or Esophageal Cancer Resection	Percentage of patients undergoing resection of a lung or esophageal cancer who had their performance status recorded within two weeks of the surgery date	Society of Thoracic Surgeons	Quarter 3
0459	Risk-Adjusted Morbidity after Lobectomy for Lung cancer	Percentage of patients undergoing elective lobectomy for lung cancer that have a prolonged length of stay (>14 days)	Society of Thoracic Surgeons	Quarter 3
0460	Risk-adjusted morbidity for esophagectomy for cancer	The percentage of patients undergoing elective esophagectomy for cancer that had a prolonged length of stay (>14 days)	Society of Thoracic Surgeons	Quarter 3
0508	Inappropriate use of "probably benign" assessment category in mammography screening	Percentage of final reports for screening mammograms that are classified as "probably benign"	American Medical Association - Physician Consortium for Performance Improvement	Quarter 4
0509	Reminder system for mammograms	Percentage of patients aged 40 years and older undergoing a screening mammogram whose information is entered into a reminder system* with a target due date for the next mammogram	American Medical Association - Physician Consortium for Performance Improvement	Quarter 4
BONE/JO	INT			
0423	Functional status change for patients with hip impairments	Percentage of patients aged 18 or older with a hip impairment associated with a functional deficit that had their functional status assessed at the beginning and end of rehabilitation.	Focus on Therapeutic Outcomes, Inc	Quarter 3
0422	Functional status change for patients with knee impairments	Functional status change in patients aged 18 or older with a knee impairment associated with a functional deficit that had their functional status assessed at the beginning and end of rehabilitation	Focus on Therapeutic Outcomes, Inc	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0425	Functional status change for patients with lumbar spine impairments	Percentage of patients aged 18 or older with a lumbar spine impairment associated with a functional deficit that had their functional status assessed at the beginning and end of rehabilitation.	Focus on Therapeutic Outcomes, Inc	Quarter 3
0426	Functional status change for patients with shoulder impairments	Percentage of patients aged 18 or older with a shoulder impairment associated with a functional deficit that had their functional status assessed at the beginning and end of rehabilitation.	Focus on Therapeutic Outcomes, Inc	Quarter 3
0427	Functional status change for patients with elbow, wrist or hand impairments	Percentage of patients aged 18 or older with an elbow, wrist or hand impairment associated with a functional deficit that had their functional status assessed at the beginning and end of rehabilitation.	Focus on Therapeutic Outcomes, Inc	Quarter 3
0424	Functional status change for patients with foot/ankle impairments	Functional status change in patients aged 18 or older with a foot/ankle impairment associated with a functional deficit that had their functional status assessed at the beginning and end of rehabilitation	Focus on Therapeutic Outcomes, Inc	Quarter 3
0428	Functional status change for patients with general orthopedic impairments	Functional status change in patients aged 18 or older with a general orthopedic impairment associated with a functional deficit that had their functional status assessed at the beginning and end of rehabilitation	Focus on Therapeutic Outcomes, Inc	Quarter 3
0511	Correlation With Existing Imaging Studies for All Patients Undergoing Bone Scintigraphy	Percentage of final reports for all patients, regardless of age, undergoing bone scintigraphy that include physician documentation of correlation with existing relevant imaging studies (e.g., x-ray, MRI, CT) that were performed	American Medical Association - Physician Consortium for Performance Improvement	Quarter 4
INFECTIO	DUS DISEASE			
0394	Hepatitis C: Counseling Regarding Use of Contraception Prior to Antiviral Treatment	Percentage of female patients aged 18 to 44 years and all men aged 18 years and older with a diagnosis chronic hepatitis C who are receiving antiviral treatment who were counseled regarding contraception prior to the initiation of antiviral treatment	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0401	Hepatitis C: Counseling Regarding Risk of Alcohol Consumption	Percentage of patients aged 18 years and older with a diagnosis of hepatitis C who received counseling regarding the risk of alcohol consumption at least once within the 12 month reporting period	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0395	Paired Measure: Hepatitis C RNA Testing Before Initiating Treatment (paired with 0396)	Percentage of patients aged 18 years and older with a diagnosis of chronic hepatitis C who are receiving antiviral treatment for whom quantitative HCV RNA testing was performed within 6 months prior to initiation of antiviral treatment	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0396	Paired Measure: HCV Genotype Testing Prior to Treatment (paired with 0395)	Percentage of patients aged 18 years and older with a diagnosis of chronic hepatitis C who are receiving antiviral treatment for whom HCV genotype testing was performed within 6 months prior to initiation of antiviral treatment	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0397	Hepatitis C: Prescribed Antiviral Therapy	Percentage of patients aged 18 years and older with a diagnosis of chronic hepatitis C who were prescribed peginterferon and ribavirin therapy within the 12 month reporting period	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0398	Hepatitis C: HCV RNA Testing at Week 12 of Treatment	Percentage of patients aged 18 years and older with a diagnosis of chronic hepatitis C who are receiving antiviral treatment for whom quantitative HCV RNA testing was performed at 12 weeks from initiation of antiviral treatment	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0403	Medical visit	Percentage of patients, regardless of age, with a diagnosis of HIV/AIDS with at least one medical visit in each 6 month period with a minimum of 60 days between each visit	National Committee for Quality Assurance	Quarter 3
0404	CD4+ Cell Count	Percentage of patients, regardless of age, with a diagnosis of HIV/AIDS with a CD4+ cell count or CD4+ cell percentage performed at least once every 6 months	National Committee for Quality Assurance	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0405	PCP prophylaxis	Percentage of patients aged 1 month or older who were prescribed Pneumocystis jiroveci pneumonia (PCP) prophylaxis.	National Committee for Quality Assurance	Quarter 3
0406	Adolescent and adult clients with AIDS who are prescribed potent ART (PROCESS MEASURE)	Percentage of patients who were prescribed potent antiretroviral therapy	National Committee for Quality Assurance	Quarter 3
0407	HIV RNA control after six months of potent antiretroviral therapy	Percentage of patients with viral load below limits of quantification OR patients with viral load not below limits of quantification who have a documented plan of care	National Committee for Quality Assurance	Quarter 3
0408	TB Screening	Percentage of patients for whom there was documentation that a tuberculosis (TB) screening test was placed and read at least once since the diagnosis of HIV infection.	National Committee for Quality Assurance	Quarter 3
0409	STD - Chlamydia and Gonorrhea Screenings	Percentage of patients screened for chlamydia and gonorrhea at least once since the diagnosis of HIV infection	National Committee for Quality Assurance	Quarter 3
0410	STD - Syphilis Screening	Percentage of patients aged 13 years and older with a diagnosis of HIV/AIDS for whom syphilis screening was performed in the last 12 months	National Committee for Quality Assurance	Quarter 3
0411	Hepatitis B Screening	Percentage of patients for whom Hepatitis B screening was performed at least once since the diagnosis of HIV infection or for whom there is documented immunity	National Committee for Quality Assurance	Quarter 3
0412	Hepatitis B Vaccination	Percentage of patients, regardless of age, with a diagnosis of HIV/AIDS who have received at least one hepatitis B vaccination, or who have documented immunity	National Committee for Quality Assurance	Quarter 3
0413	Screening for High Risk Sexual Behaviors	Percentage of patients, aged 13 years and older, who were screened* at least once in a 12-month measurement period for high risk sexual behavior	National Committee for Quality Assurance	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0431	Influenza Vaccination Coverage among Healthcare Personnel	Percentage of healthcare personnel (HCP) who receive the influenza vaccination.	Centers for Disease Control and Prevention	Quarter 3
0414	HIV/AIDS: Hepatitis C	Percentage of patients for whom Hepatitis C (Hep C) screening was performed at least once since the diagnosis of HIV infection or for whom there is documented immunity	National Committee for Quality Assurance	Quarter 3
0415	Screening for Injection Drug Use	Percentage of patients, aged 13 years and older, who were screened, at least once in a 12-month measurement period, for injection drug use	National Committee for Quality Assurance, American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0399	Paired Measure: Hepatitis C: Hepatitis A Vaccination (paired with 0400)	Percentage of patients aged 18 years and older with a diagnosis of hepatitis C who have received hepatitis A vaccination, or who have documented immunity	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0400	Paired Measure: Hepatitis C: Hepatitis B Vaccination (paired with 0399)	Percentage of patients aged 18 years and older with a diagnosis of hepatitis C who have received hepatitis B vaccination, or who have documented immunity.	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0393	Hepatitis C: Testing for Chronic Hepatitis C – Confirmation of Hepatitis C Viremia	Percentage of patients aged 18 years and older with a diagnosis of hepatitis C seen for an initial evaluation who had HCV RNA testing ordered or previously performed	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0402	Screening foreign-born adults for chronic hepatitis B	Percentage of adults aged 18 years and above born in an HBV-endemic country and tested for hepatitis B surface antigen and antibody	Asian Liver Center at Stanford University	Quarter 3
0475	Measurement of Hepatitis B Vaccine Administration to All Newborns Prior to Hospital or Birthing Facility Discharge	Percentage of newborns administered hepatitis B vaccine prior to discharge from the birthing facility or hospital, subtract the number of newborns who died prior to discharge, and divide this number by the number of live newborns discharged from the birthing facility or hospital during a given time period (perhaps annually) to identify the hepatitis B vaccine coverage rate for newborns at a single birthing facility or hospital.	Centers for Disease Control and Prevention	Quarter 4
MENTAL	HEALTH			
0418	Screening for Clinical Depression	Percentage of patients aged 18 years and older screened for clinical depression using a standardized tool and follow up plan documented.	Centers for Medicare & Medicaid Services	Quarter 3
CHILD HE				
0474	Birth Trauma Rate: Injury to Neonates (PSI #17)	Percentage of neonates with specific birth trauma per 1000 births. Exclude infants with injury to skeleton and osteogenesis imperfecta, subdural or cerebral hemorrhage in preterm infant.	Agency for Healthcare Research and Quality, National Perinatal Information Center	Quarter 4
0478	Nosocomial Blood Stream Infections in Neonates (NQI #3)	Percentage of qualifying neonates with selected bacterial blood stream infections	Agency for Healthcare Research and Quality	Quarter 4
0504	Pediatric Weight Documented in Kilograms	Percent of emergency department patients < 18 years of age with a current weight in kilograms documented in the ED record	American Academy of Pediatrics	Quarter 4
PERINAT	AL	·	·	·
0473	Appropriate DVT prophylaxis in women undergoing cesarean delivery	Measure adherence to current ACOG, ACCP recommendations for use of DVT prophylaxis in women undergoing cesarean delivery	Hospital Corporation of America	Quarter 4

Measure Number	Title	Description	Steward	Anticipated Review
0469	Elective delivery prior to 39 completed weeks gestation	Percentage of babies electively delivered prior to 39 completed weeks gestation	Hospital Corporation of America	Quarter 4
0470	Incidence of Episiotomy	Percentage of vaginal deliveries during which an episiotomy is performed	Christiana Care Health System	Quarter 4
OTHER	·			
0454	Anesthesiology and Critical Care: Perioperative Temperature Management	Percentage of patients, regardless of age, undergoing surgical or therapeutic procedures under general or neuraxial anesthesia of 60 minutes duration or longer for whom either active warming was used intraoperatively for the purpose of maintaining normothermia, OR at least one body temperature equal to or greater than 36 degrees Centigrade (or 96.8 degrees Fahrenheit) was recorded within the 30 minutes immediately before or the 30 minutes immediately after anesthesia end time	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0419	Universal Documentation and Verification of Current Medications in the Medical Record	Percentage of patients aged 18 years and older with a list of current medications with dosages (includes prescription, over-the-counter, herbals, vitamin/mineral/dietary [nutritional] supplements) and verified with the patient or authorized representative documented by the provider.	Centers for Medicare & Medicaid Services	Quarter 3
0464	Anesthesiology and Critical Care: Prevention of Catheter-Related Bloodstream Infections (CRBSI) – Central Venous Catheter (CVC) Insertion Protocol	Percentage of patients who undergo CVC insertion for whom CVC was inserted with all elements of maximal sterile barrier technique (cap AND mask AND sterile gown AND sterile gloves AND a large sterile sheet AND hand hygiene AND 2% chlorhexidine for cutaneous antisepsis) followed	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0450	Postoperative DVT or PE (PSI 12)	Percent of adult surgical discharges with a secondary diagnosis code of deep vein thrombosis or pulmonary embolism	Agency for Healthcare Research and Quality	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0271	Discontinuation of Prophylactic Antibiotics (Non- Cardiac Procedures)	Percentage of non-cardiac surgical patients aged 18 years and older undergoing procedures with the indications for prophylactic antibiotics AND who received a prophylactic antibiotic, who have an order for discontinuation of prophylactic antibiotics within 24 hours of surgical end time	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0268	Selection of Prophylactic Antibiotic: First OR Second Generation Cephalosporin	Percentage of surgical patients aged 18 years and older undergoing procedures with the indications for a first OR second generation cephalosporin prophylactic antibiotic, who had an order for cefazolin OR cefuroxime for antimicrobial prophylaxis	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0420	Pain Assessment Prior to Initiation of Patient Therapy	Percentage of patients with documentation of a pain assessment (if pain is present, including location, intensity and description) through discussion with the patient including the use of a standardized tool on each initial evaluation prior to initiation of therapy and documentation of a follow up plan.	Centers for Medicare & Medicaid Services	Quarter 3
0270	Timing of Antibiotic Prophylaxis: Ordering Physician	Percentage of surgical patients aged 18 years and older undergoing procedures with the indications for prophylactic parenteral antibiotics, who have an order for prophylactic antibiotic to be given within one hour (if fluoroquinolone or vancomycin, two hours), prior to the surgical incision (or start of procedure when no incision is required)	American Medical Association - Physician Consortium for Performance Improvement	Quarter 3
0486	Adoption of Medication e- Prescribing	Documents whether provider has adopted a qualified e-Prescribing system and the extent of use in the ambulatory setting.	Centers for Medicare & Medicaid Services	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0488	Adoption of Health Information Technology	Documents whether provider has adopted and is using health information technology. To qualify, the provider must have adopted and be using a certified/qualified electronic health record (EHR).	Centers for Medicare & Medicaid Services	Quarter 3
0490	The Ability to use Health Information Technology to Perform Care Management at the Point of Care	Documents the extent to which a provider uses a certified/qualified electronic health record (EHR) system capable of enhancing care management at the point of care. To qualify, the facility must have implemented processes within their EHR for disease management that incorporate the principles of care management at the point of care which include: a. The ability to identify specific patients by diagnosis or medication use b. The capacity to present alerts to the clinician for disease management, preventive services and wellness c. The ability to provide support for standard care plans, practice guidelines, and protocol	Centers for Medicare & Medicaid Services	Quarter 3
0489	The Ability for Providers with HIT to Receive Laboratory Data Electronically Directly into their Qualified/Certifie d EHR System as Discrete Searchable Data Elements	Documents the extent to which a provider uses certified/qualified electronic health record (EHR) system that incorporates an electronic data interchange with one or more laboratories allowing for direct electronic transmission of laboratory data into the EHR as discrete searchable data elements.	Centers for Medicare & Medicaid Services	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0421	Adult Weight Screening and Follow-Up	Percentage of patients aged 18 years and older with a calculated BMI documented in the medical record AND if the most recent BMI is outside the parameters, a follow up plan is documented. Parameters: age 65 and older BMI > or = 30 or < 22; age 18-64 BMI > or = 25 or< 18.5	Centers for Medicare & Medicaid Services	Quarter 3
0491	Tracking of Clinical Results Between Visits	Documentation of the extent to which a provider uses a certified/qualified electronic health record (EHR) system to track pending laboratory tests, diagnostic studies (including common preventive screenings) or patient referrals. The Electronic Health Record includes provider reminders when clinical results are not received within a predefined timeframe.	Centers for Medicare & Medicaid Services	Quarter 3
0487	EHR with EDI prescribing used in encounters where a prescribing event occurred.	Of all patient encounters within the past month that used an electronic health record (EHR) with electronic data interchange (EDI) where a prescribing event occurred, how many used EDI for the prescribing event.	City of New York Department of Health and Mental Hygiene	Quarter 3
0492	Participation in a Practice-based or individual Quality Database Registry with a standard measure set.	This Registry should be capable of a. generating population based reports relating to published guideline goals or benchmarking data b. providing comparisons to the practitioner c. providing feedback that is related to guideline goals d. capturing data for one or more chronic disease conditions (i.e. diabetes) or preventive care measures (i.e. USPTF recommendations) for all patients eligible for the measures	Centers for Medicare & Medicaid Services	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0493	Participation by a physician or other clinician in systematic clinical database registry that includes consensus endorsed quality measures	Participation in a systematic qualified clinical database registry involves: a. Physician or other clinician submits standardized data elements to registry b. Data elements are applicable to consensus endorsed quality measures c. Registry measures shall include at least two (2) representative NQF consensus endorsed measures for registry's clinical topic(s) and report on all patients eligible for the selected measures. d. Registry provides calculated measures results, benchmarking, and quality improvement information to individual physicians and clinicians. e. Registry must receive data from more than 5 separate practices and may not be located (warehoused) at an individual group's practice. Participation in a national or state- wide registry is encouraged for this measure. f. Registry may provide feedback directly to the provider's local registry if one exists.	Centers for Medicare & Medicaid Services	Quarter 3
0429	Change in Basic Mobility as Measured by the AM-PAC:	The Activity Measure for Post Acute Care (AM-PAC) is a functional status assessment instrument developed specifically for use in facility and community dwelling post acute care (PAC) patients: Basic Mobility scale	CREcare	Quarter 3
0430	Change in Daily Activity Function as Measured by the AM-PAC:	The Activity Measure for Post Acute Care (AM-PAC) is a functional status assessment instrument developed specifically for use in facility and community dwelling post acute care (PAC) patients: Daily Activity scale	CREcare	Quarter 3

Measure Number	Title	Description	Steward	Anticipated Review
0453	Urinary catheter removed on Postoperative Day 1 (POD1) or Postoperative Day 2 (POD2) with day of surgery being day zero.	Surgical patients with urinary catheter removed on Postoperative Day 1 or Postoperative Day 2 with day of surgery being day zero.	Center for Medicare & Medicaid Services	Quarter 3
0452	Surgery Patients with Perioperative Temperature Management	Surgery patients for whom either active warming was used intraoperatively for the purpose of maintaining normothermia or who had at least one body temperature equal to or greater than 96.8° F/36° C recorded within the 30 minutes immediately prior to or the 15 minutes immediately after Anesthesia End Time.	Center for Medicare & Medicaid Services	Quarter 3
0500	Severe Sepsis and Septic Shock: Management Bundle	Initial steps in the management of the patient presenting with infection (severe sepsis or septic shock)	Henry Ford Hospital	Quarter 4
0502	Pregnancy test for female abdominal pain patients.	Pregnancy test for female abdominal pain patients.	American College of Emergency Physicians	Quarter 4
0510	Exposure time reported for procedures using fluoroscopy	Percentage of final reports for procedures using fluoroscopy that include documentation of radiation exposure or exposure time	American Medical Association - Physician Consortium for Performance Improvement	Quarter 4
0495	Median Time from ED Arrival to ED Departure for Admitted ED Patients	Median time from emergency department arrival to time of departure from the emergency room for patients admitted to the facility from the emergency department	Centers for Medicare & Medicaid Services	Quarter 4