

**Behavioral Health Endorsement Maintenance Phase II: NQF-Endorsed® Maintenance Standards**

Updated on September 19th, 2012

<b>Measure Number</b>	<b>Title</b>	<b>Description</b>	<b>Measure Steward</b>
0008	Experience of Care and Health Outcomes (ECHO) Survey	52- questions including patient demographic information. The survey measures patient experiences with behavioral health care (mental health and substance abuse treatment) and the organization that provides or manages the treatment and health outcomes. Level of analysis: health plan- HMO, PPO, Medicare, Medicaid, commercial	Agency for Healthcare Research and Quality
0103	Major Depressive Disorder (MDD): Diagnostic Evaluation	Percentage of patients aged 18 years and older with a new diagnosis or recurrent episode of MDD who met the DSM-IV criteria during the visit in which the new diagnosis or recurrent episode was identified during the measurement period	American Medical Association - Physician Consortium for Performance Improvement (AMA-PCPI)
0104	Major Depressive Disorder (MDD): Suicide Risk Assessment	Percentage of patients aged 18 years and older with a new diagnosis or recurrent episode of MDD who had a suicide risk assessment completed at each visit during the measurement period	American Medical Association - Physician Consortium for Performance Improvement (AMA-PCPI)
0105	Antidepressant Medication Management	The percentage of members 18 years of age and older who were diagnosed with a new episode of major depression and treated with antidepressant medication, and who remained on an antidepressant medication treatment. Two rates are reported. a) Effective Acute Phase Treatment. The percentage of newly diagnosed and treated members who remained on an antidepressant medication for at least 84 days (12 weeks). b) Effective Continuation Phase Treatment. The percentage of newly diagnosed and treated members who remained on an antidepressant medication for at least 180 days (6 months).	National Committee for Quality Assurance

**Behavioral Health Endorsement Maintenance Phase II: NQF-Endorsed® Maintenance Standards**

Updated on September 19th, 2012

Measure Number	Title	Description	Measure Steward
0106	Diagnosis of attention deficit hyperactivity disorder (ADHD) in primary care for school age children and adolescents	Percentage of patients newly diagnosed with attention deficit hyperactivity disorder (ADHD) whose medical record contains documentation of Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) or Diagnostic and Statistical Manual for Primary Care (DSM-PC) criteria being addressed.	Institute for Clinical Systems Improvement
0107	Management of attention deficit hyperactivity disorder (ADHD) in primary care for school age children and adolescents	Percentage of patients treated with psychostimulant medication for the diagnosis of attention deficit hyperactivity disorder (ADHD) whose medical record contains documentation of a follow-up visit at least twice a year.	Institute for Clinical Systems Improvement
0108	Follow-Up Care for Children Prescribed ADHD Medication (ADD)	<p>The percentage of children newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed. Two rates are reported.</p> <ul style="list-style-type: none"> <li>• Initiation Phase. The percentage of members 6–12 years of age as of the IPSD with an ambulatory prescription dispensed for ADHD medication, who had one follow-up visit with practitioner with prescribing authority during the 30-day Initiation Phase.</li> <li>• Continuation and Maintenance (C&amp;M) Phase. The percentage of members 6–12 years of age as of the IPSD with an ambulatory prescription dispensed for ADHD medication, who remained on the medication for at least 210 days and who, in addition to the visit in the Initiation Phase, had at least two follow-up visits with a practitioner within 270 days (9 months) after the Initiation Phase ended.</li> </ul>	National Committee for Quality Assurance

**Behavioral Health Endorsement Maintenance Phase II: NQF-Endorsed® Maintenance Standards**

Updated on September 19th, 2012

<b>Measure Number</b>	<b>Title</b>	<b>Description</b>	<b>Measure Steward</b>
0418	Screening for Clinical Depression	Percentage of patients aged 18 years and older screened for clinical depression using a standardized tool and follow up plan documented.	Centers for Medicare and Medicaid Services
0518	Depression Assessment Conducted	Percent of patients who were screened for depression (using a standardized depression screening tool) at start or resumption of home health care	Centers for Medicare and Medicaid Services
0552	HBIPS-4: Patients discharged on multiple antipsychotic medications.	Patients discharged on multiple antipsychotic medications overall and stratified by age groups: Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years). Note: this is a paired measure with HBIPS-5: Patients discharged on multiple antipsychotic medications with appropriate justification.	The Joint Commission
0557	HBIPS-6 Post discharge continuing care plan created	Patients discharged from a hospital-based inpatient psychiatric setting with a continuing care plan created overall and stratified by age groups: Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years). Note: this is a paired measure with HBIPS-7: Post discharge continuing care plan transmitted to next level of care provider upon discharge.	The Joint Commission
0558	HBIPS-7 Post discharge continuing care plan transmitted to next level of care provider upon discharge	Patients discharged from a hospital-based inpatient psychiatric setting with a continuing care plan provided to the next level of care clinician or entity overall and stratified by age groups: Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years). Note: this is a paired measure with HBIPS-6: Post discharge continuing care plan created.	The Joint Commission

**Behavioral Health Endorsement Maintenance Phase II: NQF-Endorsed® Maintenance Standards**

Updated on September 19th, 2012

Measure Number	Title	Description	Measure Steward
0560	HBIPS-5 Patients discharged on multiple antipsychotic medications with appropriate justification	Patients discharged from a hospital-based inpatient psychiatric setting on two or more antipsychotic medications with appropriate justification overall and stratified by age groups: Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years). Note: this is a paired measure with HBIPS-4: Patients discharged on multiple antipsychotic medications.	The Joint Commission
0640	HBIPS-2 Hours of physical restraint use	The number of hours that all patients admitted to a hospital-based inpatient psychiatric setting were maintained in physical restraint per 1000 psychiatric inpatient hours, overall and stratified by age groups: : Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years).	The Joint Commission
0641	HBIPS-3 Hours of seclusion use	The number of hours that all patients admitted to a hospital-based inpatient psychiatric setting were held in seclusion per 1000 psychiatric inpatient hours, overall and stratified by age groups: Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years).	The Joint Commission