

NQF Behavioral Health Project
Phase II Submitted Measures

Measure #	Title	Description	Developer
0103	Major Depressive Disorder (MDD): Diagnostic Evaluation	Percentage of patients aged 18 years and older with a new diagnosis or recurrent episode of MDD who met the DSM-IV criteria during the visit in which the new diagnosis or recurrent episode was identified during the measurement period	AMA-PCPI
0104	Major Depressive Disorder (MDD): Suicide Risk Assessment	Percentage of patients aged 18 years and older with a new diagnosis or recurrent episode of MDD who had a suicide risk assessment completed at each visit during the measurement period	AMA-PCPI
0105	Antidepressant Medication Management	The percentage of members 18 years of age and older who were diagnosed with a new episode of major depression and treated with antidepressant medication, and who remained on an antidepressant medication treatment. Two rates are reported. a) Effective Acute Phase Treatment. The percentage of newly diagnosed and treated members who remained on an antidepressant medication for at least 84 days (12 weeks). b) Effective Continuation Phase Treatment. The percentage of newly diagnosed and treated members who remained on an antidepressant medication for at least 180 days (6 months).	NCQA
0418	Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan	Percentage of patients aged 12 years and older screened for clinical depression on the date of encounter using an age appropriate standardized depression screening tool AND, if positive, a follow-up plan is documented on the date of the positive screen.	CMS / Quality Insights of PA
0518	Depression Assessment Conducted	Percent of patients who were screened for depression (using a standardized depression screening tool) at start or resumption of home health care	CMS/Acumen

NQF Behavioral Health Project
Phase II Submitted Measures

Measure #	Title	Description	Developer
0552	HBIPS-4 Patients discharged on multiple antipsychotic medications	Patients discharged on multiple antipsychotic medications overall and stratified by age groups: Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years). Note: this is a paired measure with HBIPS-5: Patients discharged on multiple antipsychotic medications with appropriate justification.	The Joint Commission
0557	HBIPS-6 Post discharge continuing care plan created	Patients discharged from a hospital-based inpatient psychiatric setting with a continuing care plan created overall and stratified by age groups: Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years). Note: this is a paired measure with HBIPS-7: Post discharge continuing care plan transmitted to next level of care provider upon discharge.	The Joint Commission
0558	HBIPS-7 Post discharge continuing care plan transmitted to next level of care provider upon discharge	Patients discharged from a hospital-based inpatient psychiatric setting with a continuing care plan provided to the next level of care clinician or entity overall and stratified by age groups: Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years). Note: this is a paired measure with HBIPS-6: Post discharge continuing care plan created.	The Joint Commission
0560	HBIPS-5 Patients discharged on multiple antipsychotic medications with appropriate justification	Patients discharged from a hospital-based inpatient psychiatric setting on two or more antipsychotic medications with appropriate justification overall and stratified by age groups: Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years). Note: this is a paired measure with HBIPS-4: Patients discharged on multiple antipsychotic medications.	The Joint Commission

NQF Behavioral Health Project
Phase II Submitted Measures

Measure #	Title	Description	Developer
0640	HBIPS-2 Hours of physical restraint use	The number of hours that all patients admitted to a hospital-based inpatient psychiatric setting were maintained in physical restraint per 1000 psychiatric inpatient hours, overall and stratified by age groups: : Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years).	The Joint Commision
0641	HBIPS-3 Hours of seclusion use	The number of hours that all patients admitted to a hospital-based inpatient psychiatric setting were held in seclusion per 1000 psychiatric inpatient hours, overall and stratified by age groups: Children (Age 1 through 12 years), Adolescents (Age 13 through 17 years), Adults (Age 18 through 64 years), Older Adults (Age greater than and equal to 65 years).	The Joint Commision
1651	TOB-1 Tobacco Use Screening	Hospitalized patients age 18 years and older who are screened during the hospital stay for tobacco use (cigarettes, smokeless tobacco, pipe and cigars) within the past 30 days. This measure is intended to be used as part of a set of 4 linked measures addressing Tobacco Use (TOB-2 Tobacco Use Treatment Provided or Offered (during the hospital stay); TOB-3 Tobacco Use Treatment Provided or offered at Discharge; TOB-4 Tobacco Use: Assessing Status After Discharge.)	The Joint Commision
1654	TOB - 2 Tobacco Use Treatment Provided or Offered and the subset measure TOB-2a Tobacco Use Treatment	The measure is reported as an overall rate which includes all hospitalized patients 18 years of age and older to whom tobacco use treatment was provided during the hospital stay, or offered and refused, and a second rate, a subset of the first, which includes only those patients who received tobacco use treatment during the hospital stay. Refer to section 2a1.10 Stratification Details/Variables for the rationale for the addition of the subset measure. These measures are intended to be used as part of a set of 4 linked measures addressing Tobacco Use (TOB-1 Tobacco Use Screening; TOB-3 Tobacco Use Treatment Provided or Offered at Discharge; TOB-4 Tobacco Use: Assessing Status After Discharge.)	The Joint Commision

NQF Behavioral Health Project
Phase II Submitted Measures

Measure #	Title	Description	Developer
1656	TOB-3 Tobacco Use Treatment Provided or Offered at Discharge and the subset measure TOB-3a Tobacco Use Treatment at Discharge	The measure is reported as an overall rate which includes all hospitalized patients 18 years of age and older to whom tobacco use treatment was provided, or offered and refused, at the time of hospital discharge, and a second rate, a subset of the first, which includes only those patients who received tobacco use treatment at discharge. Treatment at discharge includes a referral to outpatient counseling and a prescription for one of the FDA-approved tobacco cessation medications. Refer to section 2a1.10 Stratification Details/Variables for the rationale for the addition of the subset measure. These measures are intended to be used as part of a set of 4 linked measures addressing Tobacco Use (TOB-1 Tobacco Use Screening; TOB 2 Tobacco Use Treatment Provided or Offered During the Hospital Stay; TOB-4 Tobacco Use: Assessing Status After Discharge).	The Joint Commission
1657	TOB-4 Tobacco Use: Assessing Status after Discharge	Hospitalized patients 18 years of age and older who are identified through the screening process as having used tobacco products (cigarettes, smokeless tobacco, pipe, and cigars) within the past 30 days who are contacted between 15 and 30 days after hospital discharge and follow-up information regarding tobacco use status is collected. This measure is intended to be used as part of a set of 4 linked measures addressing Tobacco Use (TOB-1 Tobacco Use Screening; TOB-2 Tobacco Use Treatment Provided or Offered (during hospital stay); TOB-3 Tobacco Use Treatment Provided or Offered at Discharge).	The Joint Commission
1661	SUB-1 Alcohol Use Screening	Hospitalized patients 18 years of age and older who are screened during the hospital stay using a validated screening questionnaire for unhealthy alcohol use. This measure is intended to be used as part of a set of 4 linked measures addressing Substance Use (SUB-1 Alcohol Use Screening ; SUB-2 Alcohol Use Brief Intervention Provided or Offered; SUB-3 Alcohol and Other Drug Use Disorder Treatment Provided or Offered at Discharge; SUB-4 Alcohol and Drug Use: Assessing Status after Discharge).	The Joint Commission

NQF Behavioral Health Project
Phase II Submitted Measures

Measure #	Title	Description	Developer
1663	SUB-2 Alcohol Use Brief Intervention Provided or Offered and SUB-2a Alcohol Use Brief Intervention	<p>The measure is reported as an overall rate which includes all hospitalized patients 18 years of age and older to whom a brief intervention was provided, or offered and refused, and a second rate, a subset of the first, which includes only those patients who received a brief intervention. The Provided or Offered rate (SUB-2), describes patients who screened positive for unhealthy alcohol use who received or refused a brief intervention during the hospital stay. The Alcohol Use Brief Intervention (SUB-2a) rate describes only those who received the brief intervention during the hospital stay. Those who refused are not included.</p> <p>These measures are intended to be used as part of a set of 4 linked measures addressing Substance Use (SUB-1 Alcohol Use Screening ; SUB-2 Alcohol Use Brief Intervention Provided or Offered; SUB-3 Alcohol and Other Drug Use Disorder Treatment Provided or Offered at Discharge; SUB-4 Alcohol and Drug Use: Assessing Status after Discharge).</p>	The Joint Commision

NQF Behavioral Health Project
Phase II Submitted Measures

Measure #	Title	Description	Developer
1664	SUB-3 Alcohol & Other Drug Use Disorder Treatment Provided or Offered at Discharge and SUB-3a Alcohol & Other Drug Use Disorder Treatment at Discharge	<p>The measure is reported as an overall rate which includes all hospitalized patients 18 years of age and older to whom alcohol or drug use disorder treatment was provided, or offered and refused, at the time of hospital discharge, and a second rate, a subset of the first, which includes only those patients who received alcohol or drug use disorder treatment at discharge. The Provided or Offered rate (SUB-3) describes patients who are identified with alcohol or drug use disorder who receive or refuse at discharge a prescription for FDA-approved medications for alcohol or drug use disorder, OR who receive or refuse a referral for addictions treatment. The Alcohol and Other Drug Disorder Treatment at Discharge (SUB-3a) rate describes only those who receive a prescription for FDA-approved medications for alcohol or drug use disorder OR a referral for addictions treatment. Those who refused are not included.</p> <p>These measures are intended to be used as part of a set of 4 linked measures addressing Substance Use (SUB-1 Alcohol Use Screening ; SUB-2 Alcohol Use Brief Intervention Provided or Offered; SUB-3 Alcohol and Other Drug Use Disorder Treatment Provided or Offered at Discharge; SUB-4 Alcohol and Drug Use: Assessing Status after Discharge).</p>	The Joint Commision
1665	SUB-4 Alcohol & Drug Use: Assessing Status After Discharge	<p>Hospitalized patients age 18 years and older who screened positive for unhealthy alcohol use or who received a diagnosis of alcohol or drug disorder during their inpatient stay, who are contacted between 7 and 30 days after hospital discharge and follow-up information regarding their alcohol or drug use status post discharge is collected.</p> <p>This measure is intended to be used as part of a set of 4 linked measures addressing Substance Use (SUB-1) Alcohol Use Screening ; SUB-2 Alcohol Use Brief Intervention Provided or Offered; SUB-3 Alcohol and Other Drug Use Disorder Treatment Provided or Offered at Discharge; SUB-4 Alcohol and Drug Use: Assessing Status after Discharge).</p>	The Joint Commision

NQF Behavioral Health Project
Phase II Submitted Measures

Measure #	Title	Description	Developer
1880	Adherence to Mood Stabilizers for Individuals with Bipolar I Disorder	The measure calculates the percentage of individuals 18 years of age or greater as of the beginning of the measurement period with bipolar I disorder who are prescribed a mood stabilizer medication, with adherence to the mood stabilizer medication [defined as a Proportion of Days Covered (PDC)] of at least 0.8 during the measurement period (12 consecutive months).	CMS/ FMQAI
1884	Depression Response at Six Months- Progress Towards Remission	Adult patients age 18 and older with major depression or dysthymia and an initial PHQ-9 score > 9 who demonstrate a response to treatment at six months defined as a PHQ-9 score that is reduced by 50% or greater from the initial PHQ-9 score. This measure applies to both patients with newly diagnosed and existing depression identified during the defined measurement period whose current PHQ-9 score indicates a need for treatment.	MN Community Measurement
1885	Depression Response at Twelve Months - Progress Towards Remission	Adult patients age 18 and older with major depression or dysthymia and an initial PHQ-9 score > 9 who demonstrate a response to treatment at twelve months defined as a PHQ-9 score that is reduced by 50% or greater from the initial PHQ-9 score. This measure applies to both patients with newly diagnosed and existing depression identified during the defined measurement period whose current PHQ-9 score indicates a need for treatment.	MN Community Measurement
1922	HBIPS-1 Admission Screening	The proportion of patients admitted to a hospital-based inpatient psychiatric setting who are screened within the first three days of hospitalization for all of the following: risk of violence to self or others, substance use, psychological trauma history and patient strengths. This measure is a part of a set of seven nationally implemented measures that address hospital-based inpatient psychiatric services (HBIPS-2: Physical Restraint, HBIPS-3: Seclusion, HBIPS-4: Multiple Antipsychotic Medications at Discharge, HBIPS-5: Multiple Antipsychotic Medications at Discharge with Appropriate Justification, HBIPS-6: Post Discharge Continuing Care Plan and HBIPS-7: Post Discharge Continuing Care Plan Transmitted) that are used in The Joint Commission's accreditation process.	The Joint Commision

NQF Behavioral Health Project
Phase II Submitted Measures

Measure #	Title	Description	Developer
2152	Preventive Care and Screening: Unhealthy Alcohol Use: Screening & Brief Counseling	Percentage of patients aged 18 years and older who were screened for unhealthy alcohol use at least once during the two-year measurement period using a systematic screening method AND who received brief counseling if identified as an unhealthy alcohol user	AMA-PCPI