

	ASTHMA	ASTHMA/COPD	COPD	PNEUMONIA	IMAGING	CRITICAL CARE
NQF Portfolio	<ul style="list-style-type: none">• Asthma in Younger Adults Admission Rate (PQI 15) (NQF #0283)• Asthma Admission Rate (pediatric) (NQF #0728)• Asthma: Pharmacologic Therapy for Persistent Asthma (NQF #0047)• Asthma Medication Ratio (AMR) (NQF #1800)	<ul style="list-style-type: none">• Chronic Obstructive Pulmonary Disease (COPD) or Asthma in Older Adults Admission Rate (PQI 5) (NQF #0275)	<ul style="list-style-type: none">• COPD: spirometry evaluation (NQF #0091)• COPD: inhaled bronchodilator therapy (NQF #0102)• Use of Spirometry Testing in the Assessment and Diagnosis of COPD (NQF #0577)• Health-related Quality of Life in COPD patients before and after Pulmonary Rehabilitation (NQF #0700)• Functional Capacity in COPD patients before and after Pulmonary Rehabilitation (NQF #0701)• Hospital 30-Day, All-Cause, Risk-Standardized Readmission Rate (RSRR) following Chronic Obstructive Pulmonary Disease (COPD) Hospitalization (NQF #1891)• Hospital 30-Day, All-Cause, Risk-Standardized Mortality Rate (RSMR) following Chronic Obstructive Pulmonary Disease (COPD) Hospitalization (NQF #1893)• Pharmacotherapy Management of COPD Exacerbation (NQF #2856)	<ul style="list-style-type: none">• Pneumonia Mortality Rate (IQI #20) (NQF #0231)• Bacterial Pneumonia Admission Rate (PQI 11) (NQF #0279)• Hospital 30-day, all-cause, risk-standardized mortality rate (RSMR) following pneumonia hospitalization (NQF #0468)• Hospital 30-day, all-cause, risk-standardized readmission rate (RSRR) following pneumonia hospitalization (NQF #0506)• Proportion of Patients Hospitalized with Pneumonia that have a Potentially Avoidable Complication (during the Index Stay or in the 30-day Post-Discharge Period) (NQF #0708)• Hospital-level, risk-standardized payment associated with a 30-day episode of care for pneumonia (NQF #2579)	<ul style="list-style-type: none">• Thorax CT: Use of Contrast Material (NQF #0513)	<ul style="list-style-type: none">• PICU Severity-adjusted Length of Stay (NQF #0334)• PICU Unplanned Readmission Rate (NQF #0335)• Intensive Care Unit Venous Thromboembolism Prophylaxis (NQF #0372 and #3053)• Patients Admitted to ICU who Have Care Preferences Documented (NQF #1626)• Proportion of patients who died from cancer admitted to the ICU in the last 30 days of life (NQF #0213)
MIPS	<ul style="list-style-type: none">• Optimal Asthma Control ((NQF #2852-Not Endorsed)• Medication Management for People with Asthma (MMA) (NQF #1799-Endorsement Removed)	<ul style="list-style-type: none">• None	<ul style="list-style-type: none">• COPD: spirometry evaluation (NQF #0091)• COPD: inhaled bronchodilator therapy (NQF #0102)	<ul style="list-style-type: none">• None	<ul style="list-style-type: none">• None	<ul style="list-style-type: none">• None

Hospital Compare	<ul style="list-style-type: none">• None	<ul style="list-style-type: none">• None	<ul style="list-style-type: none">• Initial antibiotic selection for community-acquired pneumonia (CAP) in immunocompetent patients (NQF #0147- Endorsement Removed)• Hospital 30-Day, All-Cause, Risk-Standardized Readmission Rate (RSRR) following Chronic Obstructive Pulmonary Disease (COPD) Hospitalization (NQF #1891)• Hospital 30-Day, All-Cause, Risk-Standardized Mortality Rate (RSMR) following Chronic Obstructive Pulmonary Disease (COPD) Hospitalization (NQF #1893)	<ul style="list-style-type: none">• Hospital 30-day, all-cause, risk-standardized mortality rate (RSMR) following pneumonia hospitalization (NQF #0468)• Hospital 30-day, all-cause, risk-standardized readmission rate (RSRR) following pneumonia hospitalization (NQF #0506)• Hospital-level, risk-standardized payment associated with a 30-day episode of care for pneumonia (NQF #2579)	<ul style="list-style-type: none">• Thorax CT: Use of Contrast Material (NQF #0513)• 	<ul style="list-style-type: none">• None
Choosing Wisely	<ul style="list-style-type: none">• American Academy of Allergy, Asthma & Immunology: 1/10 for asthma, 1 related (but not exact) NQF measure<ul style="list-style-type: none">○ Don't diagnose or manage asthma without spirometry.	<ul style="list-style-type: none">• None	<ul style="list-style-type: none">• None	<ul style="list-style-type: none">• None	<ul style="list-style-type: none">• American College of Radiology: 1/5 for PCC Imaging, No NQF relevant measures<ul style="list-style-type: none">○ Avoid admission or preoperative chest x-rays for ambulatory patients with unremarkable history and physical exam.	<ul style="list-style-type: none">• None