



Table of Endocrine Project: NQF-Endorsed® Maintenance Standards

Measure Number	Title	Description	Measure Steward
0037	Osteoporosis testing in older women	Percentage of female patients aged 65 and older who reported receiving a bone density test (BMD) to check for osteoporosis	National Committee for Quality Assurance
0045	Osteoporosis: Communication with the Physician Managing Ongoing Care Post Fracture of Hip, Spine or Distal Radius for Men and Women Aged 50 Years and Older	The percentage of patients aged 50 years and older treated for a hip, spine or distal radial fracture with documentation of communication with the physician managing the patient's ongoing care that a fracture occurred and that the patient was or should be tested or treated for osteoporosis.	National Committee for Quality Assurance
0046	Osteoporosis: Screening or Therapy for Women Aged 65 Years and Older	Percentage of female patients aged 65 years and older who had a central dual-energy X-ray absorptiometry (DXA) measurement ordered or performed at least once since age 60 or pharmacologic therapy prescribed within 12 months.	National Committee for Quality Assurance
0048	Osteoporosis: Management Following Fracture of Hip, Spine or Distal Radius for Men and Women Aged 50 Years and Older	Percentage of patients aged 50 years or older with fracture of the hip, spine or distal radius that had a central DXA measurement ordered or performed or pharmacologic therapy prescribed	National Committee for Quality Assurance
0049	Osteoporosis: Pharmacologic Therapy for Men and Women Aged 50 Years and Older	Percentage of patients aged 50 years and older with a diagnosis of osteoporosis who were prescribed pharmacologic therapy within 12 months	National Committee for Quality Assurance
0053	Osteoporosis Management in Women Who Had a Fracture	The percentage of women 67 years of age and older who suffered a fracture and who had either a bone mineral density (BMD) test or prescription for a drug to treat or prevent osteoporosis in the six months after the date of fracture.	National Committee for Quality Assurance
0055	Comprehensive Diabetes Care: Eye Exam	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) who received a retinal or dilated eye exam during the measurement year or a negative retinal or dilated eye exam in the year prior to the measurement year.	National Committee for Quality Assurance

Measure Number	Title	Description	Measure Steward
0056	Diabetes: Foot exam	The percentage of patients 18-75 years of age with diabetes (type 1 and type 2) who received a foot exam (visual inspection with either a sensory exam or a pulse exam) during the measurement year.	National Committee for Quality Assurance
0057	Comprehensive Diabetes Care: Hemoglobin A1c (HbA1c) testing	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) who received an HbA1c test during the measurement year.	National Committee for Quality Assurance
0059	Comprehensive Diabetes Care: Hemoglobin A1c (HbA1c) Poor Control (>9.0%)	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) whose most recent HbA1c level during the measurement year was greater than 9.0% (poor control) or was missing a result, or if an HbA1c test was not done during the measurement year.	National Committee for Quality Assurance
0060	Hemoglobin A1c (HbA1c) Testing for Pediatric Patients	Percentage of pediatric patients aged 5-17 years of age with diabetes who received an HbA1c test during the measurement year.	National Committee for Quality Assurance
0061	Comprehensive Diabetes Care: Blood Pressure Control (<140/90 mm Hg)	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) whose most recent blood pressure (BP) reading is <140/90 mm Hg during the measurement year.	National Committee for Quality Assurance
0062	Comprehensive Diabetes Care: Medical Attention for Nephropathy	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) who received a nephropathy screening test or had evidence of nephropathy during the measurement year.	National Committee for Quality Assurance
0063	Comprehensive Diabetes Care: LDL-C Screening	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) who received an LDL-C test during the measurement year.	National Committee for Quality Assurance
0064	Comprehensive Diabetes Care: LDL-C Control <100 mg/dL	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) whose most recent LDL-C test is <100 mg/dL during the measurement year.	National Committee for Quality Assurance
0416	Diabetic Foot & Ankle Care, Ulcer Prevention – Evaluation of Footwear	Percentage of patients aged 18 years and older with a diagnosis of diabetes mellitus who were evaluated for proper footwear and sizing during one or more office visits within 12 month	American Podiatric Medical Association
0417	Diabetic Foot & Ankle Care, Peripheral Neuropathy – Neurological Evaluation	Percentage of patients, 18 years or older, with diabetes aged 18 years and older with who had a lower extremity neurological exam with risk categorization performed and a treat plan established at least once within 12 months a diagnosis of diabetes mellitus who had a neurological examination of their lower extremities during one or more office visits within 12 months	American Podiatric Medical Association

Measure Number	Title	Description	Measure Steward
0519	Diabetic Foot Care and Patient Education Implemented	Percent of diabetic patients for whom physician-ordered monitoring for the presence of skin lesions on the lower extremities and patient education on proper foot care were implemented during their episode of care	Centers for Medicare and Medicaid Services
0545	Adherence to Chronic Medications for Individuals with Diabetes Mellitus	<p>The measure addresses adherence to three types of chronic medications; statins, angiotensin converting enzyme inhibitors (ACEIs)/angiotensin receptor blockers (ARBs) and oral hypoglycemic agents. The measure is divided into three submeasures:</p> <p>Measure A: The percentage of eligible individuals who had at least two prescriptions for statins and who have a Proportion of Days Covered (PDC) of at least 0.8 during the measurement period (12 consecutive months).</p> <p>Measure B: The percentage of eligible individuals who had at least two prescriptions for ACEIs/ARBs and who have a PDC of at least 0.8 during the measurement period (12 consecutive months).</p> <p>Measure C: The percentage of eligible individuals who had at least two prescriptions for a single oral hypoglycemic agent or at least two prescriptions for multiple agents within an antidiabetic class and who have a PDC of at least 0.8 for at least 1 antidiabetic class during the measurement period (12 consecutive months).</p>	Centers for Medicare and Medicaid Services
0546	Diabetes: Appropriate Treatment of Hypertension	The percentage of patients who were dispensed a medication for diabetes and hypertension that are receiving an angiotensin-converting -enzyme-inhibitor (ACEI) or angiotensin receptor blocker (ARB) or direct renin inhibitor (DRI) renin-angiotensin-antagonist medication.	Pharmacy Quality Alliance
0575	Comprehensive Diabetes Care: Hemoglobin A1c (HbA1c) Control (<8.0%)	The percentage of members 18 - 75 years of age with diabetes (type 1 and type 2) whose most recent HbA1c level is <8.0% during the measurement year.	National Committee for Quality Assurance
0603	Adult(s) taking insulin with evidence of self-monitoring blood glucose testing.	This measure identifies patients with diabetes mellitus taking insulin that had evidence of self-monitoring blood glucose testing in last 12 reported months.	Optum
0604	Adult(s) with diabetes mellitus that had a serum creatinine in last 12 reported months.	This measure identifies adults with diabetes mellitus that had a serum creatinine test in last 12 reported months.	Optum
0614	Steroid Use - Osteoporosis Screening	The percentage of patients aged 18 years and older who have been on chronic steroids for at least 180 days and had a bone density evaluation or were taking osteoporosis treatment	ActiveHealth Management

Measure Number	Title	Description	Measure Steward
0729	Optimal Diabetes Care	<p>The percentage of adult diabetes patients who have optimally managed modifiable risk factors (A1c, LDL, blood pressure, tobacco non-use and daily aspirin usage for patients with diagnosis of ischemic vascular disease) with the intent of preventing or reducing future complications associated with poorly managed diabetes.</p> <p>Patients ages 18 - 75 with a diagnosis of diabetes, who meet all the numerator targets of this composite measure: A1c < 8.0, LDL < 100, Blood Pressure < 140/90, Tobacco non-user and for patients with diagnosis of ischemic vascular disease daily aspirin use unless contraindicated.</p> <p>Please note that while the all-or-none composite measure is considered to be the gold standard, reflecting best patient outcomes, the individual components may be measured as well. This is particularly helpful in quality improvement efforts to better understand where opportunities exist in moving the patients toward achieving all of the desired outcomes. Please refer to the additional numerator logic provided for each component.</p>	MN Community Measurement
0731	Comprehensive Diabetes Care	<p>The percentage of patients 18–75 years of age with diabetes (type 1 and type 2) who had each of the following:</p> <ul style="list-style-type: none"> - Hemoglobin A1c (HbA1c) testing (NQF#0057) - HbA1c poor control (>9.0%) (NQF#0059) - HbA1c control (<8.0%) (NQF#0575) - HbA1c control (<7.0%) for a selected population* - Eye exam (retinal) performed (NQF#0055) - LDL-C screening (NQF#0063) - LDL-C control (<100 mg/dL) (NQF#0064) - Medical attention for nephropathy (NQF#0062) - BP control (<140/90 mm Hg) (NQF#0061) - Smoking status and cessation advice or treatment 	National Committee for Quality Assurance