



NATIONAL QUALITY FORUM

Driving measurable health
improvements together

Memo

July 28, 2020

To: Consensus Standards Approval Committee (CSAC)

From: Geriatrics and Palliative Care Project Team

Re: Geriatrics and Palliative Care Fall 2019, Track 1 Measures

COVID-19 Updates

Considering the recent COVID-19 global pandemic, many organizations needed to focus their attention on the public health crisis. To provide greater flexibility for stakeholders and continue the important work in quality measurement, the National Quality Forum (NQF) extended commenting periods and adjusted measure endorsement timelines for the Fall 2019 cycle.

Commenting periods for all measures evaluated in the Fall 2019 cycle were extended from 30 days to 60 days. Based on the comments received during this 60-day extended commenting period, measures entered one of two tracks:

Track 1: Measures Continuing in Fall 2019 Cycle

Measures that did not receive public comments or only received comments in support of the Standing Committees' recommendations will be reviewed by the CSAC.

- **Exceptions**

Exceptions were granted to measures if non-supportive comments received during the extended post-comment period were similar to those received during the pre-evaluation meeting period and have already been adjudicated by the respective Standing Committees during the measure evaluation Fall 2019 meetings.

Track 2: Measures Deferred to Spring 2020 Cycle

Fall 2019 measures requiring further action or discussion from a Standing Committee were deferred to the Spring 2020 cycle. This includes measures where consensus was not reached or those that require a response to public comments received. Measures undergoing maintenance review will retain endorsement during that time. Track 2 measures will be reviewed during the CSAC's meeting in November.

During the CSAC meeting on July 28-29, the CSAC will review Fall 2019 measures assigned to Track 1. Evaluation summaries for measures in track 1 have been described in this memo and related Geriatrics and Palliative Care draft report. A list of measures assigned to Track 2 can be found in the Executive Summary section of the Geriatrics and Palliative Care draft report for tracking purposes and will be described further in a subsequent report. Measures in track 2 will be reviewed by the CSAC on November 17-18, 2020.

CSAC Action Required

The CSAC will review recommendations from the Geriatrics and Palliative Care project at its July 28-29, 2020 meeting and vote on whether to uphold the recommendations from the Committee.

<http://www.qualityforum.org>

This memo includes a summary of the project and measure recommendations. The following documents accompany this memo:

- **Geriatrics and Palliative Care Draft Report.** The draft report includes measure evaluation details on one measure assigned to Track 1. The measure assigned to Track 2 will be reviewed during the CSAC's meeting in November. The complete draft report and supplemental materials are available on the project webpage.

Background

Since 2006, when it first developed a measurement framework for palliative and end-of-life care, and endorsed 38 evidence-based preferred practices for high quality palliative care programs, NQF has endorsed more than 30 measures in this topic area, many of which currently are used in federal quality improvement and public reporting programs. In 2017, NQF expanded the scope of the Standing Committee charged with the oversight of NQF's portfolio of palliative and end-of-life care measures by adding measures specifically relevant to the geriatric population. This renamed "Geriatrics and Palliative Care Standing Committee" has the requisite expertise to evaluate and assume oversight of measures that focus on key issues specific to older adults.

Draft Report

The Geriatrics and Palliative Care Fall 2019 draft report presents the results of the evaluation of one measure considered under the Consensus Development Process (CDP). This measure is recommended for endorsement.

	Maintenance	New	Total
Measures under consideration	1	0	1
Measures recommended for endorsement	1	0	1

CSAC Action Required

Pursuant to the CDP, the CSAC is asked to consider endorsement of one candidate consensus measure.

Measure Recommended for Endorsement

- [1623: Bereaved Family Survey \(Department of Veterans Affairs\)](#)

Overall Suitability for Endorsement: Yes-16; No-1

Comments and Their Disposition

NQF did not receive any comments pertaining to measure 1623: Bereaved Family Survey.

Member Expression of Support

Throughout the 16-week continuous public commenting period, NQF members had the opportunity to express their support ('support' or 'do not support') for each measure submitted for endorsement consideration to inform the Committee's recommendations. No NQF members provided their expression of support for measure 1623: Bereaved Family Survey.

Appendix A: CSAC Checklist

The table below lists the key considerations to inform the CSAC's review of the measures submitted for endorsement consideration.

Key Consideration	Yes/No	Notes
Were there any process concerns raised during the CDP project? If so, briefly explain.	No	
Did the Standing Committee receive requests for reconsideration? If so, briefly explain.	No	
Did the Standing Committee overturn any of the Scientific Methods Panel's ratings of Scientific Acceptability? If so, state the measure and why the measure was overturned.	Yes	The two main concerns raised by the SMP were that the risk adjustment model did not include SDS, particularly race/ethnicity; and that the beta-binomial values presented as part of the construct validity were too low. In discussion with the Committee, the developer shared that they have updated testing results demonstrating stronger beta-binomial values and strong odds ratios. Regarding the measure's risk adjustment model, the developer clarified that this measure is developed for use by the U.S. Department of Veterans Affairs (VA) and that the VA's strong preference is to not apply risk adjustment to measures. There is concern about obscuring the source of variation in measure performance. Per NQF process, the SMP may recommend discussion points to the Committee regarding the use of SDS in risk adjustment models, but may not fail a measure solely for this reason. The Committee felt this was sufficient rationale to overturn the SMP rating.
If a recommended measure is a related and/or competing measure, was a rationale provided for the Standing Committee's recommendation? If not, briefly explain.	N/A	
Were any measurement gap areas addressed? If so, identify the areas.	No	
Are there additional concerns that require CSAC discussion? If so, briefly explain.	No	

Appendix B: Details of Measure Evaluation

1623 Bereaved Family Survey

[Submission](#) | [Specifications](#)

Description: This measure calculates the proportion of Veteran decedent's family members who rate overall satisfaction with the Veteran decedent's end-of-life care in an inpatient setting as "Excellent" versus "Very good," "good," "fair," or "poor."

Numerator Statement: The numerator is comprised of completed surveys (at least 12 of 17 structured items completed), where the global item question has an optimal response. The global item question asks "Overall, how would you rate the care that [Veteran] received in the last month of life" and the possible answer choices are: Excellent, Very good, Good, Fair, or Poor. The optimal response is Excellent.

Denominator Statement: The denominator consists of all inpatient deaths for which a survey was completed (at least 12 of 17 structured items completed), excluding: 1) deaths within 24 hours of admission (unless the Veteran had a previous hospitalization in the last month of life); 2) deaths that occur in the Emergency Department (unless the Veteran had a prior hospitalization of at least 24 hours in the last 31 days of life); Additional exclusion criteria include: 1) Veterans for whom a family member knowledgeable about their care cannot be identified (determined by the family member's report); or contacted (no current contacts listed or no valid addresses on file); 2) absence of a working telephone available to the family member.

Exclusions:

- Veterans for whom a family member knowledgeable about their care cannot be identified (determined by family member's report)
- Absence of a current address and/or working telephone number for a family member or emergency contact.
- Deaths within 24 hours of admission without a prior hospitalization of last least 24 hours in the last 31 days of life.
- Deaths that occur in the operating room during an outpatient procedure.
- Deaths due to a suicide or accident
- Surveys in which less than 12 items were answered.

Adjustment/Stratification: Statistical Risk Model

Level of Analysis: Facility, Other

Setting of Care: Inpatient/Hospital, Post-Acute Care

Type of Measure: Outcome: PRO-PM

Data Source: Instrument-Based Data

Measure Steward: Department of Veterans Affairs / Hospice and Palliative Care

STANDING COMMITTEE MEETING 2/20/2020 and 2/25/2020

1. Importance to Measure and Report: The measure meets the Importance criteria

(1a. Evidence, 1b. Performance Gap)

1a. Evidence: **Rating accepted from previous review cycle**; 1b. Performance Gap: **H-7; M-9; L-0; I-0**

Rationale:

- During the prior review in 2015, the developer provided a logic model stating that receiving a palliative care consult or dying in a hospice unit results in a greater likelihood of families rating end-of-life inpatient care as excellent. The developer included a recommendation from the 2009 version of the Clinical Practice Guidelines for Quality Palliative Care. In addition to the guideline recommendation, the developer stated that physical symptoms such as pain, nausea, constipation, and dyspnea are common at end of life, and that clinicians do not always recognize these symptoms or manage them appropriately. The developer stated that studies have found that providers do not communicate with patients about patients' healthcare preferences and that providers' treatment decisions may not be consistent with patients' preferences.
- The Committee agreed there was no change in evidence from previous endorsement, and agreed to accept the decision and vote from the previous review cycle, which was that the measure passes the evidence criterion.

1623 Bereaved Family Survey

- The developer provided results from 2017 (n=146 VA facilities) demonstrating a 65% mean overall score, a score range from 13%-100%, and IQR of 85 and 72. The Committee felt that there is a clear performance gap that warrants a national performance measure.

2. Scientific Acceptability of Measure Properties: The measure meets the Scientific Acceptability criteria

(2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity)

2a. Reliability: **Yes-16; No-0**; 2b. Validity: **H-2; M-11; L-2; I-1**

Rationale:

- Because the SMP rated this measure low on validity, Standing Committee deliberations started with a discussion of the SMP's rating, the rationale for that rating, and a vote on whether the Committee chose to accept that rating. Accepting the rating would have removed endorsement and ended discussion of the measure. The two main concerns raised by the SMP were that the risk adjustment model did not include SDS, particularly race/ethnicity; and that the beta-binomial values presented as part of the construct validity were too low. Per NQF process, the SMP may recommend discussion points to the Committee regarding the use of SDS in risk adjustment models, but may not fail a measure solely for this reason.
- In discussion with the Committee, the developer shared that they have updated testing results demonstrating stronger beta-binomial values and strong odds ratios, and that they would be happy to share this formally during the post-meeting public comment period. The Committee felt this was sufficient rationale to overturn the SMP rating and continue discussion of the measure.
- The Committee voted to overturn the SMP's validity rating: Accept-2; Overturn-14
- The Committee asked for some clarifications on the measure specifications, including the use of male pronouns in the survey, the exact scope and inclusions of the survey, and the grade level of some survey questions. The developer clarified that there are separate surveys for male and female patients, each with corresponding pronouns. The developer further clarified that the measure encompasses all deaths in a VA facility (and only in a VA facility), regardless of setting of care (hospice vs. intensive care). The developer noted that they offer an "unsure" option if caregivers are not sure how to answer a question, but agreed that appropriate grade-level content is a worthy goal. The developer hopes to include more survey questions in future endorsement submissions and will review the readability.
- The Committee asked the developer to elaborate on the rationale for the measure's risk adjustment model. The developer clarified that this measure is developed for use by the VA and that the VA's strong preference is to not apply risk adjustment to measures. There is concern about obscuring the source of variation in measure performance. The developer noted that they felt some risk adjustment was necessary, and they had developed their model to be closely aligned with the model for measure 2651. The Committee noted that measure 2651 does not include race/ethnicity in its risk adjustment model, yet this was not raised as a significant concern by either the SMP or the Committee. The Committee was satisfied with the explanation and rationale around risk adjustment, and the discussion turned to the construct validity concern. The developer reported that they have updated testing results that show beta-binomial values of 0.13-1.57 at the facility level and odds ratios of 1.44--19.16 at the national level between the measure under review and other accepted process measures. The developer stated they will be sharing these results through the commenting process. The Committee was satisfied that the measure meets the validity criterion.
- The Committee noted that the SMP rated the measure high on reliability.
- The SMP's ratings for reliability: H-3; M-2; L-0; I-1
- The Committee did not have any concerns with the measure meeting this criterion and voted unanimously to accept the SMP's rating.

3. Feasibility: H-2; M-15; L-0; I-0

(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/unintended consequences identified; 3d. Data collection strategy can be implemented)

Rationale:

- Committee members noted that while some of the data elements are available in the electronic health record, the key responses have to be gathered through mail or telephone surveys. The developer

1623 Bereaved Family Survey
<p>stated they have been refining both procedures for gathering electronic data and survey contact procedures for more efficient survey administration.</p> <ul style="list-style-type: none"> The Committee acknowledged that the measure developer is also the measure's main user and that this should result in a very feasible measure.
<p>4. Use and Usability</p> <p><i>4a. Use; 4a1. Accountability and transparency; 4a2. Feedback on the measure by those being measured and others; 4b. Usability; 4b1. Improvement; 4b2. The benefits to patients outweigh evidence of unintended negative consequences to patients)</i></p> <p>4a. Use: Pass-14; No Pass-0 4b. Usability: H-4; M-12; L-0; I-1</p> <p><u>Rationale:</u></p> <ul style="list-style-type: none"> The Committee had questions for NQF staff about the use criterion and for the developer about the current use of the measure. NQF clarified that use is currently a must-pass criterion for maintenance measures such as this measure, and that NQF defined use as publicly reported within six years of initial endorsement and use within an accountability program within three years. The developer reported that the measure is used for accountability across all VA facilities. They have been working to put a plan in place for public reporting. For VA patients, facility choice is based almost entirely on location. Publicly reporting the survey results would not assist in choosing care. The developer has been working to expand use of the measure in private facilities and health systems. They are also working to report results for nursing homes, where there may be more patient choice available, especially since the enactment of the VA MISSION Act of 2018. This reporting would require authorization from the Secretary of Veterans Affairs. A Committee member pointed out that veterans with Medicare coverage also have additional facility choices, and this could be a potential focus area for reporting. The Committee was willing to accept the developer's plan for public reporting, but strongly encouraged the measure be publicly reported. The Committee stated they expect to see the measure reported when it returns for its next maintenance endorsement. The Committee had no concerns about the usability of the measure.
<p>5. Related and Competing Measures</p> <ul style="list-style-type: none"> This measure is related to NQF 2651 CAHPS Hospice Survey. <ul style="list-style-type: none"> The developer stated that the populations are different for these two measures, as measure 1623 is focused on deaths in a VA inpatient setting. The Committee engaged in a brief discussion of 1623 and 2651 as related measures. The Committee felt there was a clear difference between the two measures, and stated they are different measures with different populations. The Committee felt the differences between the VA and other health systems justified different measures. Committee members did identify areas, such as questions around supports, where the content of the questions could be more aligned, stating there is strong evidence around best practices in these areas.
<p>6. Standing Committee Recommendation for Endorsement: Y-16; N-1</p>
<p>7. Public and Member Comment</p> <ul style="list-style-type: none"> No public and member comments were received.
<p>8. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X</p>
<p>9. Appeals</p>



**NATIONAL
QUALITY FORUM**

Driving measurable health
improvements together

<http://www.qualityforum.org>

Geriatrics and Palliative Care Fall 2019 Review Cycle

CSAC Review and Endorsement

July 28 and 29, 2020

Standing Committee Recommendations

- Two measures reviewed for Fall 2019
 - ▣ Two measures reviewed by the Scientific Methods Panel
- One measure recommended for endorsement
 - ▣ NQF 1623: Bereaved Family Survey
- One measure deferred to Spring 2020 due to COVID-19 extended commenting periods
 - ▣ NQF 2651: CAHPS® Hospice Survey (experience with care)

Overarching Issues

- Concerns Raised by the Scientific Methods Panel and Discussed by the Committee
 - ▣ Concerns that the risk adjustment model did not include SDS, particularly race/ethnicity
 - ▣ Concerns that the beta-binomial values presented as part of the construct validity were too low
- Applying NQF's Use Criterion
 - ▣ Measure is used for accountability across all VA facilities but is not publicly reported
 - ▣ Current criteria guidance defines use as publicly reported within six years of initial endorsement and use within an accountability program within three years
 - ▣ The Committee accepted the developer's plan for public reporting



Public and Member Comment and Member Expressions of Support

- No comments were received on NQF 1623
- No NQF member of expressions of support received



Timeline and Next Steps

Process Step	Timeline
CSAC Endorsement Meeting	July 28 - 29, 2020
Appeals Period	August 3 – September 1, 2020



Questions?

- Project team:
 - ▣ Katie Goodwin, Director
 - ▣ Erin Buchanan, Manager
 - ▣ Mike DiVecchia, Project Manager
 - ▣ Ngozi Ihenacho, Analyst
- Project webpage:
[https://www.qualityforum.org/Geriatrics and Palliative Care.aspx](https://www.qualityforum.org/Geriatrics_and_Palliative_Care.aspx)
- Project email address: Palliative@qualityforum.org

THANK YOU.

NATIONAL QUALITY FORUM

<http://www.qualityforum.org>

Geriatrics and Palliative Care, Fall 2019 Cycle, Track 1: CDP Report

**DRAFT REPORT FOR CSAC REVIEW
JULY 28-29, 2020**

This report is funded by the Department of Health and Human Services under contract HHSM-500-2017-00060I Task Order HHSM-500-T0001

<http://www.qualityforum.org>

Contents

Executive Summary.....	3
Introduction	3
NQF Portfolio of Performance Measures for Geriatrics and Palliative Care Conditions	5
Table 1. NQF Geriatrics and Palliative Care Portfolio of Measures	5
Geriatrics and Palliative Care Measure Evaluation	6
Table 2. Geriatrics and Palliative Care Measure Evaluation Summary, Fall 2019 – Track 1	6
Comments Received Prior to Committee Evaluation	6
Comments Received After Committee Evaluation	6
Summary of Measure Evaluation: Fall 2019 Measures, Track 1	7
References.....	9
Appendix A: Details of Measure Evaluation	11
Measures Recommended	11
1623 Bereaved Family Survey	11
Appendix B: Geriatrics and Palliative Care Portfolio—Use in Federal Programs	15
Appendix C: Geriatrics and Palliative Care Standing Committee and NQF Staff.....	18
Appendix D: Measure Specifications.....	21
1623 Bereaved Family Survey	21
Appendix E: Related and Competing Measures	24

Executive Summary

Improving the quality of palliative and end-of-life care, and geriatric care more generally, is becoming increasingly important due to factors that have intensified the need for individualized, person-centered care. Some of these factors include the aging U.S. population; the projected increases in the number of Americans with chronic illnesses, disabilities, and functional limitations; and increases in ethnic and cultural diversity.

Palliative care is patient- and family-centered care that optimizes quality of life by anticipating, preventing, and alleviating suffering throughout the continuum of a person's illness by addressing physical, intellectual, emotional, social, and spiritual needs and facilitating patient autonomy, access to information, and choice.¹ Palliative care is holistic, thus requiring an interdisciplinary, team-based approach to care. With its focus on improving quality of life, palliative care is distinct from care intended to cure an illness or condition, although it can be delivered concurrently with curative therapies, and can begin at any point in the disease progression. It can be provided in any setting, including outpatient care settings and at home.

To date, the National Quality Forum (NQF) has endorsed more than 30 measures that address geriatric care, palliative care, and end-of-life care. These measures address physical, spiritual, and legal aspects of care, as well as the care of patients nearing the end of life.

Due to circumstances around the COVID-19 global pandemic, commenting periods for all measures evaluated in the Fall 2019 cycle were extended from 30 days to 60 days. Based on the comments received during this 60-day extended commenting period, measures entered into one of two tracks:

Track 1: measures continuing its review in Fall 2019 Cycle:

Recommended for Endorsement

- **NQF 1623:** Bereaved Family Survey

Track 2: measures deferred to Spring 2020 Cycle:

- **NQF 2651:** CAHPS® Hospice Survey (experience with care)

This report contains details of the evaluation of the measure assigned to *Track 1* and are continuing in the Fall 2019 cycle. The detailed evaluation summary of the measure assigned to *Track 2* and deferred to the Spring 2020 cycle will be included in a subsequent report. Brief summaries of the Fall 2019 *Track 1* measures currently under review are included in the body of the report; detailed summaries of the Committee's discussion and ratings of the criteria for each measure are in [Appendix A](#).

Introduction

Improving the quality of both palliative and end-of-life care, and geriatric care more generally, is becoming increasingly important due to factors that have intensified the need for individualized, person-centered care. Some of these factors include the aging U.S. population; the projected increases in the

number of Americans with chronic illnesses, disabilities, and functional limitations; and increases in ethnic and cultural diversity.² In 2018, the 65-and-older population numbered 50.9 million individuals (15.6 percent of the U.S. population), and this figure is expected to increase to 94.7 million by 2060.³ As many as 35 percent of older Americans have some type of disability (e.g., vision, hearing, ambulation, cognition), while 46 percent of those 75 and older report limitations in physical functioning.⁴ Additionally, data indicate that 46 percent of the noninstitutionalized U.S. population aged 65 or older have two or three chronic conditions, and 15 percent have four or more.⁵

Palliative care is patient- and family-centered care that optimizes quality of life by anticipating, preventing, and alleviating suffering throughout the continuum of a person's illness by addressing physical, intellectual, emotional, social, and spiritual needs and facilitating patient autonomy, access to information, and choice.⁶ Palliative care is holistic, thus requiring an interdisciplinary, team-based approach to care. With its focus on improving quality of life, palliative care is distinct from care intended to cure an illness or condition, although it can be delivered concurrently with curative therapies, and can begin at any point in the disease progression. It can be provided in any setting, including outpatient care settings and at home.

Although palliative care is still provided primarily by specially trained teams of professionals in hospitals and through hospice, there is increased focus on provision of palliative care in the community,⁷ often by clinicians who are not palliative care specialists. The provision of palliative care has been shown to increase patient and family satisfaction with care,⁸ reduce emergency department visits, hospital admissions, and hospital readmissions,⁹ and decrease costs to the healthcare system.^{10,11} However, access to hospital-based specialty palliative care continues to vary by hospital size and location, and even when programs are available, not all patients who could benefit actually receive those services.¹²

Palliative care is appropriate for those who are expected to recover, as well as for those who have chronic, progressive, and/or terminal illness. For those with a terminal illness, high quality end-of-life care is comprehensive care that addresses medical, emotional, spiritual, and social needs during the last stages of illness.¹³ Much end-of-life care is palliative, when life-prolonging interventions are no longer appropriate, effective, or desired.¹⁴ Thus, for patients nearing the end of life, there often will be a greater emphasis on palliative care over curative treatment. In many instances, this care is provided in the form of hospice.

Hospice is a service delivery system that relies on an interdisciplinary approach that emphasizes symptom management for patients near the end of life. While hospice care is covered through Medicaid and most private insurance plans, approximately 85 percent of hospice enrollees receive coverage through the Medicare hospice benefit.¹⁵ Almost 1.5 million Medicare beneficiaries and their families received hospice care in 2017.¹⁶ For these individuals, the average length of stay was 76.1 days; however, the median length of stay was only 24 days, meaning that many enrolled in hospice too late to fully realize the benefits of the program.¹⁷ Beginning in 2014, Medicare-certified hospices were required to report performance on quality measures as part of the Hospice Quality Reporting Program (HQRP); those not reporting face a reduction in payments from Medicare. Performance rates for these measures are publicly reported on the Centers for Medicare and Medicaid Services (CMS) Hospice Compare website.¹⁸

Since 2006, when it first developed a measurement framework for palliative and end-of-life care, and endorsed 38 evidence-based preferred practices for high quality palliative care programs,¹⁹ NQF has endorsed more than 30 measures in this topic area, many of which currently are used in federal quality improvement and public reporting programs.

In 2017, NQF expanded the scope of the Standing Committee charged with the oversight of the palliative and end-of-life care measures portfolio by adding measures specifically relevant to older adults (i.e., the geriatric population). Several previously seated and new members of this renamed “Geriatrics and Palliative Care Standing Committee” are geriatric healthcare professionals. Thus, the Committee has the requisite expertise to assume oversight of measures that focus on key issues specific to older adults, such as multimorbidity and frailty. At present, measures specifically relevant to the geriatric population remain aspirational. Thus, for the time being, the geriatrics measures evaluated by this Committee include setting-specific measures that primarily affect older individuals. Examples of such measures include those that assess care provided by home health agencies or other home-based care providers.

NQF Portfolio of Performance Measures for Geriatrics and Palliative Care Conditions

The Geriatrics and Palliative Care Standing Committee ([Appendix C](#)) oversees NQF’s portfolio of Geriatrics and Palliative Care measures ([Appendix B](#)). This portfolio contains 36 measures: 18 process measures, 17 outcome measures, and one composite measure (see table below).

Table 1. NQF Geriatrics and Palliative Care Portfolio of Measures

	Process	Outcome	Composite
Palliative/End-of-Life Care			
Physical Aspects of Care	9	–	–
Psychological and Psychiatric Aspects of Care	–	–	–
Social Aspects of Care	–	–	–
Spiritual, Religious, and Existential Aspects of Care	1	–	–
Cultural Aspects of Care	–	–	–
Care of the Patient Nearing the End of Life	3	12	1
Ethical and Legal Aspects of Care	3	–	–
Geriatrics	2	5	–
Total	18	17	1

Some of the measures in the Geriatrics and Palliative Care portfolio will be evaluated by other NQF standing committees. These include a cultural communication measure (Patient Experience and Function Committee) and pain measures for cancer patients (Cancer Committee).

Geriatrics and Palliative Care Measure Evaluation

On February 20 and 25, 2020, the Geriatrics and Palliative Care Standing Committee evaluated two measures undergoing maintenance review against NQF's [standard measure evaluation criteria](#). One measure was assigned to *Track 1* and is continuing in the Fall 2019 cycle. The detailed evaluation summary of one measure assigned to *Track 2* and deferred to the Spring 2020 cycle will be included in a subsequent report.

Table 2. Geriatrics and Palliative Care Measure Evaluation Summary, Fall 2019 – Track 1

	Maintenance	New	Total
Measures under consideration	1	0	1
Measures recommended for endorsement	1	0	1

Comments Received Prior to Committee Evaluation

NQF solicits comments on endorsed measures on an ongoing basis through the [Quality Positioning System \(QPS\)](#). In addition, NQF solicits comments for a continuous 16-week period during each evaluation cycle via an online tool located on the project webpage. For this evaluation cycle, the commenting period opened on December 11, 2019. NQF did not receive any comments on the measures as of January 31, 2020.

Comments Received After Committee Evaluation

Considering the recent COVID-19 global pandemic, many organizations needed to focus their attention on the public health crisis. In order to provide greater flexibility for stakeholders and continue the important work in quality measurement, the National Quality Forum (NQF) extended commenting periods and adjusted measure endorsement timelines for the Fall 2019 cycle.

Commenting periods for all measures evaluated in the Fall 2019 cycle were extended from 30 days to 60 days. Based on the comments received during this 60-day extended commenting period, measures entered one of two tracks:

Track 1: Measures Continuing in Fall 2019 Cycle

Measures that did not receive public comments or only received comments in support of the Standing Committees' recommendations will move forward to the CSAC for review and discussion during its meeting on July 28-29.

○ Exceptions

Exceptions were granted to measures if non-supportive comments received during the extended post-comment period were similar to those received during the pre-evaluation meeting period and have already been adjudicated by the respective Standing Committees during the measure evaluation Fall 2019 meetings.

Track 2: Measures Deferred to Spring 2020 Cycle

Fall 2019 measures requiring further action or discussion from a Standing Committee were

deferred to the Spring 2020 cycle. This includes measures where consensus was not reached or those that require a response to public comments received. Measures undergoing maintenance review will retain endorsement during that time.

The extended public commenting period with NQF member support closed on May 28, 2020. NQF did not receive any comments pertaining to the measure assigned to Track 1.

During the Fall 2019 CSAC meeting on July 28-29, the Consensus Standards Approval Committee (CSAC) will review all measures assigned to Track 1. A list of measures assigned to Track 2 can be found in the [Executive Summary section](#) of this report for tracking purposes, but these measures will be reviewed by CSAC on November 17 and 18, 2020.

Summary of Measure Evaluation: Fall 2019 Measures, Track 1

The following brief summaries of the measure evaluation highlight the major issues that the Committee considered. Details of the Committee's discussion and ratings of the criteria for each measure are included in [Appendix A](#).

1623 Bereaved Family Survey (Department of Veterans Affairs/Hospice and Palliative Care): Recommended

Description: This measure calculates the proportion of veteran decedent's family members who rate overall satisfaction with the veteran decedent's end-of-life care in an inpatient setting as "Excellent" versus "Very good," "good," "fair," or "poor"; **Measure Type:** Outcome: PRO-PM; **Level of Analysis:** Facility, Other; **Setting of Care:** Inpatient/Hospital, Post-Acute Care; **Data Source:** Instrument-Based Data

Research has emphasized the extent to which end-of-life care in the United States needs to be improved. Commonly experienced symptoms like constipation, dyspnea, pain, and nausea are often not recognized by clinicians and therefore not adequately managed. Additionally, healthcare preferences are not always communicated between patients and providers, and those preferences are not always consistent with providers' treatment decision. The strategy of assessing the quality of end of life with a post-death survey of family members has become known as critically important as an essential source of data that define the quality of end-of-life care for patients and their family members.

Because the Scientific Methods Panel (SMP) rated this measure low on validity, Committee deliberations started with a discussion of the SMP's rating, the rationale for that rating, and a vote on whether the Committee chose to accept that rating. Accepting the rating would end discussion of the measure and the measure would lose endorsement. Rejection of the rating would allow the Committee to continue discussion of the measure. The two main concerns raised by the SMP were that the risk adjustment model did not include socio-demographic status (SDS), particularly race/ethnicity, and that the beta-binomial values presented as part of the construct validity were too low. Per NQF process, the SMP may recommend discussion points to the Committee regarding the use of SDS in risk adjustment models, but may not fail a measure solely for this reason. In discussion with the Committee, the developer shared that they have updated testing results demonstrating stronger beta-binomial values and strong odds ratios, and that they would be happy to share this formally during the post-meeting public comment

period. The Committee felt this was sufficient rationale to overturn the SMP rating and continue discussion of the measure.

The Committee asked the developer to elaborate on the rationale for the measure's risk adjustment model. The developer clarified that this measure is developed for use by the U.S. Department of Veterans Affairs (VA) and that the VA's strong preference is to not apply risk adjustment to measures. There is concern about obscuring the source of variation in measure performance. The developer noted that they felt some risk adjustment was necessary and they had developed their model to be closely aligned with the model for measure 2651. The Committee noted that measure 2651 does not include race/ethnicity in its risk adjustment model, yet this was not raised as a significant concern by either the SMP or the Committee. The Committee was satisfied with the explanation and rationale around risk adjustment and the discussion turned to the construct validity concern. The developer reported that they have updated testing results that show beta-binomial values of 0.13-1.57 at the facility level and odds ratios of 1.44-19.16 at the national level between the measure under review and other accepted process measures. The developer stated they will be sharing these results through the commenting process. The Committee was satisfied that the measure meets the validity criterion.

The Committee agreed there was no change in evidence from the previous endorsement review, and agreed to accept the decision and vote from the measure's previous review cycle, which was that the measure passes the evidence criterion. The Committee agreed that there is a clear performance gap that warrants a national performance measure. The Committee noted that the SMP rated the measure high on reliability. The Committee asked for some clarifications on the measure specifications, including the use of male pronouns in the survey, the exact scope and inclusions of the survey, and the grade level of some survey questions. The Committee accepted the developer's response and was satisfied that the measure met the reliability criterion.

The Committee noted that the measure developer is also the measure's main user and that this should result in a very feasible measure. The Committee had questions about the current use of the measure. The developer reported that the measure is used for accountability across all VA facilities, and they have been working to put a plan in place for public reporting. The developer has been working to expand use of the measure in private facilities and health systems. They are also working to report results for nursing homes, where there may be more patient choice available, especially since the enactment of the VA MISSION Act of 2018. The Committee accepted the developer's plan for public reporting but strongly encouraged the measure be publicly reported. The Committee stated they expect to see the measure reported when it returns for its next maintenance endorsement. The Committee had no concerns about the usability of the measure.

References

- ¹ National Quality Forum (NQF). *A National Framework and Preferred Practices for Palliative and Hospice Care Quality: A Consensus Report*. Washington, DC: NQF; 2006.
<http://www.qualityforum.org/WorkArea/linkit.aspx?LinkIdentifier=id&ItemID=22041>. Last accessed June 2016.
- ² Institute of Medicine (IOM). *Dying in America: Improving Quality and Honoring Individual Preferences near the End of Life*. Washington, DC: The National Academies Press; 2014.
<http://www.nationalacademies.org/hmd/Reports/2014/Dying-In-America-Improving-Quality-and-Honoring-Individual-Preferences-Near-the-End-of-Life.aspx>. Last accessed June 2016.
- ³ Administration for Community Living. *2018 Profile of Older Americans*. Washington, DC: U.S. Department of Health and Human Services; April 2018.
<https://acl.gov/sites/default/files/Aging%20and%20Disability%20in%20America/2017OlderAmericansProfile.pdf>. Last accessed July 2019.
- ⁴ Administration for Community Living. *2018 Profile of Older Americans*. Washington, DC: U.S. Department of Health and Human Services; April 2018.
<https://acl.gov/sites/default/files/Aging%20and%20Disability%20in%20America/2017OlderAmericansProfile.pdf>. Last accessed July 2019.
- ⁵ Ward BW, Schiller JS. Prevalence of multiple chronic conditions among US adults: estimates from the National Health Interview Survey, 2010. *Prev Chronic Dis*. 2013;10:E65.
- ⁶ National Quality Forum (NQF). *A National Framework and Preferred Practices for Palliative and Hospice Care Quality: A Consensus Report*. Washington, DC: NQF; 2006.
<http://www.qualityforum.org/WorkArea/linkit.aspx?LinkIdentifier=id&ItemID=22041>. Last accessed June 2016.
- ⁷ Aldridge MD, Hasselaar J, Garralda E, et al. Education, implementation, and policy barriers to greater integration of palliative care: a literature review. *Palliat Med*. 2016;30(3):224-239.
- ⁸ Casarett DS, Shreve C, Luhrs K, et al. Measuring families' perceptions of care across a health care system: preliminary experience with the Family Assessment of Treatment at End of Life Short form (FATE-S). *J Pain Symptom Manage*. 2010;40(6):801-809.
- ⁹ Scibetta C, Kerr K, McGuire J, Rabow MW. The Costs of Waiting: Implications of the Timing of Palliative Care Consultation Among a Cohort of Decedents at a Comprehensive Cancer Center. *J Palliat Med*. 2016;19(1):69-75.
- ¹⁰ Lustbader D, Mudra M, Romano C, et al. The impact of a home-based palliative care program in an accountable care organization. *J Palliat Med*. 2016;20(1):23-28.

¹¹ Krakauer, R, Spettell C, Reisman L, et al. Opportunities to improve the quality of care for advanced illness. *Health Aff (Millwood)*. 2009;28(5):1357-1359.

¹² Morrison RS, Meier DE, Dumanovsky T, Rogers M. *America's Care of Serious Illness: 2015 State-By-State Report Card on Access to Palliative Care in Our Nation's Hospitals*. New York: Center to Advance Palliative Care and National Palliative Care Research Center; 2015. <https://reportcard.capc.org/>. Last accessed June 2016.

¹³ Institute of Medicine (IOM). *Dying in America: Improving Quality and Honoring Individual Preferences near the End of Life*. Washington, DC: The National Academies Press; 2014. <http://www.nationalacademies.org/hmd/Reports/2014/Dying-In-America-Improving-Quality-and-Honoring-Individual-Preferences-Near-the-End-of-Life.aspx>. Last accessed June 2016.

¹⁴ National Quality Forum (NQF). *A National Framework and Preferred Practices for Palliative and Hospice Care Quality: A Consensus Report*. Washington, DC: NQF; 2006. <http://www.qualityforum.org/WorkArea/linkit.aspx?LinkIdentifier=id&ItemID=22041>. Last accessed June 2016.

¹⁵ National Hospice and Palliative Care Organization (NHPCO). *NHPCO's Facts and Figures: Hospice Care in America 2015 Edition*. Alexandria, VA: NHPCO; 2015. http://www.nhpc.org/sites/default/files/public/Statistics_Research/2015_Facts_Figures.pdf. Last accessed March 2019.

¹⁶ National Hospice and Palliative Care Organization (NHPCO). *NHPCO Facts and Figures 2018 Edition*. Alexandria, VA: NHPCO; 7/2/2019. https://39k5cm1a9u1968hg74aj3x51-wpengine.netdna-ssl.com/wp-content/uploads/2019/07/2018_NHPCO_Facts_Figures.pdf. Last accessed July 2019.

¹⁷ National Hospice and Palliative Care Organization (NHPCO). *NHPCO Facts and Figures 2018 Edition*. Alexandria, VA: NHPCO; 2018. 7/2/2019. https://39k5cm1a9u1968hg74aj3x51-wpengine.netdna-ssl.com/wp-content/uploads/2019/07/2018_NHPCO_Facts_Figures.pdf. Last accessed July 2019.

¹⁸ Centers for Medicare & Medicaid Services. Hospice Compare website. <https://www.medicare.gov/hospicecompare/>. Last accessed March 2019.

¹⁹ National Quality Forum (NQF). *A National Framework and Preferred Practices for Palliative and Hospice Care Quality: A Consensus Report*. Washington, DC: NQF; 2006. <http://www.qualityforum.org/WorkArea/linkit.aspx?LinkIdentifier=id&ItemID=22041>. Last accessed June 2016.

Appendix A: Details of Measure Evaluation

Rating Scale: H=High; M=Moderate; L=Low; I=Insufficient; NA=Not Applicable

Measures Recommended

1623 Bereaved Family Survey
Submission Specifications
<p>Description: This measure calculates the proportion of Veteran decedent's family members who rate overall satisfaction with the Veteran decedent's end-of-life care in an inpatient setting as "Excellent" versus "Very good," "good," "fair," or "poor."</p> <p>Numerator Statement: The numerator is comprised of completed surveys (at least 12 of 17 structured items completed), where the global item question has an optimal response. The global item question asks "Overall, how would you rate the care that [Veteran] received in the last month of life" and the possible answer choices are: Excellent, Very good, Good, Fair, or Poor. The optimal response is Excellent.</p> <p>Denominator Statement: The denominator consists of all inpatient deaths for which a survey was completed (at least 12 of 17 structured items completed), excluding: 1) deaths within 24 hours of admission (unless the Veteran had a previous hospitalization in the last month of life); 2) deaths that occur in the Emergency Department (unless the Veteran had a prior hospitalization of at least 24 hours in the last 31 days of life); Additional exclusion criteria include: 1) Veterans for whom a family member knowledgeable about their care cannot be identified (determined by the family member's report); or contacted (no current contacts listed or no valid addresses on file); 2) absence of a working telephone available to the family member.</p> <p>Exclusions:</p> <ul style="list-style-type: none"> - Veterans for whom a family member knowledgeable about their care cannot be identified (determined by family member's report) - Absence of a current address and/or working telephone number for a family member or emergency contact. - Deaths within 24 hours of admission without a prior hospitalization of last least 24 hours in the last 31 days of life. - Deaths that occur in the operating room during an outpatient procedure. - Deaths due to a suicide or accident - Surveys in which less than 12 items were answered. <p>Adjustment/Stratification: Statistical Risk Model</p> <p>Level of Analysis: Facility, Other</p> <p>Setting of Care: Inpatient/Hospital, Post-Acute Care</p> <p>Type of Measure: Outcome: PRO-PM</p> <p>Data Source: Instrument-Based Data</p> <p>Measure Steward: Department of Veterans Affairs / Hospice and Palliative Care</p>
<p>STANDING COMMITTEE MEETING 2/20/2020 and 2/25/2020</p> <p>1. Importance to Measure and Report: <u>The measure meets the Importance criteria</u> (1a. Evidence, 1b. Performance Gap)</p> <p>1a. Evidence: Rating accepted from previous review cycle; 1b. Performance Gap: H-7; M-9; L-0; I-0</p> <p><u>Rationale:</u></p> <ul style="list-style-type: none"> • During the prior review in 2015, the developer provided a logic model stating that receiving a palliative care consult or dying in a hospice unit results in a greater likelihood of families rating end-of-life inpatient care as excellent. The developer included a recommendation from the 2009 version of the Clinical Practice Guidelines for Quality Palliative Care. In addition to the guideline recommendation, the developer stated that physical symptoms such as pain, nausea, constipation, and dyspnea are common at end of life, and that clinicians do not always recognize these symptoms or manage them appropriately. The developer stated that studies have found that providers do not communicate with

1623 Bereaved Family Survey

patients about patients' healthcare preferences and that providers' treatment decisions may not be consistent with patients' preferences.

- The Committee agreed there was no change in evidence from previous endorsement, and agreed to accept the decision and vote from the previous review cycle, which was that the measure passes the evidence criterion.
- The developer provided results from 2017 (n=146 VA facilities) demonstrating a 65% mean overall score, a score range from 13%-100%, and IQR of 85 and 72. The Committee felt that there is a clear performance gap that warrants a national performance measure.

2. Scientific Acceptability of Measure Properties: The measure meets the Scientific Acceptability criteria

(2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity)

2a. Reliability: **Yes-16; No-0**; 2b. Validity: **H-2; M-11; L-2; I-1**

Rationale:

- Because the SMP rated this measure low on validity, Standing Committee deliberations started with a discussion of the SMP's rating, the rationale for that rating, and a vote on whether the Committee chose to accept that rating. Accepting the rating would have removed endorsement and ended discussion of the measure. The two main concerns raised by the SMP were that the risk adjustment model did not include SDS, particularly race/ethnicity; and that the beta-binomial values presented as part of the construct validity were too low. Per NQF process, the SMP may recommend discussion points to the Committee regarding the use of SDS in risk adjustment models, but may not fail a measure solely for this reason.
- In discussion with the Committee, the developer shared that they have updated testing results demonstrating stronger beta-binomial values and strong odds ratios, and that they would be happy to share this formally during the post-meeting public comment period. The Committee felt this was sufficient rationale to overturn the SMP rating and continue discussion of the measure.
- The Committee voted to overturn the SMP's validity rating: **Accept-2; Overturn-14**
- The Committee asked for some clarifications on the measure specifications, including the use of male pronouns in the survey, the exact scope and inclusions of the survey, and the grade level of some survey questions. The developer clarified that there are separate surveys for male and female patients, each with corresponding pronouns. The developer further clarified that the measure encompasses all deaths in a VA facility (and only in a VA facility), regardless of setting of care (hospice vs. intensive care). The developer noted that they offer an "unsure" option if caregivers are not sure how to answer a question, but agreed that appropriate grade-level content is a worthy goal. The developer hopes to include more survey questions in future endorsement submissions and will review the readability.
- The Committee asked the developer to elaborate on the rationale for the measure's risk adjustment model. The developer clarified that this measure is developed for use by the VA and that the VA's strong preference is to not apply risk adjustment to measures. There is concern about obscuring the source of variation in measure performance. The developer noted that they felt some risk adjustment was necessary, and they had developed their model to be closely aligned with the model for measure 2651. The Committee noted that measure 2651 does not include race/ethnicity in its risk adjustment model, yet this was not raised as a significant concern by either the SMP or the Committee. The Committee was satisfied with the explanation and rationale around risk adjustment, and the discussion turned to the construct validity concern. The developer reported that they have updated testing results that show beta-binomial values of 0.13-1.57 at the facility level and odds ratios of 1.44--19.16 at the national level between the measure under review and other accepted process measures. The developer stated they will be sharing these results through the commenting process. The Committee was satisfied that the measure meets the validity criterion.
- The Committee noted that the SMP rated the measure high on reliability.
- The SMP's ratings for reliability: **H-3; M-2; L-0; I-1**
- The Committee did not have any concerns with the measure meeting this criterion and voted unanimously to accept the SMP's rating.

3. Feasibility: **H-2; M-15; L-0; I-0**

1623 Bereaved Family Survey

(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/unintended consequences identified; 3d. Data collection strategy can be implemented)

Rationale:

- Committee members noted that while some of the data elements are available in the electronic health record, the key responses have to be gathered through mail or telephone surveys. The developer stated they have been refining both procedures for gathering electronic data and survey contact procedures for more efficient survey administration.
- The Committee acknowledged that the measure developer is also the measure's main user and that this should result in a very feasible measure.

4. Use and Usability

4a. Use; 4a1. Accountability and transparency; 4a2. Feedback on the measure by those being measured and others; 4b. Usability; 4b1. Improvement; 4b2. The benefits to patients outweigh evidence of unintended negative consequences to patients)

4a. Use: **Pass-14; No Pass-0** 4b. Usability: **H-4; M-12; L-0; I-1**

Rationale:

- The Committee had questions for NQF staff about the use criterion and for the developer about the current use of the measure. NQF clarified that use is currently a must-pass criterion for maintenance measures such as this measure, and that NQF defined use as publicly reported within six years of initial endorsement and use within an accountability program within three years.
- The developer reported that the measure is used for accountability across all VA facilities. They have been working to put a plan in place for public reporting. For VA patients, facility choice is based almost entirely on location. Publicly reporting the survey results would not assist in choosing care.
- The developer has been working to expand use of the measure in private facilities and health systems. They are also working to report results for nursing homes, where there may be more patient choice available, especially since the enactment of the VA MISSION Act of 2018. This reporting would require authorization from the Secretary of Veterans Affairs. A Committee member pointed out that veterans with Medicare coverage also have additional facility choices, and this could be a potential focus area for reporting.
- The Committee was willing to accept the developer's plan for public reporting, but strongly encouraged the measure be publicly reported. The Committee stated they expect to see the measure reported when it returns for its next maintenance endorsement. The Committee had no concerns about the usability of the measure.

5. Related and Competing Measures

- This measure is related to NQF 2651 CAHPS Hospice Survey.
 - The developer stated that the populations are different for these two measures, as measure 1623 is focused on deaths in a VA inpatient setting.
- The Committee engaged in a brief discussion of 1623 and 2651 as related measures. The Committee felt there was a clear difference between the two measures, and stated they are different measures with different populations. The Committee felt the differences between the VA and other health systems justified different measures. Committee members did identify areas, such as questions around supports, where the content of the questions could be more aligned, stating there is strong evidence around best practices in these areas.

6. Standing Committee Recommendation for Endorsement: Y-16; N-1**7. Public and Member Comment**

- No public and member comments were received.
- During the comment period, the developer provided updated testing results to supplement the committee's recommendation to endorse the measure.

8. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X

1623 Bereaved Family Survey
9. Appeals

Appendix B: Geriatrics and Palliative Care Portfolio—Use in Federal Programs^a

NQF #	Title	Federal Programs: Finalized or Implemented as of January 13, 2020
0167	Improvement in Ambulation and Locomotion	Home Health Value Based Purchasing (Implemented)
0174	Improvement in Bathing	Home Health Value Based Purchasing (Implemented)
0175	Improvement in Bed Transferring	Home Health Value Based Purchasing (Implemented)
0176	Improvement in Management of Oral Medications	Home Health Value Based Purchasing (Implemented) Home Health Quality Reporting (Implemented)
0177	Improvement in pain interfering with activity	Home Health Value Based Purchasing (Implemented) Home Health Quality Reporting (Implemented)
0209	Comfortable Dying: Pain Brought to a Comfortable Level Within 48 Hours of Initial Assessment	N/A
0383	Oncology: Plan of Care for Pain – Medical Oncology and Radiation Oncology (paired with 0384)	Hospital Care (Implemented) Prospective Payment System-Exempt Cancer Hospital Quality Reporting (Implemented) Merit-Based Incentive Payment System (MIPS) Program (Finalized)
0384	Oncology: Medical and Radiation - Pain Intensity Quantified (paired with 0383)	Merit-Based Incentive Payment System (MIPS) Program (Implemented) Medicaid Promoting Interoperability Program (Proposed)
0420	Pain Assessment and Follow-Up	N/A
1617	Patients Treated with an Opioid who are Given a Bowel Regimen	Hospice Quality Reporting (Implemented)
1628	Patients with Advanced Cancer Screened for Pain at Outpatient Visits	Prospective Payment System-Exempt Cancer Hospital Quality Reporting (Considered)
1634	Hospice and Palliative Care — Pain Screening	Hospice Quality Reporting (Implemented)
1637	Hospice and Palliative Care — Pain Assessment	Hospice Quality Reporting (Implemented)
1638	Hospice and Palliative Care — Dyspnea Treatment	Hospice Quality Reporting (Implemented)
1639	Hospice and Palliative Care — Dyspnea Screening	Hospice Quality Reporting (Implemented)

^a Per CMS Measures Inventory Tool as of 03/10/2020

NQF #	Title	Federal Programs: Finalized or Implemented as of January 13, 2020
1647	Beliefs and Values - Percentage of hospice patients with documentation in the clinical record of a discussion of spiritual/religious concerns or documentation that the patient/caregiver did not want to discuss	Hospice Quality Reporting (Implemented)
0326	Advance Care Plan	Home Health Value Based Purchasing (Implemented) Merit-Base Incentive Payment System (MIPS) Program (Finalized) Ambulatory Surgical Center Quality Reporting (Considered) Hospital Outpatient Quality Reporting (Considered)
1626	Patients Admitted to ICU who Have Care Preferences Documented	N/A
1641	Hospice and Palliative Care – Treatment Preferences	Prospective Payment System-Except Cancer Hospital Quality Reporting (Considered) Hospice Quality Reporting (Implemented)
0210	Proportion receiving chemotherapy in the last 14 days of life	Merit-Base Incentive Payment System (MIPS) Program (Finalized) Hospital Compare (Finalized) Prospective Payment System – Exempt Cancer Hospital Quality Reporting: (Finalized)
0213	Proportion admitted to the ICU in the last 30 days of life	Merit-Base Incentive Payment System (MIPS) Program (Finalized) Hospital Compare (Finalized) Prospective Payment System – Exempt Cancer Hospital Quality Reporting: (Finalized)
0215	Proportion not admitted to hospice	Merit-Base Incentive Payment System (MIPS) Program (Finalized) Hospital Compare (Finalized) Prospective Payment System – Exempt Cancer Hospital Quality Reporting: (Finalized)
0216	Proportion admitted to hospice for less than 3 days	Merit-Base Incentive Payment System (MIPS) Program (Finalized) Hospital Compare (Finalized) Prospective Payment System – Exempt Cancer Hospital Quality Reporting: (Finalized)
1623	Bereaved Family Survey	N/A

NQF #	Title	Federal Programs: Finalized or Implemented as of January 13, 2020
1625	Hospitalized Patients Who Die an Expected Death with an ICD that Has Been Deactivated	N/A
2651	CAHPS Hospice Survey (Experience with Care): 8 PRO-PMs: (Hospice Team Communication; Getting Timely Care; Getting Emotional and Religious Support; Getting Hospice Training; Rating of the Hospice Care; Willingness to Recommend the Hospice; Treating Family Member with Respect; Getting Help for Symptoms)	Hospice Quality Reporting (Implemented)
3235	Hospice and Palliative Care Composite Process Measure—Comprehensive Assessment at Admission	Hospice Quality Reporting (Implemented)

Appendix C: Geriatrics and Palliative Care Standing Committee and NQF Staff

STANDING COMMITTEE

R. Sean Morrison, MD (Co-Chair)

Patty and Jay Baker National Palliative Care Center; Director, National Palliative Care Research Center;
Director, Hertzberg Palliative Care Institute, Icahn School of Medicine at Mount Sinai
New York, New York

Deborah Waldrop, PhD, LMSW, ACSW (Co-Chair)

Professor, University of Buffalo, School of Social Work
Buffalo, New York

Margie Atkinson, D Min, BCC

Director, Pastoral Care, Ethics and Palliative Care, Morton Plant Mease/Bay Care Health System
Palm Harbor, Florida

Sree Battu, MD

Senior Associate Consultant, Mayo Clinic
Beachwood, Ohio

Samira Beckwith, LCSW, FACHE, LHD

President and CEO, Hope HealthCare Services
Fort Myers, Florida

Amy J. Berman, BSN

Senior Program Officer, John A. Hartford Foundation
New York, New York

Cleanne Cass, DO, FAAHPM, FAFAP

Director of Community Care and Education, Hospice of Dayton
Dayton, Ohio

Marian Grant, DNP, CRNP

Senior Regulatory Advisor, Coalition to Transform Advanced Care (C-TAC)
Washington, DC

George Handzo, BCC, CSSBB

Director, Health Services Research and Quality, HealthCare Chaplaincy
Los Angeles, California

Arif H. Kamal, MD, MBA, MHS, FACP, FAAHPM

Physician Quality and Outcomes Officer, Duke Cancer Institute
Durham, North Carolina

Suzanne Johnson, MPH, RN

Chief Operating Officer, National Hospice and Palliative Care Organization
Alexandria, Virginia

Janice Knebl, DO, MBA, FACOI, FACP

Director and Chief, Center for Geriatrics, University of North Texas Health Science Center at Fort Worth
Fort Worth, Texas

Christopher Laxton, CAE

Executive Director, AMDA - The Society for Post-Acute and Long-Term Care Medicine
Columbia, Maryland

Katherine Lichtenberg, DO, MPH, FAAFP

Physician Director, Enhanced Personal Health Care, Anthem Blue Cross and Blue Shield
Saint Louis, Missouri

Kelly Michaelson, MD, MPH, FCCM, FAP

Professor of Pediatrics and Julia and David Uihlein Professor of Bioethics and Medical Humanities
Director, Center for Bioethics and Medical Humanities
Northwestern University Feinberg School of Medicine
Attending Physician, Ann and Robert H. Lurie Children's Hospital of Chicago
Chicago, Illinois

Douglas Nee, Pharm D, MS

Clinical Pharmacist, Self
San Diego, California

Laura Porter, MD

Medical Advisor and Senior Patient Advocate, Colon Cancer Alliance
Washington, District of Columbia

Lynn Reinke, PhD, ARNP, FAAN

Research Investigator/Nurse Practitioner, VA Puget Sound Health Care System
Seattle, Washington

Tracy Schroepfer, PhD, MSW

Associate Professor of Social Work, University of Wisconsin, Madison, School of Social Work
Madison, Wisconsin

Linda Schwimmer, JD

Attorney, President and CEO, New Jersey Health Care Quality Institute
Pennington, New Jersey

Christine Seel Ritchie, MD, MSPH

Professor of Medicine in Residence, Harris Fishbon Distinguished Professor for Clinical Translational Research
in Aging, University of California San Francisco, Jewish Home of San Francisco Center for Research on Aging
San Francisco, California

Janelle Shearer, RN, BSN, MA, CPHQ

Program Manager, Stratis Health
Bloomington, Minnesota

Karl Steinberg, MD, CMD, HMDC

Chief Medical Officer, Mariner Health Central; Medical Director, Life Care Center of Vista, Carlsbad by the Sea
Care Center, Hospice by the Sea
Oceanside, California

Paul E. Tatum, MD, MSPH, CMD, FAAHPM, AGSF

Associate Professor in the Division of Geriatrics and Palliative Medicine at the Dell Medical School, University
of Texas, Austin
Austin, Texas

Sarah Thirlwell, MSc, MSc(A), RN, CHPN, CHPCA, AOCNS

Supportive Care Director, H. Lee Moffitt Cancer Center and Research Institute Hospital, Inc.
Tampa, Florida

NQF STAFF

Kathleen Giblin, RN

Acting Senior Vice President, Quality Measurement

Apryl Clark, MHSA

Acting Vice President, Quality Measurement

Amy Moyer, MS, PMP

Director

Kathryn Goodwin, MS

Director

Hannah Ingber, MPH

Project Analyst

Ngozi Ihenacho, MPH

Project Analyst

Appendix D: Measure Specifications

	1623 Bereaved Family Survey
Steward	Department of Veterans Affairs / Hospice and Palliative Care
Description	This measure calculates the proportion of Veteran decedent's family members who rate overall satisfaction with the Veteran decedent's end-of-life care in an inpatient setting as "Excellent" versus "Very good", "good", "fair", or "poor".
Type	Outcome: PRO-PM
Data Source	Instrument-Based Data For 2a1.25 - Family reported data/survey. For 2a1.26 - Bereaved Family Survey
Level	Facility, Other
Setting	Inpatient/Hospital, Post-Acute Care
Numerator Statement	The numerator is comprised of completed surveys (at least 12 of 17 structured items completed), where the global item question has an optimal response. The global item question asks "Overall, how would you rate the care that [Veteran] received in the last month of life" and the possible answer choices are: Excellent, Very good, Good, Fair, or Poor. The optimal response is Excellent.
Numerator Details	Included are those patients included in the denominator with completed surveys (at least 12 of 17 structured items completed) that receive an optimal response on the global item question.
Denominator Statement	The denominator consists of all inpatient deaths for which a survey was completed (at least 12 of 17 structured items completed), excluding: 1) deaths within 24 hours of admission (unless the Veteran had a previous hospitalization in the last month of life); 2) deaths that occur in the Emergency Department (unless the Veteran had a prior hospitalization of at least 24 hours in the last 31 days of life); Additional exclusion criteria include: 1) Veterans for whom a family member knowledgeable about their care cannot be identified (determined by the family member's report); or contacted (no current contacts listed or no valid addresses on file); 2) absence of a working telephone available to the family member.
Denominator Details	<p>The purpose of this measure is to assess families' perceptions of the quality of care that Veterans received from the VA in the last month of life. The BFS consists of 19 items (17 structured and 2 open-ended). The BFS items were selected from a longer survey that was developed and validated with the support of a VA HSR&D Merit Award and have been approved for use by the Office of Management and Budget.</p> <p>Seventeen items in the survey have predefined response options and ask family members to rate aspects of the care that the Veteran received from the VA in the last month of life. These items cover areas of care such as communication, emotional and spiritual support. Two additional items are open-ended and give family members the opportunity to provide comments regarding the care the patient received.</p> <p>A growing body of research has underscored the degree to which end-of-life care in the United States needs to be improved. The challenges of end-of-life care are particularly significant in the U.S. Department of Veterans Affairs Health Care system because the VA provides care for an increasingly older population with multiple comorbid conditions. In FY2000, approximately 104,000 enrolled Veterans died in the U.S., and approximately 27,200 Veterans died in VA facilities. At least 30% of the Veterans are over age 65 now, and 46% will be over 65 by 2030. Therefore, it is clear that the number of deaths in VA facilities will increase substantially as the World War II and Korean War Veterans age. These demographic trends mean that, like other healthcare systems, the VA will face substantial challenges of providing care to Veterans near the end-of-life.</p> <p>The VA has addressed this challenge aggressively in the last 5 years, however the VA has not yet developed and implemented measures of the quality of end-of-life care it provides to Veterans. There are at least 3 reasons why adoption of a quality measurement tool is</p>

	1623 Bereaved Family Survey
	<p>essential. First, it would make it possible to define and compare the quality of end-of-life care at each VA facility and to identify opportunities for improvement. Second, facilities and VISNs (geographic service divisions within the VA system) would be able to monitor the effectiveness of efforts to improve care locally and nationally, and would enable monitoring of the impact of the Comprehensive End of Life Care Initiative, ensuring that expenditures are producing improvements in care. Third, it will help the VA to recognize those facilities that provide outstanding end-of-life care, so that successful processes and structures of care can be identified and disseminated throughout the VA.</p> <p>The BFS's 17 close-ended items ask family members to rate aspects of the care that the Veteran received from the VA in the last month of life. These items cover areas of care such as communication, emotional and spiritual support, pain management and personal care needs. Two additional items (not used in scoring) are open-ended and give family members the opportunity to provide comments regarding the care the patient received. The BFS has undergone extensive development and has been pilot-tested for all inpatient deaths in Q4FY2008 in seven VISNs (1,2,4,5,8,11, and 22). As of October 1, 2009, Q1FY2010, all inpatient deaths in all VISNs were included in the project.</p> <p>The indicator denominator is comprised of the number of Veterans who die in an inpatient VA facility (intensive care, acute care, hospice unit, nursing home care or community living center) for whom a survey is completed. Completed surveys are defined as those with at least 12 of the 17 structured items completed.</p>
Exclusions	<ul style="list-style-type: none"> - Veterans for whom a family member knowledgeable about their care cannot be identified (determined by family member's report) - Absence of a current address and/or working telephone number for a family member or emergency contact. - Deaths within 24 hours of admission without a prior hospitalization of last least 24 hours in the last 31 days of life. - Deaths that occur in the operating room during an outpatient procedure. - Deaths due to a suicide or accident - Surveys in which less than 12 items were answered. -
Exclusion details	Name, address, and phone number of patient's family member or emergency contact are required for determining exclusion. In addition, information regarding the patient's admission(s) during the last 31 days of life, and including length of stay are also required to determine exclusion.
Risk Adjustment	Statistical risk model
Stratification	Variables necessary to stratify the measure are VISN, facility, quarter, year, outcome. VISN refers to "Veterans Integrated Service Network" and is a geographic area of the country where a facility is located. Facility is the actual VA medical center or affiliated community living center where the Veteran died. Quarter is the 3 month time period in which the patient died. Year is the VA fiscal year (runs from Oct 1 to Sept 30). Outcome refers to whether or not a survey was completed.
Type Score	Rate/proportion better quality = higher score
Algorithm	<p>The purpose of this measure is to assess families' perceptions of the quality of care that Veterans received from the VA in the last month of life. The BFS consists of 19 items (17 structured and 2 open-ended). The BFS items were selected from a longer survey that was developed and validated with the support of a VA HSR&D Merit Award and have been approved for use by the Office of Management and Budget.</p> <p>Seventeen items in the survey have predefined response options and ask family members to rate aspects of the care that the Veteran received from the VA in the last month of life. These items cover areas of care such as communication, emotional and spiritual support.</p>

	1623 Bereaved Family Survey
	<p>Two additional items are open-ended and give family members the opportunity to provide comments regarding the care the patient received.</p> <p>A growing body of research has underscored the degree to which end-of-life care in the United States needs to be improved. The challenges of end-of-life care are particularly significant in the U.S. Department of Veterans Affairs Health Care system because the VA provides care for an increasingly older population with multiple comorbid conditions. In FY2000, approximately 104,000 enrolled Veterans died in the U.S., and approximately 27,200 Veterans died in VA facilities. At least 30% of the Veterans are over age 65 now, and 46% will be over 65 by 2030. Therefore, it is clear that the number of deaths in VA facilities will increase substantially as the World War II and Korean War Veterans age. These demographic trends mean that, like other healthcare systems, the VA will face substantial challenges of providing care to Veterans near the end-of-life.</p> <p>The VA has addressed this challenge aggressively in the last 5 years, however the VA has not yet developed and implemented measures of the quality of end-of-life care it provides to Veterans. There are at least 3 reasons why adoption of a quality measurement tool is essential. First, it would make it possible to define and compare the quality of end-of-life care at each VA facility and to identify opportunities for improvement. Second, facilities and VISNs (geographic service divisions within the VA system) would be able to monitor the effectiveness of efforts to improve care locally and nationally, and would enable monitoring of the impact of the Comprehensive End of Life Care Initiative, ensuring that expenditures are producing improvements in care. Third, it will help the VA to recognize those facilities that provide outstanding end-of-life care, so that successful processes and structures of care can be identified and disseminated throughout the VA.</p> <p>The BFS's 17 close-ended items ask family members to rate aspects of the care that the Veteran received from the VA in the last month of life. These items cover areas of care such as communication, emotional and spiritual support, pain management and personal care needs. Two additional items (not used in scoring) are open-ended and give family members the opportunity to provide comments regarding the care the patient received. The BFS has undergone extensive development and has been pilot-tested for all inpatient deaths in Q4FY2008 in seven VISNs (1,2,4,5,8,11, and 22). As of October 1, 2009, Q1FY2010, all inpatient deaths in all VISNs were included in the project.</p> <p>The 17 structured items of the Bereaved Family Survey are scored as either "1" (optimal response) or "0" (all other answer choices). A score of "1" indicates that the family member perceived that the care they and/or the Veteran received was the best possible care (Excellent). A score of "0" reflects all other possible responses (Very good, Good, Fair, Poor). Items are coded as missing if respondents cannot or refuse to answer the item. Thus, the score for each item can be expressed as a fraction corresponding to the number of families who reported that the Veteran received optimal care (numerator), divided by the number of valid, non-missing responses for that item (denominator). Similarly, the score for the 17-item survey is calculated based on the global question item (Overall, how would you rate the care received in the last month of life? - Excellent, Very Good, Good, Fair, Poor). The global item is scored as the # of optimal responses/# of valid, non-missing responses for all completed surveys (12 of 17 structured items answered). This scoring system produces a facility- or VISN-level score that reflects the proportion of Veterans who received the best possible care overall (BFS score) and in specific areas corresponding to BFS items (e.g. pain management, communication, personal care, etc).</p> <p>We then add nonresponse and patient case mix weights to the model. All adjusted scores are reported. 122841 141015 146971 135548</p>
Copyright / Disclaimer	<p>This material is based upon work supported (or supported in part) by the Department of Veterans Affairs, Veterans Health Administration, Office of Research and Development, HSR&D. Use or publication of any materials used in the Bereaved Family Survey is prohibited.</p>

Appendix E: Related and Competing Measures

Comparison of NQF 1623 and NQF 2651

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
Steward	Department of Veterans Affairs / Hospice and Palliative Care	Centers for Medicare and Medicaid Services
Description	This measure calculates the proportion of Veteran decedent's family members who rate overall satisfaction with the Veteran decedent's end-of-life care in an inpatient setting as "Excellent" versus "Very good", "good", "fair", or "poor".	<p>The measures submitted here are derived from the CAHPS® Hospice Survey, which is a 47-item standardized questionnaire and data collection methodology. The survey is intended to measure the care experiences of hospice patients and their primary caregivers. Respondents to the survey are the primary informal caregivers of patients who died under hospice care. These are typically family members but can be friends. The hospice identifies the primary informal caregiver from their administrative records. Data collection for sampled decedents/caregivers is initiated two months following the month of the decedent's death.</p> <p>The publicly reported measures described here include the following six multi-item measures.</p> <ul style="list-style-type: none"> • Hospice Team Communication • Getting Timely Care • Treating Family Member with Respect • Getting Emotional and Religious Support • Getting Help for Symptoms • Getting Hospice Training <p>In addition, there are two global rating items that are publicly-reported measures.</p> <ul style="list-style-type: none"> • Rating of the hospice care • Willingness to recommend the hospice <p>Below we list each multi-item measure and its constituent items, along with the two global rating items. Then we briefly provide some general background information about CAHPS surveys.</p> <p>List of CAHPS Hospice Survey Measures</p> <p>Multi-Item Measures</p>

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
		<p>Hospice Team Communication (Composed of 6 items)</p> <ul style="list-style-type: none"> + While your family member was in hospice care, how often did the hospice team keep you informed about when they would arrive to care for your family member? + While your family member was in hospice care, how often did the hospice team explain things in a way that was easy to understand? + How often did the hospice team listen carefully to you when you talked with them about problems with your family member's hospice care? + While your family member was in hospice care, how often did the hospice team keep you informed about your family member's condition? + While your family member was in hospice care, how often did the hospice team listen carefully to you? + While your family member was in hospice care, how often did anyone from the hospice team give you confusing or contradictory information about your family member's condition or care? <p>Getting Timely Care (Composed of 2 items)</p> <ul style="list-style-type: none"> + While your family member was in hospice care, when you or your family member asked for help from the hospice team, how often did you get help as soon as you needed it? + How often did you get the help you needed from the hospice team during evenings, weekends, or holidays? <p>Treating Family Member with Respect (Composed of 2 items)</p> <ul style="list-style-type: none"> + While your family member was in hospice care, how often did the hospice team treat your family member with dignity and respect? + While your family member was in hospice care, how often did you feel that the hospice team really cared about your family member? <p>Providing Emotional Support (Composed of 3 items)</p> <ul style="list-style-type: none"> + While your family member was in hospice care, how much emotional support did you get from the hospice team? + In the weeks after your family member died, how much emotional support did you get from the hospice team? + Support for religious or spiritual beliefs includes talking, praying, quiet time, or other ways of meeting your religious or spiritual needs. While your family member was in hospice care, how much support for your religious and spiritual beliefs did you get from the hospice team?

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
		<p>Getting Help for Symptoms (Composed of 4 items)</p> <ul style="list-style-type: none"> + Did your family member get as much help with pain as he or she needed? + How often did your family member get the help he or she needed for trouble breathing? + How often did your family member get the help he or she needed for trouble with constipation? + How often did your family member receive the help he or she needed from the hospice team for feelings of anxiety or sadness? <p>Getting Hospice Care Training (Composed of 5 items)</p> <ul style="list-style-type: none"> + Did the hospice team give you enough training about what side effects to watch for from pain medicine? + Did the hospice team give you the training you needed about if and when to give more pain medicine to your family member? + Did the hospice team give you the training you needed about how to help your family member if he or she had trouble breathing? + Did the hospice team give you the training you needed about what to do if your family member became restless or agitated? + Side effects of pain medicine include things like sleepiness. Did any member of the hospice team discuss side effects of pain medicine with your or your family member? <p>Global Rating Measures:</p> <p>In addition to the multi-item measures, there are two “global” ratings measures. These single-item measures provide families and patients looking for care with overall evaluations of the care provided by the hospice. The items are rating of hospice care and willingness to recommend the hospice.</p> <ul style="list-style-type: none"> + Rating of Hospice Care: Using any number from 0 to 10, where 0 is the worst hospice care possible and 10 is the best hospice care possible, what number would you use to rate your family member’s hospice care? + Willingness to Recommend Hospice: Would you recommend this hospice to your friends and family? <p>The CAHPS Hospice Survey is part of the CAHPS family of experience of care surveys. English and other translations of the survey are available at http://www.hospicecahpssurvey.org/en/survey-instruments/ . CMS initiated national implementation of the CAHPS Hospice Survey in 2015. Hospices meeting CMS eligibility criteria were required to administer the survey for a</p>

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
		<p>“dry run” for at least one month of sample from the first quarter of 2015. Beginning with the second quarter of 2015, hospices are required to participate on an ongoing monthly basis in order to receive their full Annual Payment Update from CMS. Information regarding survey content and national implementation requirements, including the latest versions of the survey instrument and standardized protocols for data collection and submission, are available at: http://www.hospicecahpsurvey.org/. Public reporting of the survey-based measures on Hospice Compare started in February 2018 (www.medicare.gov Choose find hospice care)</p> <p>A list of the CAHPS Hospice Survey measures, including the components of the multi-item measures can be found in Appendix A</p>
Type	Outcome: PRO-PM	Outcome: PRO-PM
Data Source	<p>Instrument-Based Data For 2a1.25 - Family reported data/survey.</p> <p>For 2a1.26 - Bereaved Family Survey</p> <p>Available in attached appendix at A.1 No data dictionary</p>	<p>Instrument-Based Data CAHPS Hospice Survey; please see S.16 for information regarding modes of data collection. The survey instrument is available in English, Spanish, Chinese, Russian, Portuguese, Vietnamese, Polish and Korean.</p> <p>Available at measure-specific web page URL identified in S.1 No data dictionary</p>
Level	Facility, Other	Facility
Setting	Inpatient/Hospital, Post-Acute Care	Other
Numerator Statement	The numerator is comprised of completed surveys (at least 12 of 17 structured items completed), where the global item question has an optimal response. The global item question asks "Overall, how would you rate the care that [Veteran] received in the last month of life" and the possible answer choices are: Excellent, Very good, Good, Fair, or Poor. The optimal response is Excellent.	CMS calculates CAHPS Hospice Survey measure scores using top-, middle- and bottom- box scoring. The top-box score refers to the percentage of caregiver respondents that give the most positive response(s). The bottom box score refers to the percentage of caregiver respondents that give the least positive response(s). The middle box is the proportion remaining after the top and bottom boxes have been calculated; see below for details. Details regarding the definition of most and least positive response(s) are noted in Section S.5 below.
Numerator Details	Included are those patients included in the denominator with completed surveys (at least 12 of 17 structured items completed) that receive an optimal response on the global item question.	<p>For each survey item, the top and bottom box numerators are the number of respondents who selected the most and least positive response category(ies), respectively, as follows:</p> <p>For items using a “Never/Sometimes/Usually/Always” response scale, the top box numerator is the number of respondents who answer “Always” and the bottom box numerator is the number of respondents who answer “Never” or “Sometimes.” The one exception to this guidance is for the Q10</p>

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
		<p>“While your family member was in hospice care, how often did anyone from the hospice team give you confusing or contradictory information about your family member’s condition or care?” For this item, the top box numerator is the number of respondents who answer “Never” and the bottom box numerator is the number of respondents who answer “Always” or “Usually.”</p> <p>For items using a “Yes, definitely/Yes, somewhat/No” response scale, the top box numerator is the number of respondents who answer “Yes, definitely” and the bottom box numerator is the number of respondents who answer “No.”</p> <p>For items using a “Too Little/Right Amount/Too Much” response scale, the top box numerator is the number of respondents who answer “Right Amount” and the bottom box numerator is the number of respondents who answer “Too little” or “Too much.” (There is no middle box for items using this response scale.)</p> <p>The top box numerator for the Rating of Hospice item is the number of respondents who answer 9 or 10 for the item (on a scale of 0 to 10, where 10 is the “Best Hospice Care Possible”); the bottom box numerator is the number of respondents who answer 0 to 6.</p> <p>The top box numerator for the Willingness to Recommend item is the number of respondents who answer “Definitely Yes” (on a scale of “Definitely No/Probably No/Probably Yes/Definitely Yes”); the bottom box numerator is the number of respondents who answer “Probably No” or “Definitely No.”</p> <p>Calculation of hospice-level multi-item measures</p> <ol style="list-style-type: none"> 0. Score each item using top- box method, possible values of 0 or 100 1. Calculate mode- adjusted scores for each item for each respondent 2. Calculate case-mix adjusted scores for each item for each hospice 3. Take the unweighted means of the mode- and case-mix-adjusted hospice-level items to form multi-item measures <p>Here is an example of calculations for the measure “Getting Timely Care.”</p> <ol style="list-style-type: none"> 0. Score each item using top box method, possible values of 0 or 100 <p>Both items in “Getting Care Quickly” have four response options: Never, Sometimes, Usually, Always. Recode each item as 100 for “Always” and 0 for “Never”, “Sometimes”, or “Usually”.</p>

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
		<p>Item #1. While your family member was in hospice care, when you or your family member asked for help from the hospice team, how often did you get help as soon as you needed it?</p> <p>Item #2. How often did you get the help you needed from the hospice team during evenings, weekends, or holidays?</p> <ol style="list-style-type: none"> 1. Calculate mode-adjusted scores for each item for each respondent 2. Calculate case-mix adjusted scores for each item for each hospice <p>Each item is case mix adjusted separately; this step produces case-mix adjusted item-level scores for each hospice.</p> <ol style="list-style-type: none"> 3. Take the unweighted means of the case-mix adjusted hospice-level items to form multi-item measures. <p>If the case-mix adjusted scores for a hospice are 95 for item #1 and 90 for item #2, then the hospice-level 'Getting Timely Care' would be calculated as $(\text{Item1} + \text{Item2}) / 2 = (95 + 90) / 2 = 92.5$.</p>
Denominator Statement	<p>The denominator consists of all inpatient deaths for which a survey was completed (at least 12 of 17 structured items completed), excluding: 1) deaths within 24 hours of admission (unless the Veteran had a previous hospitalization in the last month of life); 2) deaths that occur in the Emergency Department (unless the Veteran had a prior hospitalization of at least 24 hours in the last 31 days of life); Additional exclusion criteria include: 1) Veterans for whom a family member knowledgeable about their care cannot be identified (determined by the family member's report); or contacted (no current contacts listed or no valid addresses on file); 2) absence of a working telephone available to the family member.</p>	<p>CAHPS® Hospice Survey measure scores are calculated only for hospices that had at least 30 completed questionnaires over the most recent eight quarters of data collection.</p> <p>The target population for the survey are the adult primary caregivers of hospice decedents. Respondent eligibility and exclusions are defined in detail in the sections that follow. A survey is defined as completed when at least 50 percent of the questions applicable to all decedents/caregivers are answered (Questions 1 – 4, 6 – 13, 15, 17, 21, 24, 26, 28, 30 – 32, and 35 – 47). The survey uses screener questions to identify respondents eligible to respond to subsequent items. Therefore, denominators vary by survey item (and corresponding multi-item measures, if applicable) according to the eligibility of respondents for each item. In addition, for the Getting Hospice Care Training measure, scores are calculated only among those respondents who indicate that their family member received hospice care at home or in an assisted living facility.</p>
Denominator Details	<p>The purpose of this measure is to assess families' perceptions of the quality of care that Veterans received from the VA in the last month of life. The BFS consists of 19 items (17 structured and 2 open-ended). The BFS items were selected from a longer survey that was developed and validated with the support of a VA</p>	<p>For each item in a multi-item measure, as well as for the ratings measures, the top box denominator is the number of respondents per hospice who answered the item. For each multi-item measure score, the denominator is the number of respondents who answer at least one item within the multi-item measure. Multi-item measure scores are the average proportion of</p>

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
	<p>HSR&D Merit Award and have been approved for use by the Office of Management and Budget.</p> <p>Seventeen items in the survey have predefined response options and ask family members to rate aspects of the care that the Veteran received from the VA in the last month of life. These items cover areas of care such as communication, emotional and spiritual support. Two additional items are open-ended and give family members the opportunity to provide comments regarding the care the patient received.</p> <p>A growing body of research has underscored the degree to which end-of-life care in the United States needs to be improved. The challenges of end-of-life care are particularly significant in the U.S. Department of Veterans Affairs Health Care system because the VA provides care for an increasingly older population with multiple comorbid conditions. In FY2000, approximately 104,000 enrolled Veterans died in the U.S., and approximately 27,200 Veterans died in VA facilities. At least 30% of the Veterans are over age 65 now, and 46% will be over 65 by 2030. Therefore, it is clear that the number of deaths in VA facilities will increase substantially as the World War II and Korean War Veterans age. These demographic trends mean that, like other healthcare systems, the VA will face substantial challenges of providing care to Veterans near the end-of-life.</p> <p>The VA has addressed this challenge aggressively in the last 5 years, however the VA has not yet developed and implemented measures of the quality of end-of-life care it provides to Veterans. There are at least 3 reasons why adoption of a quality measurement tool is essential. First, it would make it possible to define and compare the quality of end-of-life care at each VA facility and to identify opportunities for improvement. Second, facilities and VISNs (geographic service divisions within the VA system) would be able to monitor the</p>	<p>respondents that gave responses in the most positive category across the items in the multi-item measure (as discussed in S.6).</p> <p>Survey population: Primary caregivers of patients who died while receiving care from a given hospice in a given month.</p> <p>Denominator for Multi-Item Measures: The number of respondents who answer at least one item within the multi-item measure.</p> <p>Denominator for Rating Measures: The number of respondents who answered the item.</p>

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
	<p>effectiveness of efforts to improve care locally and nationally, and would enable monitoring of the impact of the Comprehensive End of Life Care Initiative, ensuring that expenditures are producing improvements in care. Third, it will help the VA to recognize those facilities that provide outstanding end-of-life care, so that successful processes and structures of care can be identified and disseminated throughout the VA.</p> <p>The BFS's 17 close-ended items ask family members to rate aspects of the care that the Veteran received from the VA in the last month of life. These items cover areas of care such as communication, emotional and spiritual support, pain management and personal care needs. Two additional items (not used in scoring) are open-ended and give family members the opportunity to provide comments regarding the care the patient received. The BFS has undergone extensive development and has been pilot-tested for all inpatient deaths in Q4FY2008 in seven VISNs (1,2,4,5,8,11, and 22). As of October 1, 2009, Q1FY2010, all inpatient deaths in all VISNs were included in the project.</p> <p>The indicator denominator is comprised of the number of Veterans who die in an inpatient VA facility (intensive care, acute care, hospice unit, nursing home care or community living center) for whom a survey is completed. Completed surveys are defined as those with at least 12 of the 17 structured items completed.</p>	
Exclusions	<ul style="list-style-type: none"> - Veterans for whom a family member knowledgeable about their care cannot be identified (determined by family member's report) - Absence of a current address and/or working telephone number for a family member or emergency contact. - Deaths within 24 hours of admission without a prior hospitalization of last least 24 hours in the last 31 days of life. 	<p>The eight measures included here are calculated only for hospices that have at least 30 completed surveys over eight quarters of data collection.</p> <p>The exclusions noted in here are those who are ineligible to participate in the survey. The one exception is caregivers who report on the survey that they “never” oversaw or took part in the decedent’s care; these respondents are instructed to complete the “About You” and “About Your Family Member” sections of the survey only.</p> <p>Cases are excluded from the survey target population if:</p> <ul style="list-style-type: none"> • The hospice patient is still alive

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
	<ul style="list-style-type: none"> - Deaths that occur in the operating room during an outpatient procedure. - Deaths due to a suicide or accident - Surveys in which less than 12 items were answered. - 	<ul style="list-style-type: none"> • The decedent's age at death was less than 18 • The decedent died within 48 hours of his/her last admission to hospice care • The decedent had no caregiver of record • The decedent had a caregiver of record, but the caregiver does not have a U.S. or U.S. Territory home address • The decedent had no caregiver other than a nonfamilial legal guardian • The decedent or caregiver requested that they not be contacted (i.e., by signing a no publicity request while under the care of hospice or otherwise directly requesting not to be contacted) • The caregiver is institutionalized, has mental/physical incapacity, has a language barrier, or is deceased • The caregiver reports on the survey that he or she "never" oversaw or took part in decedent's hospice care
Exclusion Details	Name, address, and phone number of patient's family member or emergency contact are required for determining exclusion. In addition, information regarding the patient's admission(s) during the last 31 days of life, and including length of stay are also required to determine exclusion.	Please see S.10.The CAHPS Hospice Survey Quality Assurance Guidelines (available at: http://www.hospicecahpsurvey.org/en/quality-assurance-guidelines/) contain detailed information regarding how to code decedent/caregiver cases, and how to code appropriately and inappropriately skipped items, as well as items with multiple responses.
Risk Adjustment	Statistical risk model	Statistical risk model
Stratification	Variables necessary to stratify the measure are VISN, facility, quarter, year, outcome. VISN refers to "Veterans Integrated Service Network" and is a geographic area of the country where a facility is located. Facility is the actual VA medical center or affiliated community living center where the Veteran died. Quarter is the 3 month time period in which the patient died. Year is the VA fiscal year (runs from Oct 1 to Sept 30). Outcome refers to whether or not a survey was completed.	CAHPS Hospice Survey measure scores are used for reporting at the hospice-level (i.e., not stratified by region or other characteristics).
Type Score	Rate/proportion better quality = higher score	Rate/proportion better quality = higher score

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
Algorithm	<p>The purpose of this measure is to assess families' perceptions of the quality of care that Veterans received from the VA in the last month of life. The BFS consists of 19 items (17 structured and 2 open-ended). The BFS items were selected from a longer survey that was developed and validated with the support of a VA HSR&D Merit Award and have been approved for use by the Office of Management and Budget.</p> <p>Seventeen items in the survey have predefined response options and ask family members to rate aspects of the care that the Veteran received from the VA in the last month of life. These items cover areas of care such as communication, emotional and spiritual support. Two additional items are open-ended and give family members the opportunity to provide comments regarding the care the patient received.</p> <p>A growing body of research has underscored the degree to which end-of-life care in the United States needs to be improved. The challenges of end-of-life care are particularly significant in the U.S. Department of Veterans Affairs Health Care system because the VA provides care for an increasingly older population with multiple comorbid conditions. In FY2000, approximately 104,000 enrolled Veterans died in the U.S., and approximately 27,200 Veterans died in VA facilities. At least 30% of the Veterans are over age 65 now, and 46% will be over 65 by 2030. Therefore, it is clear that the number of deaths in VA facilities will increase substantially as the World War II and Korean War Veterans age. These demographic trends mean that, like other healthcare systems, the VA will face substantial challenges of providing care to Veterans near the end-of-life.</p> <p>The VA has addressed this challenge aggressively in the last 5 years, however the VA has not yet developed and implemented measures of the quality of end-of-life care it provides to Veterans. There are at least 3 reasons why</p>	<p>Top Box Score Calculation:</p> <ol style="list-style-type: none"> 1) Identify target respondent population (i.e., primary caregivers of hospice patients who died while receiving hospice care from a given hospice in a given month) 2) Identify any exclusions from the respondent population (as described above in S.10) 3) Score each item using top box method, possible values of 0 or 100 4) Calculate mode adjusted top box scores for each item. 5) Calculate case-mix adjusted top box scores for each item for each hospice; case-mix adjustment is a linear regression based approach that adjusts for all variables listed in S.14. Specifically, a regression model predicting item scores is fit using the case-mix adjustor variables and fixed effects for hospices. Adjusted hospice means are then calculated (e.g., using LSMEANS in SAS). 6) Top-box scores are averaged across the items within each multi-item measure, weighting each item equally. If data are missing for a respondent for an item(s) within a multi-item measure, the respondent's answers to other items within the measure are still used in the calculation of multi-item measure scores. (Please see S.22 below for more details). <p>Top Box Score Calculation:</p> <ol style="list-style-type: none"> 1) Identify target respondent population (i.e., primary caregivers of hospice patients who died while receiving hospice care from a given hospice in a given month) 2) Identify any exclusions from the respondent population (as described above in S.10) 3) Score each item using top box method, possible values of 0 or 100 4) Calculate mode adjusted top box scores for each item. 5) Calculate case-mix adjusted top box scores for each item for each hospice; case-mix adjustment is a linear regression based approach that adjusts for all variables listed in S.14. Specifically, a regression model predicting item scores is fit using the case-mix adjustor variables and fixed effects for hospices. Adjusted hospice means are then calculated (e.g., using LSMEANS in SAS). 6) Top-box scores are averaged across the items within each multi-item measure, weighting each item equally. If data are missing for a respondent

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
	<p>adoption of a quality measurement tool is essential. First, it would make it possible to define and compare the quality of end-of-life care at each VA facility and to identify opportunities for improvement. Second, facilities and VISNs (geographic service divisions within the VA system) would be able to monitor the effectiveness of efforts to improve care locally and nationally, and would enable monitoring of the impact of the Comprehensive End of Life Care Initiative, ensuring that expenditures are producing improvements in care. Third, it will help the VA to recognize those facilities that provide outstanding end-of-life care, so that successful processes and structures of care can be identified and disseminated throughout the VA.</p> <p>The BFS's 17 close-ended items ask family members to rate aspects of the care that the Veteran received from the VA in the last month of life. These items cover areas of care such as communication, emotional and spiritual support, pain management and personal care needs. Two additional items (not used in scoring) are open-ended and give family members the opportunity to provide comments regarding the care the patient received. The BFS has undergone extensive development and has been pilot-tested for all inpatient deaths in Q4FY2008 in seven VISNs (1,2,4,5,8,11, and 22). As of October 1, 2009, Q1FY2010, all inpatient deaths in all VISNs were included in the project.</p> <p>The 17 structured items of the Bereaved Family Survey are scored as either "1" (optimal response) or "0" (all other answer choices). A score of "1" indicates that the family member perceived that the care they and/or the Veteran received was the best possible care (Excellent). A score of "0" reflects all other possible responses (Very good, Good, Fair, Poor). Items are coded as missing if respondents cannot or refuse to answer the item. Thus, the score for each item can be expressed as a fraction corresponding to the number of families who reported</p>	<p>for an item(s) within a multi-item measure, the respondent's answers to other items within the measure are still used in the calculation of multi-item measure scores. (Please see S.22 below for more details).</p>

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
	<p>that the Veteran received optimal care (numerator), divided by the number of valid, non-missing responses for that item (denominator). Similarly, the score for the 17-item survey is calculated based on the global question item (Overall, how would you rate the care received in the last month of life? - Excellent, Very Good, Good, Fair, Poor). The global item is scored as the # of optimal responses/# of valid, non missing responses for all completed surveys (12 of 17 structured items answered). This scoring system produces a facility- or VISN-level score that reflects the proportion of Veterans who received the best possible care overall (BFS score) and in specific areas corresponding to BFS items (e.g. pain management, communication, personal care, etc).</p> <p>We then add nonresponse and patient case mix weights to the model. All adjusted scores are reported. The purpose of this measure is to assess families' perceptions of the quality of care that Veterans received from the VA in the last month of life. The BFS consists of 19 items (17 structured and 2 open-ended). The BFS items were selected from a longer survey that was developed and validated with the support of a VA HSR&D Merit Award and have been approved for use by the Office of Management and Budget.</p> <p>Seventeen items in the survey have predefined response options and ask family members to rate aspects of the care that the Veteran received from the VA in the last month of life. These items cover areas of care such as communication, emotional and spiritual support. Two additional items are open-ended and give family members the opportunity to provide comments regarding the care the patient received.</p> <p>A growing body of research has underscored the degree to which end-of-life care in the United States needs to be improved. The challenges of end-of-life care are particularly significant in the U.S. Department of Veterans Affairs Health Care system because the VA</p>	

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
	<p>provides care for an increasingly older population with multiple comorbid conditions. In FY2000, approximately 104,000 enrolled Veterans died in the U.S., and approximately 27,200 Veterans died in VA facilities. At least 30% of the Veterans are over age 65 now, and 46% will be over 65 by 2030. Therefore, it is clear that the number of deaths in VA facilities will increase substantially as the World War II and Korean War Veterans age. These demographic trends mean that, like other healthcare systems, the VA will face substantial challenges of providing care to Veterans near the end-of-life.</p> <p>The VA has addressed this challenge aggressively in the last 5 year, however the VA has not yet developed and implemented measures of the quality of end-of-life care it provides to Veterans. There are at least 3 reasons why adoption of a quality measurement tool is essential. First, it would make it possible to define and compare the quality of end-of-life care at each VA facility and to identify opportunities for improvement. Second, facilities and VISNs (geographic service divisions within the VA system) would be able to monitor the effectiveness of efforts to improve care locally and nationally, and would enable monitoring of the impact of the Comprehensive End of Life Care Initiative, ensuring that expenditures are producing improvements in care. Third, it will help the VA to recognize those facilities that provide outstanding end-of-life care, so that successful processes and structures of care can be identified and disseminated throughout the VA.</p> <p>The BFS's 17 close-ended items ask family members to rate aspects of the care that the Veteran received from the VA in the last month of life. These items cover areas of care such as communication, emotional and spiritual support, pain management and personal care needs. Two additional items (not used in scoring) are open-ended and give family members the opportunity to</p>	

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
	<p>provide comments regarding the care the patient received. The BFS has undergone extensive development and has been pilot-tested for all inpatient deaths in Q4FY2008 in seven VISNs (1,2,4,5,8,11, and 22). As of October 1, 2009, Q1FY2010, all inpatient deaths in all VISNs were included in the project.</p> <p>The 17 structured items of the Bereaved Family Survey are scored as either "1" (optimal response) or "0" (all other answer choices). A score of "1" indicates that the family member perceived that the care they and/or the Veteran received was the best possible care (Excellent). A score of "0" reflects all other possible responses (Very good, Good, Fair, Poor). Items are coded as missing if respondents cannot or refuse to answer the item. Thus, the score for each item can be expressed as a fraction corresponding to the number of families who reported that the Veteran received optimal care (numerator), divided by the number of valid, non-missing responses for that item (denominator). Similarly, the score for the 17-item survey is calculated based on the global question item (Overall, how would you rate the care received in the last month of life? - Excellent, Very Good, Good, Fair, Poor). The global item is scored as the # of optimal responses/# of valid, non missing responses for all completed surveys (12 of 17 structured items answered). This scoring system produces a facility- or VISN-level score that reflects the proportion of Veterans who received the best possible care overall (BFS score) and in specific areas corresponding to BFS items (e.g. pain management, communication, personal care, etc). We then add nonresponse and patient case mix weights to the model. All adjusted scores are reported.</p>	
Submission items	<p>5.1 Identified measures: 2651 : CAHPS® Hospice Survey (experience with care)</p> <p>5a.1 Are specs completely harmonized? No</p>	<p>5.1 Identified measures: 0208 : Family Evaluation of Hospice Care 1623 : Bereaved Family Survey</p> <p>5a.1 Are specs completely harmonized? Yes</p>

	1623: Bereaved Family Survey	2651: CAHPS® Hospice Survey (experience with care)
	<p>5a.2 If not completely harmonized, identify difference, rationale, impact: Survey items different as well as coding of items, Target group is also different, We are specifically looking at inpatient Veteran deaths, regardless of hospice use. Currently, the BFS is the only tool assessing end of life care in a VA inpatient setting. We believe that assessing all deaths, not just hospice deaths, is critical to the VA mission of improving care for all Veterans regardless of choice of level of care at death. We do see any negative impact to interpretability or burden of data collection.</p> <p>5b.1 If competing, why superior or rationale for additive value: NQF 2651 CAHPS Hospice Survey</p> <p>Although the Bereaved Family Survey is in many ways similar to the CAHPS Hospice Survey, it provides information on a specific population (Veterans) and measures the quality of care provided a single health care system. Unlike the CAHPS-Hospice, the BFS provides a coherent measurement strategy that allows comparisons across systems of care and sites of death in a single health care system. This measure assesses the quality of care of the largest unified health care system in the United States and cares for more than 5 million patients annually. Because it is a unified health system, the VA is uniquely situated to make use of the quality data that can be easily and quickly disseminated. The BFS also measures satisfaction of care that are unique to a Veteran population (i.e, survivor and funeral benefits, PTSD). The population of Veterans and families that the VA serves is unique in several key respects: 1) Veterans and their families may face different challenges at the end of life than non-Veterans do. The costs of hospitalization are less likely to be relevant to non-VA populations.</p>	<p>5a.2 If not completely harmonized, identify difference, rationale, impact: N/A</p> <p>5b.1 If competing, why superior or rationale for additive value: 0208 Family Evaluation of Hospice Care.</p> <p>The Family Evaluation of Hospice Care Survey (FEHC), developed more than 20 years ago, assesses hospice care experiences from the perspective of bereaved family members. The CAHPS Hospice Survey covers similar domains, but includes important methodological improvements in the response task, and is adjusted for case mix and mode. Additionally, more stringent survey administration guidelines are in place to permit public reporting of the survey results and valid comparison across hospice programs. FEHC measures were maintained by the National Hospice and Palliative Care Organization (NHPCO), which operated a voluntary repository that provided hospice programs with national benchmarks for FEHC measures. With the national implementation of the CAHPS Hospice Survey, NHPCO shut down the voluntary repository. NQF endorsement of FEHC measures was removed in January 2018.</p> <p>1623 Bereaved Family Survey.</p> <p>The Department of Veterans Affairs Bereaved Family Survey assesses experiences of veterans' health care in the last month of life from the perspective of bereaved family members. Importantly, the Bereaved Family Survey assesses care for those who die in inpatient settings, regardless of whether they have received hospice care; this is distinct from respondents to the CAHPS Hospice Survey, who include informal caregivers of decedents who received hospice care across a range of care settings (including both inpatient and other settings).</p>

Appendix F: Pre-Evaluation Comments

No NQF member comments were received during the pre-commenting period.

National Quality Forum
1099 14th Street NW, Suite 500
Washington, DC 20005
<http://www.qualityforum.org>

ISBN [Assigned by Communications]
©2019 National Quality Forum