

# NQF-Endorsed Measures for Infectious Disease

DRAFT REPORT FOR VOTING

June 14, 2017



NATIONAL  
QUALITY FORUM

This report is funded by the Department of Health  
and Human Services under contract HHSM-500-  
2012-00009I Task Order HHSM-500-T0000



## Contents

<b>Executive Summary</b> .....	<b>5</b>
<b>Introduction</b> .....	<b>6</b>
HIV/AIDS.....	6
Sepsis and Septic Shock .....	6
Trends and Performance.....	7
<b>NQF Portfolio of Performance Measures for Infectious Disease Conditions</b> .....	<b>8</b>
Table 1. NQF Infectious Disease Portfolio of Measures.....	8
National Quality Strategy .....	8
Use of Measures in the Portfolio .....	9
Improving NQF’s Infectious Disease Portfolio .....	9
<b>Infectious Disease Measure Evaluation</b> .....	<b>9</b>
Table 2. Infectious Disease Measure Evaluation Summary .....	10
Comments Received Prior to Committee Evaluation .....	10
Overarching Issues .....	10
Summary of Measure Evaluation .....	13
<b>References</b> .....	<b>17</b>
<b>Appendix A: Details of Measure Evaluation</b> .....	<b>19</b>
Measures Recommended .....	19
2082 HIV Viral Load Suppression .....	19
3210 HIV Viral Load Suppression .....	22
2079 HIV Medical Visit Frequency .....	24
3209 HIV Medical Visit Frequency .....	27
2080 Gap in HIV Medical Visits .....	29
2083 Prescription of HIV Antiretroviral Therapy .....	32
3211 Prescription of HIV Antiretroviral Therapy .....	34
3215 Adult Inpatient Risk Adjusted Sepsis Mortality .....	36
0500 Severe Sepsis and Septic Shock: Management Bundle.....	40
<b>Measures Withdrawn from Consideration</b> .....	<b>54</b>
<b>Appendix B: NQF Infectious Disease Portfolio and Related Measures</b> .....	<b>55</b>
<b>Appendix C: Infectious Disease Portfolio—Use in Federal Programs</b> .....	<b>58</b>
<b>Appendix D: Project Standing Committee and NQF Staff</b> .....	<b>59</b>
<b>Appendix E: Measure Specifications</b> .....	<b>61</b>
2079 HIV Medical visit frequency .....	61
2080 Gaps in HIV medical visits.....	62

3209 HIV Medical visit frequency .....	63
2082 HIV Viral load suppression .....	66
3210 HIV Viral load suppression .....	67
2083 Prescription of HIV Antiretroviral Therapy .....	69
3211 Prescription of HIV Antiretroviral Therapy .....	70
0500 Severe Sepsis and Septic Shock Management Bundle .....	72
3215 Adult Inpatient Risk Adjusted Sepsis Mortality .....	76
<b>Appendix F: Related and Competing Measures .....</b>	<b>80</b>
<b>Appendix G: Pre-Evaluation Comments .....</b>	<b>100</b>

# Infectious Disease

## DRAFT REPORT

### Executive Summary

Many infectious diseases have been controlled or eradicated through the use of vaccines and advanced medicine, infectious disease continues to cause widespread morbidity and mortality, and rising health care costs. However, infectious disease persists, from viruses and bacteria that cause respiratory illnesses to sexually transmitted infections (STI) and more.

The Infectious Disease portfolio currently has nine endorsed measures for infectious disease addressing sepsis and septic shock, HIV/AIDS, and respiratory conditions. [Appendix B](#) details the full portfolio of infectious disease measures.

For this project, the Standing Committee evaluated four newly-submitted measures and five measures undergoing maintenance review against NQF's standard evaluation criteria. All nine measures were recommended for endorsement. The nine measures recommended by the Standing Committee are:

- 2082: HIV Viral Load Suppression (Health Resources and Services Administration - HIV/AIDS Bureau)
- 3210: HIV Viral Load Suppression – Legacy eMeasure (Health Resources and Services Administration - HIV/AIDS Bureau)
- 2079: HIV Medical Visit Frequency (Health Resources and Services Administration - HIV/AIDS Bureau)
- 3209: HIV Medical Visit Frequency – Legacy eMeasure (Health Resources and Services Administration - HIV/AIDS Bureau)
- 2080: Gap in HIV Medical Visit (Health Resources and Services Administration - HIV/AIDS Bureau)
- 2083: Prescription of HIV Antiretroviral Therapy (Health Resources and Services Administration - HIV/AIDS Bureau)
- 3211: Prescription of HIV Antiretroviral Therapy – Legacy eMeasure (Health Resources and Services Administration - HIV/AIDS Bureau)
- 0500: Severe Sepsis and Septic Shock: Management Bundle (Henry Ford Hospital)
- 3215: Adult Inpatient Risk Adjusted Sepsis Mortality (New York State Department of Health, Office of Quality and Patient Safety)

Brief summaries of the measures currently under review are included in the body of the report; detailed summaries of the Committee's discussion and ratings of the criteria for each measure are in [Appendix A](#).

## Introduction

Many infectious diseases have been controlled or eradicated through the use of vaccines and advanced medicine, infectious disease continues to cause widespread morbidity and mortality, and rising health care costs. However, infectious disease persist, from viruses and bacteria that cause respiratory illnesses to sexually transmitted infections (STI) and more. Specifically:

- Infectious diseases account for 3.9 million hospital visits per year and are a leading cause of death in the United States<sup>1</sup>;
- Each year, the nation spends more than \$120 and \$5 billion to treat infectious diseases and antibiotic resistant bacteria, respectively<sup>2</sup>;
- Septicemia is the most expensive condition treated in US hospitals at \$20.3 billion in 2011.<sup>3</sup>

Effective quality measures are critically important to improve national efforts to advance treatment of infectious disease and improve patient safety and healthcare outcomes. Providing resources, such as patient education and intervention programs along with continued scientific research for existing and emerging diseases, will reduce mortality and healthcare costs.

NQF has endorsed a number of consensus standards to evaluate the quality of care for topic areas related to infectious disease over the past decade. As quality measurement has matured, better data systems have become available, electronic health records are closer to widespread adoption, and the demand for meaningful performance measures has prompted development of more sophisticated measures of healthcare processes and outcomes for infectious disease conditions. An evaluation of the NQF endorsed infectious disease measures and consideration of new measures will ensure the currency of NQF's portfolio of voluntary consensus standards. This measurement cycle focused on measures for HIV/AIDS and Sepsis.

## HIV/AIDS

HIV (human immunodeficiency virus) is an infectious disease that attacks the immune system; if left untreated, HIV will progress to AIDS.<sup>4</sup> Fortunately, advances in care and treatment have shown that people living with HIV (PLWH) who are stable on antiretroviral medication are more likely to achieve viral suppression and are therefore less likely to transmit the virus to others.<sup>5</sup> Quality measures that allow for the monitoring of client engagement and retention in care, and the assessment of health outcomes such as viral load, are needed to reduce the incidence and prevalence of HIV infection and related morbidity and mortality.

## Sepsis and Septic Shock

Sepsis is a complication caused by the body's responses to infection.<sup>6</sup> According to the Centers for Disease Control and Prevention (CDC), sepsis can occur to anyone, at any time, from any type of infection, and can affect any part of the body.<sup>7</sup> Clinicians have identified three stages of sepsis based on the severity of the infection and the body's responses:

- Sepsis: the infection reaches the bloodstream and causes inflammation in the body.

- Severe sepsis: the infection is severe enough to affect the organs, such as the heart, brain, and kidneys.
- Septic shock: a significant drop in blood pressure that can lead to respiratory or heart failure, stroke, failure of other organs, and death.

Every year, severe sepsis strikes more than a million Americans.<sup>9</sup> Experts estimate that between 28 and 50 percent of patients with sepsis die—far more than the number of U.S. deaths from prostate cancer, breast cancer and AIDS combined.<sup>10</sup> The number of sepsis cases per year is likely rising due to a combination of factors, including increased awareness and tracking of the condition, an aging population, the increased longevity of people with chronic diseases, the spread of antibiotic-resistant organisms, an upsurge in invasive procedures and broader use of immunosuppressive and chemotherapeutic agents.<sup>11</sup>

The Agency for Healthcare Research and Quality (AHRQ) lists sepsis as the most expensive condition treated in U.S. hospitals, costing more than \$20 billion in 2011.<sup>12</sup> Readmission due to sepsis is two to three times more likely—and two or three times more costly—than readmission resulting from many other conditions, including heart failure, pneumonia and chronic obstructive pulmonary disease.<sup>13</sup>

## Trends and Performance

Approximately 1.2 million people are living with HIV in the United States today; about 87 percent of them are aware that they are infected.<sup>14</sup> In 2015, the Centers for Disease Control and Prevention (CDC) released an analysis showing that HIV diagnoses fell by 19 percent over the past decade.<sup>15</sup> This decline was driven by a decrease in HIV diagnoses among heterosexuals, people who inject drugs, and African American women and heterosexual men.<sup>16</sup> The CDC also reported that, thanks to sustained testing efforts, the proportion of Americans with HIV who know their status has reached an all-time high of 87 percent.<sup>17</sup> The implementation of the Affordable Care Act (ACA) continued to increase access to critical HIV testing, prevention, and care services nationwide.<sup>18</sup>

Recent scientific advances to stop HIV include improved testing techniques, early treatment with antiretroviral medications, and pre-exposure prophylaxis. Unfortunately, these advances are not reaching many people in need. According to the CDC, surveillance data show 57 percent of people diagnosed with HIV are receiving care for their infection, and only 55 percent have their virus suppressed through treatment.<sup>19</sup> The CDC also reports that there are substantial gaps between Southern states and the rest of the country on death rates among people with diagnosed HIV and knowledge of HIV – positive status.<sup>20</sup>

In addition to geographic disparities, progress has been uneven among racial, ethnic, gay, and bisexual men. The CDC's analysis found that HIV diagnoses dropped 18 percent among white men who have sex with men (MSM) between 2005 and 2014, but increased 24 percent among Latino MSM.<sup>21</sup> Black MSM HIV diagnoses have remained about the same since 2010 though initially increased 22 percent.<sup>22</sup>

The CDC compared national estimates of sepsis-related mortality based on death certificates with previously published sepsis mortality estimates generated using administrative claims data. Using death certificate data for the period 1999–2014, the CDC found that a total of 2,470,666 decedents (6 percent

of all deaths) had sepsis listed among the causes of death (sepsis-related deaths); for 22 percent of these decedents, sepsis was listed as the underlying cause of death.<sup>23</sup> The Centers for Medicare & Medicaid Services (CMS) recently implemented the severe sepsis and septic shock bundle (SEP-1) as part of its Inpatient Quality Reporting (IQR) program but results are not currently publicly reported.

## NQF Portfolio of Performance Measures for Infectious Disease Conditions

The Infectious Disease Standing Committee (see [Appendix D](#)) oversees NQF's portfolio of Infectious Disease measures that includes measures for HIV/AIDS, Sepsis and Septic Shock (see [Appendix B](#)). This portfolio contains nine measures: seven process measures, one outcome measure, and one composite measure (see table below).

**Table 1. NQF Infectious Disease Portfolio of Measures**

	Process	Outcome/Resource Use	Composite
<b>HIV/AIDS</b>	5	1	0
<b>Sepsis and Septic Shock</b>	0	0	1
<b>Respiratory</b>	2	0	0
<b>Total</b>	7	1	1

Additional measures related to Infectious Disease are assigned to other projects. These include various screening measures for sexually transmitted infections and Hepatitis C (Perinatal and Health and Well-being projects), vaccination measures (Perinatal and Health and Well-being projects), respiratory measures (Pulmonary project), facility acquired infection measures (Patient Safety project) and perioperative antibiotic measures (Surgery).

## National Quality Strategy

NQF-endorsed measures for Infectious Disease care support the [National Quality Strategy \(NQS\)](#). NQS serves as the overarching framework for guiding and aligning public and private efforts across all levels (local, State, and national) to improve the quality of healthcare in the U.S. The NQS establishes the "triple aim" of better care, affordable care, and healthy people/communities, focusing on six priorities to achieve those aims: *Safety, Person and Family Centered Care, Communication and Care Coordination, Effective Prevention and Treatment of Illness, Best Practices for Healthy Living, and Affordable Care*.

Quality measures for HIV/AIDS and sepsis care align with several of the NQS priorities, including:

- **Effective Prevention and Treatment.** Vaccinations against the flu or pneumonia help prevent infections that can lead to sepsis. In addition to prevention, early treatment of sepsis decreases morbidity and mortality.
- **Care Affordability.** The implementation of the Affordable Care Act (ACA) increased access to critical HIV testing, prevention, and care services nationwide.



## Use of Measures in the Portfolio

Endorsement of measures by NQF is valued not only because the evaluation process itself is both rigorous and transparent, but also because evaluations are conducted by multi-stakeholder committees comprised of clinicians and other experts from the full range of healthcare providers, employers, health plans, public agencies, community coalitions, and patients—many of whom use measures on a daily basis to ensure better care. Moreover, NQF-endorsed measures undergo routine "maintenance" (i.e., re-evaluation) to ensure that they are still the best-available measures and reflect the current science. Importantly, federal law requires that preference be given to NQF-endorsed® measures for use in federal public reporting and performance-based payment programs. NQF measures also are used by a variety of stakeholders in the private sector, including hospitals, health plans, and communities.

The measures in the infectious disease portfolio are in use in at least one federal program. The HIV/AIDS measures are included in CMS' Physician Quality Reporting System (PQRS) and HRSA's Ryan White & Global HIV/AIDS Programs. The sepsis measure is included in the CMS Hospital Inpatient Quality Reporting (IQR) program. See [Appendix C](#) for details of federal program use for the measures in the portfolio.

## Improving NQF's Infectious Disease Portfolio

During their discussions, the Committee identified additional areas where measure development is needed, including:

- Measures that underscore the value of infectious disease (ID) consultation, which studies have shown to improve outcomes. For example, the rate of ID consults in those with *Staphylococcus aureus* bacteremia, cryptococcal infection and HIV patients on ART.
- HPV screening in females with HIV.

The Committee reviewed the measurement gaps previously identified in 2012. Since that time, many of the previously identified gaps have been filled by new measures in various NQF projects. Specifically, 10 outcome measures related to infectious disease are now endorsed, measures that evaluate improvement in device associated infections in the hospital setting (two measures), measures that include follow up for screening tests (one measure), screening for STI's including HPV (three measures), and antimicrobial stewardship measures (one measure). Please see [Appendix B](#) for more details.

## Infectious Disease Measure Evaluation

On March 14, 2017, the Infectious Disease Standing Committee evaluated four new measures and five measures undergoing maintenance review against [NQF's standard evaluation criteria](#). To facilitate the evaluation, the committee and candidate standards were divided into two workgroups for preliminary review of the measures against the evaluation sub-criteria prior to consideration by the entire Standing Committee.

**Table 2. Infectious Disease Measure Evaluation Summary**

	Maintenance	New	Total
<b>Measures under consideration</b>	5	4	9
<b>Measures recommended for endorsement</b>	5	4	9
<b>Reasons for not recommending</b>	Importance – 0 Scientific Acceptability – 0 Overall – 0 Competing Measure – 0	Importance – 0 Scientific Acceptability – 0 Overall – 0 Competing Measure – 0	

## Comments Received Prior to Committee Evaluation

NQF solicits comments on endorsed measures on an ongoing basis through the [Quality Positioning System \(QPS\)](#). In addition, NQF solicits comments prior to the evaluation of the measures via an online tool located on the project webpage. For this evaluation cycle, the pre-evaluation comment period was open from February 9 – February 23, 2017. A total of six pre-evaluation comments were received ([Appendix G](#)).

All submitted comments were provided to the Committee prior to its initial deliberations during the workgroups calls.

## Overarching Issues

During the Standing Committee’s discussion of the measures, several overarching issues emerged that were factored into the Committee’s ratings and recommendations for multiple measures and are not repeated in detail with each individual measure.

### *Unintended Consequences*

The Committee expressed concerns with the potential unintended consequences associated with diagnosing and treating sepsis. The combination of symptoms associated with sepsis may be associated with other diagnoses such as pneumonia or myocardial infarction (MI). The Committee noted that patients may receive antibiotics prior to a conclusive diagnosis of sepsis – this is similar to the unintended consequences that occurred with the overuse of antibiotics in the emergency room for pneumonia. Patients may experience hypotension due to an acute MI rather than sepsis, yet, receive IV fluids in an effort to meet the data elements associated with the sepsis bundle.

### *Balancing Measures*

The Committee discussed the need for balancing measures to avoid unintended consequences as described above. The Committee suggested the following balancing measures for sepsis:

- Overuse of broad spectrum antibiotics
- Patients with heart failure and/or a MI who were overloaded with IV fluids
- Incidence of c. difficile secondary to overuse of antibiotics

NQF-endorsed measures related to antimicrobial use and *C. difficile* include #2720: National Healthcare Safety Network (NHSN) Antimicrobial Use Measure and #1717: National Healthcare Safety Network (NHSN) Facility-wide Inpatient Hospital-Onset Clostridium Difficile Infection (CDI). There are no NQF endorsed measures related to fluid overload in patients with heart failure and/or MI.

## Refining the NQF Measure Evaluation Process

In an effort to respond to evolving stakeholder needs, NQF constantly works to improve the consensus development process (CDP). In 2014, NQF transitioned to the use of standing committees for ongoing maintenance of endorsed measures and in 2015, NQF updated its Maintenance of NQF Endorsement policy to emphasize what has been learned about previously endorsed measures. Changes to the Maintenance of Endorsement policy is described below.

### Maintenance of NQF Endorsement

To streamline and improve the periodic evaluation of currently-endorsed measures, NQF has updated the way it re-evaluates measures for maintenance of endorsement. This change took effect beginning October 1, 2015. NQF's endorsement criteria have not changed, and all measures continue to be evaluated using the same criteria. However, under the new approach, there is a shift in emphasis for evaluation of currently-endorsed measures:

- **Evidence:** If the developer attests that the evidence for a measure has not changed since its previous endorsement evaluation, there is a decreased emphasis on evidence, meaning that the Committee may accept the prior evaluation of this criterion without further discussion or need for a vote. This applies only to measures that previously passed the evidence criterion without an exception. If a measure was granted an evidence exception, the evidence for that measure must be revisited.
- **Opportunity for Improvement (Gap):** For re-evaluation of endorsed measures, there is increased emphasis on current performance and opportunity for improvement. Endorsed measures that are “topped out” with little opportunity for further improvement are eligible for Inactive Endorsement with Reserve Status.
- **Reliability:**
  - Specifications: There is no change in the evaluation of the current specifications.
  - Testing: If the developer has not presented additional testing information, the Committee may accept the prior evaluation of the testing results without further discussion or need for a vote.
- **Validity:** There is less emphasis on this criterion if the developer has not presented additional testing information, and the Committee may accept the prior evaluation of this sub criterion without further discussion and vote. However, the Committee still considers whether the specifications are consistent with the evidence. In addition, for outcome measures, the Committee discusses questions required for the SDS Trial even if no change in testing is presented.
- **Feasibility:** The emphasis on this criterion is the same for both new and previously-endorsed measures, as feasibility issues might have arisen for endorsed measures that have been implemented.

- **Usability and Use:** For re-evaluation of endorsed measures, there is increased emphasis on the use of the measure, especially use for accountability purposes. There also is an increased emphasis on improvement in results over time and on unexpected findings, both positive and negative.

### *The New Endorsement and Appeals Process*

In August 2016, NQF implemented changes to its ratification and appeals process that were initiated and approved by its Board of Directors. Following public comment and voting by the NQF membership, the Consensus Standards Approval Committee (CSAC) will make the final measure endorsement decision, without ratification by another body. Additionally, the Board requested NQF to establish a five-member Appeals Board that will be responsible for adjudicating all submitted appeals regarding measure endorsement decisions. These changes apply to NQF measure endorsement projects with in-person meetings scheduled after August 2016.

The newly, constituted Appeals Board, composed of NQF Board members and former CSAC and/or committee members, will adjudicate appeals to measure endorsement decisions without a review by the CSAC. The decision of the Appeals Board will be final.

All submitted appeals will be published on the NQF website. Staff will compile the appeals for review by the Appeals Board, which will evaluate the concern(s) raised and determine if the appeal should warrant overturning the endorsement decision. Decisions on an appeal of endorsement will be publicly available on NQF's website.

Throughout the process, project staff will serve as liaisons between the CSAC, the Appeals Board, the committee, developers/stewards, and the appellant(s) to ensure the communication, cooperation, and appropriate coordination to complete the project efficiently.

## Summary of Measure Evaluation

The following brief summaries of the measure evaluation highlight the major issues that were considered by the Committee. Details of the Committee's discussion and ratings of the criteria for each measure are included in [Appendix A](#).

### *HIV/AIDS*

#### **2082 HIV Viral Load Suppression (Health Resources and Services Administration - HIV/AIDS Bureau): Recommended**

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year; **Measure Type:** Outcome; **Level of Analysis:** Facility; **Setting of Care:** Clinician Office/Clinic; **Data Source:** Laboratory, Paper Records

The main goal of antiretroviral therapy for people living with HIV (PLWH) is to inhibit HIV replication, which in turn reduces PLWH's risk of HIV-associated morbidity and mortality. This maintenance measure was last endorsed in 2012 and calculates the percentage of clients with a viral load of <200 copies/mL at the last viral load test. The Committee agreed that an opportunity for improvement in performance continues to exist. The developer provided updated reliability testing and explained that they did not update the validity testing because they intend to replace this paper-based measure with #3210, the electronically specified version of this measure. The Committee agreed the measure met the reliability and validity criteria.

#### **3210 HIV Viral Load Suppression – Legacy eMeasure (Health Resources and Services Administration - HIV/AIDS Bureau): Recommended**

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year; **Measure Type:** Outcome; **Level of Analysis:** Facility; **Setting of Care:** Clinician Office/Clinic; **Data Source:** Electronic Health Record (Only), Other

This "legacy" eMeasure is the electronically specified version of #2082, currently used in federal programs. The Committee discussed #2082 first, and because the information provided for evidence and opportunity for improvement is identical for the two measures, the Committee agreed to apply the voting results for these criteria to #3210. The developer tested the measure with the Bonnie tool using synthetic test cases. Testing results showed that the test cases passed the measure as expected which demonstrates the measure logic performs as constructed. The Committee agreed the measure met the minimum criteria for reliability and validity required for legacy eMeasures at this time.

#### **2079 HIV Medical Visit Frequency (Health Resources and Services Administration - HIV/AIDS Bureau): Recommended**

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between medical visits.; **Measure Type:** Process; **Level of Analysis:** Facility; **Setting of Care:** Clinician Office/Clinic; **Data Source:** Paper Records

Poor retention in care is associated with delayed or failed receipt of antiretroviral (ART) therapy, delayed time to viral suppression, and other behaviors or adverse clinical events such as low adherence to ART therapy and increased sexual risk transmission behaviors. This maintenance measure last endorsed in 2012 calculates the percentage of patients with HIV who had at least one medical visit in each 6-month period of a 24-month measurement period with a minimum of 60 days between medical visits. The Committee questioned the 6-month time interval and minimum 60-day interval between visits because the evidence provided does not include a specific time interval. The Committee did not agree that the evidence supported the measure focus; however, the Committee agreed retention in care was important and voted to pass the measure on evidence with an exception. The Committee agreed that an opportunity for improvement in performance continues to exist. The developer provided updated reliability testing and explained that they did not update the validity testing because they intend to replace this paper-based measure with #3209, the electronically specified version of this measure. The Committee agreed the measure met the reliability and validity criteria.

### **3209 HIV Medical Visit Frequency – *Legacy eMeasure* (Health Resources and Services Administration - HIV/AIDS Bureau): Recommended**

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in each 6-month period within 24 months with a minimum of 60 days between medical visits. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care; **Measure Type:** Process; **Level of Analysis:** Facility; **Setting of Care:** Clinician Office/Clinic; **Data Source:** Electronic Health Record (Only)

This “legacy” eMeasure is the electronically specified version of #2079, currently used in federal programs. The Committee discussed #2079 first, and because the information provided for evidence and opportunity for improvement is identical for the two measures, the Committee agreed to apply the voting results for these criteria to #2079. The developer tested the measure with the Bonnie tool using synthetic test cases. Testing results showed that the test cases passed the measure as expected which demonstrates the measure logic performs as constructed. The Committee agreed the measure met the minimum criteria for reliability and validity required for legacy eMeasures at this time.

### **2080 Gaps in HIV Medical Visits (Health Resources and Services Administration - HIV/AIDS Bureau): Recommended**

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV who did not have a medical visit in the last 6 months of the measurement year; **Measure Type:** Process; **Level of Analysis:** Clinician: Group/Practice, Facility; **Setting of Care:** Clinician Office/Clinic; **Data Source:** Other, Paper Records

Retention in care plays a critical role in assisting people living with HIV in their pursuit of achieving viral control and reducing new infections. This maintenance measure last endorsed in 2012 calculates the percentage of HIV patients who did not have a medical visit in the last 6 months of the measurement year (at least one visit every six months). The Committee noted that guidelines have changed and now recommend viral load assessment every six months but durably suppressed patients can be seen less

frequently. The Committee agreed that the evidence provided was insufficient; however, they agreed retention in care is important and voted to pass the measure on evidence with an exception. Notably, the Committee agreed that the measure could be used to help providers prioritize clients who have not been in care. The Committee agreed an opportunity for improvement continues to exist and noted disparities in care among age, gender, and racial/ethnic groups. The developer provided updated reliability testing and explained that they did not update the validity testing because they intend to replace this paper-based measure with the electronically specified version of this measure currently under development. The Committee agreed the measure met the reliability and validity criteria.

### **2083 Prescription of HIV Antiretroviral Therapy (Health Resources and Services Administration - HIV/AIDS Bureau): Recommended**

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV prescribed antiretroviral therapy for the treatment of HIV infection during the measurement year; **Measure Type:** Process; **Level of Analysis:** Facility; **Setting of Care:** Clinician Office/Clinic; **Data Source:** Other, Paper Records, Pharmacy

Antiretroviral therapy (ART) delays the progression of HIV and current treatment guidelines recommend ART for all people living with HIV (PLWH). This maintenance measure last endorsed in 2012 calculates the percentage of HIV patients with a prescription for ART. The Committee debated the evidence noting that the guideline recommends that PLWH be on ART although the focus of the measure is whether the provider prescribed ART. Ultimately, the Committee agreed the evidence provided was sufficient and accepted the previous Committee's vote on this criterion. The Committee agreed an opportunity for improvement continues to exist and noted disparities in care among age, gender, and racial/ethnic groups. The developer provided updated reliability testing and explained that they did not update the validity testing because they intend to replace this paper-based measure with #3211, the electronically specified version of this measure. The Committee agreed the measure met the reliability and validity criteria.

### **3211 Prescription of HIV Antiretroviral Therapy – Legacy eMeasure (Health Resources and Services Administration - HIV/AIDS Bureau): Recommended**

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV prescribed antiretroviral therapy for the treatment of HIV infection during the measurement year; **Measure Type:** Process; **Level of Analysis:** Facility; **Setting of Care:** Clinician Office/Clinic; **Data Source:** Electronic Health Record (Only)

This "legacy" eMeasure is the electronically specified version of #2083, currently used in federal programs. The Committee discussed #2083 first, and because the information provided for evidence and opportunity for improvement is identical for the two measures, the Committee agreed to apply the voting results for these criteria to #2083. The developer tested the measure with the Bonnie tool using synthetic test cases. Testing results showed that the test cases passed the measure as expected which demonstrates the measure logic performs as constructed. The Committee agreed the measure met the minimum criteria for reliability and validity required for legacy eMeasures at this time.



## *Sepsis and Septic Shock*

### **3215 Adult Inpatient Risk Adjusted Sepsis Mortality (New York State Department of Health): Recommended**

**Description:** Annual risk adjusted inpatient mortality rate for adult patients (aged 18 and over) admitted to acute care hospitals with diagnosis of severe sepsis or septic shock. The measure includes patients in acute care hospital settings over one year timeframe who had, either on admission, or during their hospital stay, a clinical diagnosis of severe sepsis (now referred to as 'sepsis') or septic shock using criteria described in the International Sepsis Definitions (Sepsis-2); **Measure Type:** Outcome; **Level of Analysis:** Facility; **Setting of Care:** Hospital: Acute Care Facility; **Data Source:** Claims (Other), EHRs Hybrid, Laboratory, Management Data, Non-Medical Data, Paper Records, Pharmacy, Registry

Hospitals that use early sepsis detection approaches coupled with rapid delivery of basic resuscitation interventions are able to influence mortality rates and improve patient outcomes. This newly-submitted facility-level measure estimates the probability of mortality for patients admitted to acute care hospitals with severe sepsis or septic shock. The Committee agreed that the performance rates that ranged from 1.0% to 95.0% in 2015 demonstrated a variation in hospital performance. The Committee did not express any concerns with the reliability and validity of the measure. The Committee discussed the amount of manual chart abstraction required to collect the necessary data for this measure but concluded it was feasible.

### **0500 Severe Sepsis and Septic Shock Management Bundle (Henry Ford Hospital): Recommended**

**Description:** This measure focuses on adults 18 years and older with a diagnosis of severe sepsis or septic shock. Consistent with Surviving Sepsis Campaign guidelines, the measure contains several elements, including measurement of lactate, obtaining blood cultures, administering broad-spectrum antibiotics, fluid resuscitation, vasopressor administration, reassessment of volume status and tissue perfusion, and repeat lactate measurement. As reflected in the data elements and their definitions, these elements should be performed in the early management of severe sepsis and septic shock; **Measure Type:** Composite; **Level of Analysis:** Facility; **Setting of Care:** Hospital; **Data Source:** Imaging-Diagnostic, Laboratory, Other, Paper Records, Pharmacy

Patients diagnosed with severe sepsis or septic shock who receive all the elements of care, including measurement of lactate levels (initial and repeat), blood cultures, broad spectrum antibiotics, fluid resuscitation, vasopressor administration and volume status and tissue perfusion reassessment have shown a decrease in mortality. This facility-level, all-or-none, composite measure, originally endorsed in 2012, and most recently in 2014, assesses the percentage of patients that received all of the applicable elements of care for severe sepsis or septic shock. The Committee agreed that despite the performance increase from 34.4 percent in Q4 2015 to 44.0 percent in Q2 2016, a performance gap exists. The Standing Committee thoroughly discussed the reliability and validity of the measure and concluded the testing and specifications met both criteria. Due to several concerns voiced related to unintended consequences, the Committee did not reach consensus on the Usability and Use criterion. However, the Committee supported the measure overall and recommended it for endorsement.



## References

- 1 Centers for Disease Control and Prevention (CDC). National Center for Health Statistics faststats infectious disease website. <http://www.cdc.gov/nchs/faststats/infectious-disease.htm>. Last accessed April 2017.
- 2 Research!America. *Facts about infectious diseases*. Alexandria, VA: Research!America; 2015. Investment in Research Saves Lives and Money #15. Available at <http://www.researchamerica.org/sites/default/files/uploads/factsheet15infectiousdiseases.pdf>. Last accessed April 2017.
- 3 Tono CM, Andrews RM. *National Inpatient Hospital Costs: the Most Expensive Conditions by Payer, 2011*. Rockville, MD: Agency for Healthcare Research and Quality (AHRQ). Healthcare Cost and Utilization Project (HCUP) Statistical Brief #160. Available at <https://www.hcup-us.ahrq.gov/reports/statbriefs/sb160.pdf>. Last accessed April 2017.
- 4 CDC. About HIV/AIDS what is HIV? website. <https://www.cdc.gov/hiv/basics/whatishiv.html>. Last accessed April 2017.
- 5 AIDSinfo.gov. Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents website. <https://aidsinfo.nih.gov/guidelines/html/1/adult-and-adolescent-arv-guidelines/30/adherence-to-art>. Last accessed April 2017.
- 6 CDC. Sepsis website. <https://www.cdc.gov/sepsis/index.html>. Last accessed April 2017.
- 7 CDC. Sepsis Fact Sheet. Atlanta, GA: CDC;2016. Available at <https://www.cdc.gov/sepsis/pdfs/sepsis-fact-sheet.pdf>. Last accessed April 2017.
- 8 Johnson S, Gotter A. Septic shock. *Healthline Media*. July 11, 2016. Available at <http://www.healthline.com/health/septic-shock>. Last accessed April 2017.
- 9 National Institute of General Medical Sciences. Sepsis fact sheet website. [https://www.nigms.nih.gov/education/pages/factsheet\\_sepsis.aspx](https://www.nigms.nih.gov/education/pages/factsheet_sepsis.aspx). Last accessed April 2017.
- 10 National Institute of General Medical Sciences. Sepsis fact sheet website. [https://www.nigms.nih.gov/education/pages/factsheet\\_sepsis.aspx](https://www.nigms.nih.gov/education/pages/factsheet_sepsis.aspx). Last accessed April 2017.
- 11 National Institute of General Medical Sciences. Sepsis fact sheet website. [https://www.nigms.nih.gov/education/pages/factsheet\\_sepsis.aspx](https://www.nigms.nih.gov/education/pages/factsheet_sepsis.aspx). Last accessed April 2017.
- 12 National Institute of General Medical Sciences. Sepsis fact sheet website [https://www.nigms.nih.gov/education/pages/factsheet\\_sepsis.aspx](https://www.nigms.nih.gov/education/pages/factsheet_sepsis.aspx). Last accessed April 2017.
- 13 National Institute of General Medical Sciences. Sepsis fact sheet website. [https://www.nigms.nih.gov/education/pages/factsheet\\_sepsis.aspx](https://www.nigms.nih.gov/education/pages/factsheet_sepsis.aspx). Last accessed April 2017.
- 14 CDC. Division of HIV/AIDS prevention. *Annual Report 2015. Putting Prevention Advances to Work*. <https://www.cdc.gov/hiv/pdf/policies/cdc-hiv-2015-dhap-annual-report.pdf>. Last accessed April 2017.
- 15 CDC. Division of HIV/AIDS prevention. *Annual Report 2015. Putting Prevention Advances to Work*. <https://www.cdc.gov/hiv/pdf/policies/cdc-hiv-2015-dhap-annual-report.pdf>. Last accessed April 2017.
- 16 CDC. Division of HIV/AIDS prevention. *Annual Report 2015. Putting Prevention Advances to Work*. <https://www.cdc.gov/hiv/pdf/policies/cdc-hiv-2015-dhap-annual-report.pdf>. Last accessed April 2017.
- 17 CDC. Division of HIV/AIDS prevention. *Annual Report 2015. Putting Prevention Advances to Work*. <https://www.cdc.gov/hiv/pdf/policies/cdc-hiv-2015-dhap-annual-report.pdf>. Last accessed April 2017.
- 18 CDC. Division of HIV/AIDS prevention. *Annual Report 2015. Putting Prevention Advances to Work*. <https://www.cdc.gov/hiv/pdf/policies/cdc-hiv-2015-dhap-annual-report.pdf>. Last accessed April 2017.
- 19 CDC. Division of HIV/AIDS prevention. *Annual Report 2015. Putting Prevention Advances to Work*. <https://www.cdc.gov/hiv/pdf/policies/cdc-hiv-2015-dhap-annual-report.pdf>. Last accessed April 2017.
- 20 CDC. Division of HIV/AIDS prevention. *Annual Report 2015. Putting Prevention Advances to Work*. <https://www.cdc.gov/hiv/pdf/policies/cdc-hiv-2015-dhap-annual-report.pdf>. Last accessed April 2017.
- 21 CDC. Division of HIV/AIDS prevention. *Annual Report 2015. Putting Prevention Advances to Work*. <https://www.cdc.gov/hiv/pdf/policies/cdc-hiv-2015-dhap-annual-report.pdf>. Last accessed April 2017.
- 22 CDC. Division of HIV/AIDS prevention. *Annual Report 2015. Putting Prevention Advances to Work*. <https://www.cdc.gov/hiv/pdf/policies/cdc-hiv-2015-dhap-annual-report.pdf>. Last accessed April 2017.

<sup>23</sup> Epstein L, Dantes R, Magill S, Fiore A. Varying Estimates of Sepsis Mortality Using Death Certificates and Administrative Codes — United States, 1999–2014. *MMWR Morb Mortal Wkly Rep*. 2016; 65:342–345. DOI: <http://dx.doi.org/10.15585/mmwr.mm6513a2>. Last accessed April 2017.

## Appendix A: Details of Measure Evaluation

**Rating Scale:** H=High; M=Moderate; L=Low; I=Insufficient; NA=Not Applicable; Y=Yes; N=No

### Measures Recommended

2082 HIV Viral Load Suppression
<a href="#">Submission</a>   <a href="#">Specifications</a>
<p><b>Description:</b> Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.</p> <p><b>Numerator Statement:</b> Number of patients in the denominator with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year</p> <p><b>Denominator Statement:</b> Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the measurement year</p> <p><b>Exclusions:</b> There are no patient exclusions.</p> <p><b>Adjustment/Stratification:</b> No risk adjustment or risk stratification</p> <p><b>Level of Analysis:</b> Facility</p> <p><b>Setting of Care:</b> Clinician Office/Clinic</p> <p><b>Type of Measure:</b> Outcome</p> <p><b>Data Source:</b> Laboratory, Paper Records</p> <p><b>Measure Steward:</b> Health Resources and Services Administration - HIV/AIDS Bureau</p>
<p><b>STANDING COMMITTEE MEETING [03/14/2017]</b></p> <p><b>1. Importance to Measure and Report:</b> <u>The measure meets the Importance criteria</u> (1a. Evidence, 1b. Performance Gap) 1a. Evidence: <b>Y-16; N-0</b>; 1b. Performance Gap: <b>H-10; M-3; L-0; I-0</b>;</p> <p><u>Rationale:</u></p> <ul style="list-style-type: none"> <li>For the 2013 endorsement evaluation, the developer noted that viral suppression is a main goal of HIV treatment and an indicator of treatment success and reduction of potential HIV transmission. The developer also provided multiple guidelines for the administration of antiretroviral therapy and viral load monitoring intervals for adults, adolescents and pregnant women.</li> <li>For the current maintenance of endorsement evaluation, the Committee agreed with the developer that antiretroviral therapy and viral suppression reduce morbidity and mortality associated with HIV. The developer also submitted updated guidelines for the administration of antiretroviral therapy and viral load monitoring intervals for people living with HIV (PLWH).</li> <li>A Committee member pointed out CDC data that showed viral suppression rates were overestimated by 20% when looking at the last viral load test in a measurement period. Specifically, the analysis showed that not all patients that were suppressed at the end of the year were suppressed <i>throughout</i> the year. The developer clarified that this measure is not intended to be a durable suppression measure.</li> <li>The Committee reviewed 2010-2014 performance gap data from the Ryan White HIV/AIDS Program Services Report, with 65% viral load suppression among the 10<sup>th</sup> percentile of providers, and 94% among the 90<sup>th</sup> percentile.</li> <li>Overall, the Committee agreed that the measure met this criterion.</li> </ul> <p><b>2. Scientific Acceptability of Measure Properties:</b> <u>The measure meets the Scientific Acceptability criteria</u> (2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity) 2a. Reliability: <b>H-12; M-4; L-0; I-0</b> 2b. Validity: <b>Accepted prior evaluation</b></p>

## 2082 HIV Viral Load Suppression

### Rationale:

- For the 2013 endorsement evaluation, the developer conducted a signal-to-noise analysis using nine of 18 HIV Research Network (HIVRN) sites and 11,134 patients in 2010. The clinic-specific reliability ranged from 0.93 to 0.99 with a between clinic variance of 0.0066.
- For the current evaluation, the developer provided updated reliability testing using the beta binomial method to assess the signal-to-noise-ratio in over 800 clinics participating in the RSR database; median reliability ranged from 0.95 to 0.98 indicating high reliability.
- A Committee member noted that since reliability ranged from 0.29 to 0.98 that some clinics may not be performing well on the measure. The developer stated that lower signal-to-noise ratios generally came from clinics with small patient populations (i.e., <25 patients).
- A Committee member questioned why the measure is specified at less than 200 copies/mL. The developer cited HHS guidelines, which indicate that although tests can quantify a viral load down to individual copies of the virus, it is generally accepted that 200 is the threshold for viral load suppression.
- Another Committee member asked the developer to define “comprehensive HIV care.” The developer noted that HIV care is in various models, but ultimately they define comprehensive care as being provided by someone who is addressing the patient’s HIV care (i.e., assessing whether the client is on antiretroviral medications and that viral load testing has been performed).
- One Committee member asked how the measure accounts for providers who could game the measure by choosing not to see or to create an unfavorable environment for non-adherent patients. The developer noted that it would be difficult to capture that information at the individual provider level since providers do not have a lot of control over the patients they see. The developer also noted that there is a measure (#2079) that assesses whether patients come back to the same clinic.
- In discussing what constitutes a medical visit, the Committee questioned whether non-face-to-face visits are included in the measure. The developer clarified that face-to-face, video visits, or another type of visit that is documented by the provider as a medical visit. If a patient comes in for lab work, to meet with an ancillary staff member or to pick up paperwork, the visit is not counted as a medical visit. The measure does not address whether visits are billable. The developer also clarified that patients are counted in the denominator if they came in for at least one visit.
- The Committee noted that the developer used a technical group and Ryan White grant recipients to test the face validity of the measure. A Committee member questioned whether the same group of people established the measure and assessed face validity. The developer clarified that these were the same group of people.
- A Committee member expressed concern that although the measure is not risk adjusted, since people living with HIV (PLWH) are a marginalized population, that the measure could be used in MIPS, a pay for performance program. The Committee member expressed concern that a provider could be penalized based on poor performance on this measure if they provided medical care to populations who are non-adherent to medical care (e.g., people experiencing homelessness, active substance users).
- The developer stated that existing methods for risk adjustment do not apply to this measure because it is currently not used in a payment program. However, the eMeasure version of this measure was submitted for consideration in the Centers for Medicare & Medicaid Services (CMS) Merit-based Incentive Payment System (CMS MIPS); therefore, the developer will work with CMS to develop appropriate methods for risk adjustment of the eMeasure.
- The Committee emphasized that there are concerns among PLWH that as measurement of HIV care is increasingly implemented, that providers will make an unfavorable environment for PLWH.
- The Committee questioned how the measure accounts for patients who have access to the care but choose not to receive it. The developer clarified that the intent of the measure is not to achieve 100%, nor is it a measure of the patient’s compliance or ability to participate in care. The measure is intended to assess the clinic’s ability to support a client in reaching viral load suppression.
- Upon a vote, the Committee agreed the measure met the reliability criteria.

<b>2082 HIV Viral Load Suppression</b>
<ul style="list-style-type: none"> <li>For the 2013 endorsement evaluation, face validity was established through a technical work group established for the development of the measures. This measure was found to be important, usable, and feasible by the technical work group overseeing the development of this measure and several others.</li> <li>For the current evaluation, the developer did not provide updated validity testing because they intend to replace this paper-based measure with #3210, the electronically specified version of this measure. The Committee agreed that the previous validity testing results were sufficient and accepted the prior evaluation without further discussion and vote.</li> </ul>
<p><b>3. Feasibility: H-14; M-1; L-0; I-0</b>  <i>(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/ unintended consequences identified 3d. Data collection strategy can be implemented)</i>  <u>Rationale:</u></p> <ul style="list-style-type: none"> <li>All data elements for this measure are generated during routine provision of care and are in defined fields in the electronic health record. Without further discussion, the Committee agreed the measure met this criterion.</li> </ul>
<p><b>4. Usability and Use: H-15; M-0; L-0; I-0</b>  <i>(Used and useful to the intended audiences for 4a. Accountability and Transparency; 4b. Improvement; and 4c. Benefits outweigh evidence of unintended consequences)</i>  <u>Rationale:</u></p> <ul style="list-style-type: none"> <li>The measure is publicly reported and used by accountability programs including MIPS and PQRS. The measure has shown improvement in viral load suppression and no potential harms were identified in measure implementation. Without further discussion, the Committee agreed the measure met this criterion.</li> </ul>
<p><b>5. Related and Competing Measures</b></p> <ul style="list-style-type: none"> <li>This measure is related to the following: <ul style="list-style-type: none"> <li>0407 HIV/AIDS: HIV RNA Control After Six Months of Potent Antiretroviral Therapy</li> <li>0405 HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis</li> <li>0409 HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</li> <li>2079 HIV Medical Visit Frequency</li> <li>2080 Gap in HIV Medical Visits</li> <li>2083 Prescription of HIV Antiretroviral Therapy</li> <li>3211 Prescription of HIV Antiretroviral Therapy</li> <li>3210 HIV Viral Suppression</li> <li>3209 HIV Medical Visit Frequency</li> </ul> </li> </ul> <p>Per the developer, this measure is harmonized with all measures except for #0405 and #0409; there are plans to harmonize with #0405 and #0409.</p>
Standing Committee Recommendation for Endorsement: <b>Y-16; N-0</b>
<p><b>6. Public and Member Comment</b></p> <ul style="list-style-type: none"> <li><a href="#">One comment was submitted in support of the Committee's recommendation for endorsement.</a></li> </ul>
<b>7. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X</b>
<b>8. Board of Directors Vote: Y-X; N-X</b>
<b>9. Appeals</b>

## 3210 HIV Viral Load Suppression

[Submission](#) | [Specifications](#)

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year.

**Numerator Statement:** Patients with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year. The outcome being measured is HIV viral suppression.

**Denominator Statement:** Patients, regardless of age, diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year who had at least one medical visit in the measurement year. The target population for this measure is all people living with HIV.

**Exclusions:** There are no patient exclusions.

**Adjustment/Stratification:** No risk adjustment or risk stratification

**Level of Analysis:** Facility

**Setting of Care:** Clinician Office/Clinic

**Type of Measure:** Outcome,

**Data Source:** Electronic Health Record (Only), Other

**Measure Steward:** Health Resources and Services Administration - HIV/AIDS Bureau

### STANDING COMMITTEE MEETING [03/14/2017]

#### 1. Importance to Measure and Report: The measure meets the Importance criteria

(1a. Evidence, 1b. Performance Gap)

1a. Evidence: **Applied the vote from #2082**; 1b. Performance Gap: **Applied the vote from #2082**

##### Rationale:

- The Committee acknowledged that this measure shares the same evidence as #2082. Because it is a legacy eMeasure, there are no performance data. The Committee agreed to apply the vote from #2082 to this criterion.

#### 2. Scientific Acceptability of Measure Properties: The measure meets the Scientific Acceptability criteria

(2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity)

2a. Reliability: H-0; M-15; L-1; I-0 2b. Validity: H-0; M-16; L-0; I-0

##### Rationale:

- This measure was tested using 34 synthetic cases with the Bonnie tool. Results showed 100% coverage and all 34 cases passed the measure as expected, demonstrating that the measure logic works as constructed. All test cases with missing data performed according to the HQMF standard as expected. Testing also included specific ways to search for patients that might fall at the edge of the measure specifications.
- Without further discussion, the Committee agreed the measure met these criteria.

#### 3. Feasibility: H-12; M-4; L-0; I-0

(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/unintended consequences identified 3d. Data collection strategy can be implemented)

##### Rationale:

- Feasibility testing showed that the measure is feasible but the Committee questioned why the variable "Encounter Performed: Face-to-Face Interaction" and "Patient Characteristic Payer" scored a 2 out of 3 on the feasibility scorecard. The developer clarified that "Face-to-Face Interaction" scored lower because the value set is defined in SNOMED whereas the other encounter value sets are defined in CPT. The developer further clarified that the variable "Patient Characteristic Payer" is a supplemental data element required to be submitted for measures in federal programs but is not used in the measure logic.
- The Committee asked the developer to explain why they expect feasibility to improve from 98.89% to 99.44%. The developer clarified that the addition of the SNOMED codes will increase the feasibility.
- Overall, the Committee agreed the measure met this criterion.

## 3210 HIV Viral Load Suppression

### 4. Usability and Use: H-8; M-8; L-0; I-0

*(Used and useful to the intended audiences for 4a. Accountability and Transparency; 4b. Improvement; and 4c. Benefits outweigh evidence of unintended consequences)*

#### Rationale:

- The Committee noted that the measure was reviewed by NQF's Measure Application Partnership (MAP) for consideration in CMS' MIPS program. MAP recommended support of this measure for rulemaking with the condition that the Infectious Disease Standing Committee review the performance data to ensure a gap in care continues to exist.
- The developer clarified that this measure is intended to be used in an accountability program. Without further discussion, the Committee agreed the measure met this criterion.

### 5. Related and Competing Measures

- This measure is related to the following:
  - 0407 HIV/AIDS: HIV RNA Control After Six Months of Potent Antiretroviral Therapy
  - 0405 HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis
  - 0409 HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis
  - 2079 HIV Medical Visit Frequency
  - 2080 Gap in HIV Medical Visits
  - 2083 Prescription of HIV Antiretroviral Therapy
  - 3211 Prescription of HIV Antiretroviral Therapy
  - 3210 HIV Viral Suppression
  - 3209 HIV Medical Visit Frequency
- Per the developer, this measure is harmonized with all measures except for #0405 and #0409; there are plans to harmonize with #0405 and #0409.

Standing Committee Recommendation for Endorsement: **Y-16; N-0**

### 6. Public and Member Comment

- One comment was submitted in support of the Committee's recommendation for endorsement.

**7. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X**

**8. Board of Directors Vote: Y-X; N-X**

**9. Appeals**



## 2079 HIV Medical Visit Frequency

[Submission](#) | [Specifications](#)

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between medical visits. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.

**Numerator Statement:** Number of patients in the denominator who had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period. (Measurement period is a consecutive 24-month period of time.)

**Denominator Statement:** Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the first 6 months of the 24-month measurement period.

**Exclusions:** Patients who died at any time during the 24-month measurement period.

**Adjustment/Stratification:** No risk adjustment or risk stratification

**Level of Analysis:** Facility

**Setting of Care:** Clinician Office/Clinic

**Type of Measure:** Process

**Data Source:** Paper Records

**Measure Steward:** Health Resources and Services Administration - HIV/AIDS Bureau

### STANDING COMMITTEE MEETING [03/14/2017]

#### 1. Importance to Measure and Report: The measure meets the Importance criteria

(1a. Evidence, 1b. Performance Gap)

1a. Evidence: **H-0; M-2; L-1; I-13**; Evidence with Exception: **Y-14; N-2**; 1b. Performance Gap: **H-13; M-3; L-0; I-0**

#### Rationale:

- This measure calculates the percent of clients retained in care over a 24-month time period. Patients are counted in the numerator if they have had at least one medical visit at least 60 days apart in each 6 months of the measurement period.
- The Committee acknowledged the developer submitted updated evidence for the systematic monitoring of retention in care they could include visit adherence, gaps in care, and number of visits during a specified visit. A Committee member questioned whether there was evidence to support the 6-month interval and the 60-days between medical visits, especially when the guidelines state that clinically stable clients can be seen less frequently.
- The developer stated that the 6-month time period was selected as a 'middle ground' based on DHHS guidelines that outline the frequency of labs performed and medical visits. The 60-day period between medical visits was selected so as to not count patients, who for instance, have medical visits on consecutive days.
- Another Committee member expressed concern for programs receiving Ryan White program funds who must report this measure. The committee member noted that some programs may be forced to see stable clients more frequently just to meet the measure's requirements.
- The Committee then debated that the measure should focus on frequency of viral load testing for stable clients who only need to see a provider for lab work. They also questioned how the measure accounts for not penalizing providers who have durably suppressed clients. Some Committee members supported that the measure focus was acceptable since retention in care has been associated with improved clinical outcomes. A Committee member commented that medical visit frequency is more controlled by the health system than by the individual provider.
- The developer further clarified that they do not expect to reach 100% adherence on this measure and the measure is applicable broadly across all types of providers caring for any patient living with HIV.
- The Committee was conflicted as to whether the measure should be used for public health rather than quality improvement purposes.



## 2079 HIV Medical Visit Frequency

- Upon a vote, the Committee did not pass the measure on the Evidence criterion but agreed to vote on whether the measure warranted an exception to the evidence. The Committee discussed the importance of the measure on outcomes for people living with HIV but did not have evidence that explicitly stated that the measure as specified would lead to those outcomes. After a full discussion, the Committee opted to invoke the exception to the evidence criterion and agreed that medical visit frequency needed to be assessed since patients cannot be treated if they have not been seen by a provider. Ultimately, a majority of the Committee agreed that providers should be held accountable for this measure in the absence of empiric evidence and passed the measure on this criterion.
- The Committee noted that medical visit frequency had increased from 67% to 73%, with disparities – similar to those seen in #2082 – among race/ethnic, gender, and age groups.

### **2. Scientific Acceptability of Measure Properties: The measure meets the Scientific Acceptability criteria**

(2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity)

2a. Reliability: **H-13; M-2; L-0; I-0** 2b. Validity: **Accepted previous Evaluation**

#### Rationale:

- The Committee reviewed testing data from the RSR (covering 11 sites and more than 17,000 patients) that showed a median reliability of 0.97; upon a vote, the Committee agreed the measure met this criterion.
- Committee members questioned whether the measure is stratified to determine the model of care the patient receives (i.e., care from a primary care provider or an infectious disease provider). The developer clarified that the measure is not stratified by the type of provider.
- As was discussed for #2082, the developer clarified that tele-visits or other advanced methods count as a medical visit; the measure does not specify how the visit is delivered or whether the visit is billable.
- In response to the Committee's question as to why validity testing was not updated, the developer responded they used their resources for the testing and development of the eMeasures. The Committee then noted the performance data demonstrated the measure was able to identify differences in performance among providers. The Committee chose to accept the previous evaluation on this criterion.

### **3. Feasibility: H-13; M-1; L-0; I-0**

(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/unintended consequences identified 3d. Data collection strategy can be implemented)

#### Rationale:

- The Committee agreed that the measure was feasible with all data elements collected and generated as part of routine delivery of care. Without further discussion, the Committee agreed the measure met this criterion.

### **4. Usability and Use: H-11; M-4; L-0; I-0**

(Used and useful to the intended audiences for 4a. Accountability and Transparency; 4b. Improvement; and 4c. Benefits outweigh evidence of unintended consequences)

#### Rationale:

- The measure is currently in use in CMS' Physician Quality Report System (PQRS), Value Based Payment Modifier (VBPM), and MIPS programs. Without further discussion, the Committee agreed the measure met this criterion.

### **5. Related and Competing Measures**

- This measure is related to the following:
  - 2080 Gap in HIV Medical Visits
  - 2082 HIV Viral Suppression
  - 2083 Prescription of HIV Antiretroviral Therapy
  - 3211 Prescription of HIV Antiretroviral Therapy
  - 3210 HIV Viral Suppression

2079 HIV Medical Visit Frequency	
	<ul style="list-style-type: none"> <li>○ 3209 HIV Medical Visit Frequency</li> <li>○ 0405 HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis</li> <li>○ 0409 HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</li> <li>• Per the developer, the measure is harmonized with the first six measures listed above. For these six measures the target population is the same (i.e., people living with HIV), however the measure focus is different (gaps in visit, prescription of ARV therapy, and viral load suppression).</li> <li>• NQF #0405 and #0409 have been deferred for maintenance of endorsement. There are no additional steps the developer must take since the measure focus is different (HIV patients receiving PCP prophylaxis and those screened for STDs).</li> </ul>
Standing Committee Recommendation for Endorsement: <b>Y-15; N-0</b>	
<b>6. Public and Member Comment</b>	
	<ul style="list-style-type: none"> <li>• <a href="#">One comment was submitted in support of the Committee's recommendation for endorsement.</a></li> </ul>
<b>7. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X</b>	
<b>8. Board of Directors Vote: Y-X; N-X</b>	
<b>9. Appeals</b>	

## 3209 HIV Medical Visit Frequency

[Submission](#) | [Specifications](#)

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in each 6-month period within 24 months with a minimum of 60 days between medical visits. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.

**Numerator Statement:** Patients who had at least one medical visit in each 6-month of a consecutive 24 month period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period.

**Denominator Statement:** Patients, regardless of age, diagnosed with HIV during the first 3 months of the year preceding the measurement period or prior to the measurement period with at least one medical visit in the first 6 months of the year preceding the measurement period. The target population for this measure is all people living with HIV.

**Exclusions:** Patients who died at any time during the measurement period or the 12 months preceding the measurement period.

**Adjustment/Stratification:** No risk adjustment or risk stratification

**Level of Analysis:** Facility

**Setting of Care:** Clinician Office/Clinic

**Type of Measure:** Process

**Data Source:** Electronic Health Record (Only)

**Measure Steward:** Health Resources and Services Administration - HIV/AIDS Bureau

### STANDING COMMITTEE MEETING [03/14/2017]

#### 1. Importance to Measure and Report: The measure meets the Importance criteria

(1a. Evidence, 1b. Performance Gap)

1a. Evidence: **Applied the vote from #2079**; 1b. Performance Gap: **Applied the vote from #2079**

##### Rationale:

- The Committee acknowledged that this measure shares the same evidence as #2079. Because it is a legacy eMeasure, there is no performance data. The Committee agreed to apply the vote from #2079 to this criterion.

#### 2. Scientific Acceptability of Measure Properties: The measure meets the Scientific Acceptability criteria

(2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity)

2a. Reliability: **H-0; M-10; L-5; I-0** 2b. Validity: **H-0; M-14; L-1; I-0**

##### Rationale:

- This measure was tested using 64 synthetic cases with the Bonnie tool. Results showed 100% coverage and all 64 cases passed the measure as expected, demonstrating that the measure logic works as constructed. Testing also included specific ways to search for patients that might fall at the edge of the measure specifications.
- The Committee noted that the measure excludes patients who die during the measurement period. Another Committee member stated that the measure should exclude patients who are incarcerated during the measurement period since providers could be penalized on this measure if they provide care to specific populations that experience incarceration. The developer clarified that there is no standardized variable for incarceration in electronic health records.
- Without further discussion, the Committee agreed this measure met this criterion.

#### 3. Feasibility: **H-9; M-6; L-0; I-0**

(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/unintended consequences identified 3d. Data collection strategy can be implemented)

##### Rationale:

- Feasibility testing showed that the measure is feasible at 98.21% and will increase to 98.81%.

<b>3209 HIV Medical Visit Frequency</b>
<ul style="list-style-type: none"> <li>The Committee questioned why the variable “Encounter Performed: Face-to-Face Interaction” and “Patient Characteristic Payer” scored a 2 out of 3 on the feasibility scorecard. The developer clarified in the discussion of #3210 that “Face-to-Face Interaction” scored lower because the value set is defined in SNOMED whereas the other encounter value sets are defined in CPT. The “Patient Characteristic Payer” variable is a supplemental data element required to be submitted for measures in federal programs but is not used in the measure logic.</li> <li>Overall, the Committee agreed the measure met this criterion.</li> </ul>
<p><b>4. Usability and Use: H-7; M-8; L-0; I-1</b></p> <p><i>(Used and useful to the intended audiences for 4a. Accountability and Transparency; 4b. Improvement; and 4c. Benefits outweigh evidence of unintended consequences)</i></p> <p><u>Rationale:</u></p> <ul style="list-style-type: none"> <li>The Committee noted that the measure is not publically reported. The measure is planned to be used in MIPS. Without further discussion, the Committee agreed the measure met this criterion.</li> </ul>
<p><b>5. Related and Competing Measures</b></p> <ul style="list-style-type: none"> <li>This measure is related to the following: <ul style="list-style-type: none"> <li>2080 Gap in HIV Medical Visits</li> <li>2082 HIV Viral Suppression</li> <li>2083 Prescription of HIV Antiretroviral Therapy</li> <li>3211 Prescription of HIV Antiretroviral Therapy</li> <li>3210 HIV Viral Suppression</li> <li>3209 HIV Medical Visit Frequency</li> <li>0405 HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis</li> <li>0409 HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</li> </ul> </li> <li>Per the developer, the measure is harmonized with the first six measures listed above. For these six measures the target population is the same (i.e., people living with HIV), however the measure focus is different (gaps in visit, prescription of ARV therapy, and viral load suppression).</li> <li>NQF #0405 and #0409 have been deferred for maintenance of endorsement. There are no additional steps the developer must take since the measure focus is different (HIV patients receiving PCP prophylaxis and those screened for STDs).</li> </ul>
Standing Committee Recommendation for Endorsement: <b>Y-16; N-0</b>
<p><b>6. Public and Member Comment</b></p> <ul style="list-style-type: none"> <li><u>One comment was submitted in support of the Committee’s recommendation for endorsement.</u></li> </ul>
<b>7. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X</b>
<b>8. Board of Directors Vote: Y-X; N-X</b>
<b>9. Appeals</b>

## 2080 Gap in HIV Medical Visits

[Submission](#) | [Specifications](#)

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV who did not have a medical visit in the last 6 months of the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.

**Numerator Statement:** Number of patients in the denominator who did not have a medical visit in the last 6 months of the measurement year (Measurement year is a consecutive 12-month period of time).

**Denominator Statement:** Number of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in the first 6 months of the measurement year. (The measurement year can be any consecutive 12-month period.)

**Exclusions:** Patients who died at any time during the measurement year.

**Adjustment/Stratification:** No risk adjustment or risk stratification

**Level of Analysis: Clinician :** Group/Practice, Facility

**Setting of Care:** Clinician Office/Clinic

**Type of Measure:** Process

**Data Source:** Other, Paper Records

**Measure Steward:** Health Resources and Services Administration-HIV/AIDS Bureau

### STANDING COMMITTEE MEETING [03/14/2017]

#### 1. Importance to Measure and Report: The measure meets the Importance criteria

(1a. Evidence, 1b. Performance Gap)

1a. Evidence: **H-0; M-2; L-0; I-11** 1b. Performance Gap: **H-3; M-10; L-0; I-0**; Evidence Exception: **Y-12; N-1**

#### Rationale:

- The Committee questioned the value of the measure over the related measure, #2079. The developer clarified that this measure looks at a shorter time period and that some users pair this measure with the longer term retention measure (#2079).
- A Committee member acknowledged the importance of retention in care, but did not believe there was evidence related to two medical visits and health outcomes. The Committee asked the developer to clarify the definition of a medical visit. The developer noted they do not specify the mode of the visit (face-to-face vs. tele-health) but that tele-health visits are not ruled out.
- In response to the Committee's question as to how a gap in medical care is defined, the developer stated that if a patient has a medical visit in the first six months of the measurement period but not within the second six months then the client would have experienced a gap in care. Based on this definition, their data indicates that 21% of patients had a gap in medical visits. Better performance for this measure is indicated by a lower rate e.
- A Committee member noted that the guidelines have changed to say that viral load should be assessed every six months but that durably suppressed clients can be seen less frequently.
- The Committee agreed that the evidence provided was insufficient; however, they agreed retention in care is important and voted to pass the measure on evidence with an exception. Notably, the Committee agreed that the measure could be used to help providers prioritize clients who have not been in care.
- The Committee noted that the performance gap had increased from 2010 to 2014 (i.e., more people are not getting regular care). Without further discussion, the Committee agreed the measure met this criterion.

#### 2. Scientific Acceptability of Measure Properties: The measure meets the Scientific Acceptability criteria

(2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity)

2a. Reliability: **Accepted prior evaluation** 2b. Validity: **Accepted prior evaluation**

#### Rationale:

## 2080 Gap in HIV Medical Visits

- The Committee reviewed the testing data from over 800 participants in the Ryan White HIV/AIDS Program (RWHAP) that showed a median reliability of 0.973; without further discussion agreed to accept the prior evaluation of this measure.
- In discussion of measure validity, the Committee noted that new testing was provided but that face validity was completed. They also noted the exclusion for patients who died (<1%) had minimal impact on the overall score.
- The developer clarified they are field-testing an electronic version of this measure. Without further discussion, the Committee agreed to accept the prior evaluation of this measure.

### 3. Feasibility: H-10; M-3; L-0; I-0

*(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/unintended consequences identified 3d. Data collection strategy can be implemented)*

#### Rationale:

- The Committee acknowledged that data are generated or collected by and used by health care professionals during the provision of care. All data elements are in defined fields in electronic health records. There are no fees, licenses, or other requirements to use this measure.
- A Committee member commented that the measure is calculated in the inverse and it could be confusing to those who use the measure.

### 4. Usability and Use: H-10; M-3; L-0; I-0

*(Used and useful to the intended audiences for 4a. Accountability and Transparency; 4b. Improvement; and 4c. Benefits outweigh evidence of unintended consequences)*

#### Rationale:

- The Committee asked the developer to explain the measure selection process for recipients of RWHAP grant funds. The developer clarified that RWHAP does not specify which measures recipients must use.
- The Committee discussed that the measure is the most useful to determine how to schedule patients, with Committee members anecdotally sharing they find the measure valuable.
- A Committee member questioned how the measure is implemented in paper records. Other Committee members responded that smaller providers with paper records have to manually count the measure and that larger centers with paper records may find the manual implementation of the measure to be difficult.

### 5. Related and Competing Measures

- This measure is related to the following:
  - 2079 HIV Medical Visit Frequency
  - 2082 HIV Viral Suppression
  - 2083 Prescription of HIV Antiretroviral Therapy
  - 3211 Prescription of HIV Antiretroviral Therapy
  - 3210 HIV Viral Suppression
  - 3209 HIV Medical Visit Frequency
  - 0405 HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis
  - 0409 HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis
- Per the developer, the measure is harmonized with the first six measures listed above. For these six measures the target population is the same (i.e., people living with HIV), however the measure focus is different (medical visit frequency, prescription of ARV therapy, and viral load suppression).
- NQF #0405 and #0409 have been deferred for maintenance of endorsement. There are no additional steps the developer must take since the measure focus is different (HIV patients receiving PCP prophylaxis and those screened for STDs).

Standing Committee Recommendation for Endorsement: **Y-13; N-0**

### 6. Public and Member Comment

- [One comment was submitted in support of the Committee's recommendation for endorsement.](#)

<b>2080 Gap in HIV Medical Visits</b>
<b>7. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X</b>
<b>8. Board of Directors Vote: Y-X; N-X</b>
<b>9. Appeals</b>

## 2083 Prescription of HIV Antiretroviral Therapy

[Submission](#) | [Specifications](#)

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV prescribed antiretroviral therapy for the treatment of HIV infection during the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.

**Numerator Statement:** Number of patients from the denominator prescribed HIV antiretroviral therapy during the measurement year.

**Denominator Statement:** Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the measurement year.

**Exclusions:** There are no patient exclusions.

**Adjustment/Stratification:**

**Level of Analysis:** Facility

**Setting of Care:** Clinician Office/Clinic

**Type of Measure:** Process

**Data Source:** Other, Paper Records, Pharmacy

**Measure Steward:** Health Resources and Services Administration - HIV/AIDS Bureau

### STANDING COMMITTEE MEETING [03/14/2017]

#### 1. Importance to Measure and Report: The measure meets the Importance criteria

(1a. Evidence, 1b. Performance Gap)

1a. Evidence: **Accepted previous evaluation**; 1b. Performance Gap: **H-12; M-0; L-0; I-0**;

Rationale:

- A Committee member noted that the evidence base for this measure addresses viral suppression but not the prescription of antiretroviral (ARV) therapy, which is the measure's focus. The Committee member pointed out that they should consider whether the mere *prescription* of antiretroviral therapy rather than the *receipt* of the therapy by the patient is adequate evidence.
- The developer noted that there is not a way to measure whether patients are picking up their medications from the pharmacy and that this measure is used in tandem with #2082.
- The Committee acknowledged that the measure is based on a strong recommendation for ARV (i.e., that the patient is taking the medication) but noted there is a difference between writing a prescription (i.e., the focus of this measure) and a patient actually receiving the medication. The Committee stated that although the measure is valuable, there does not seem to be a link between *prescribing* ARV and viral suppression.
- Ultimately, the Committee decided not to re-vote on the evidence and accepted the previous evaluation.
- From 2010 to 2014, measure performance improved from 68.4% to 77.6%, with a median score of 90% in 2014. The Committee reviewed data that showed disparities among age, gender, and racial/ethnic groups. Overall, the Committee agreed the measure met this criterion.

#### 2. Scientific Acceptability of Measure Properties: The measure does meet the Scientific Acceptability criteria

(2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity)

2a. Reliability: **Accepted prior evaluation** 2b. Validity: **Accepted prior evaluation**

Rationale:

- The Committee reviewed testing data from the Ryan White HIV/AIDS Program Services Report (RSR) (covering 2000 RWHAP recipients) showing reliability at 0.99 and found the testing method appropriate; without further discussion the Committee accepted the previous evaluation on this criterion.
- The Committee noted that the measure is abstracted from paper and electronic records and questioned why the measure does not exclude patient death. The developer sited that some providers



<b>2083 Prescription of HIV Antiretroviral Therapy</b>
<p>are starting to use electronic health records and noted that death was not an exclusion for this measure because HIV mortality is extremely low and the increased burden of data collection and analysis does not add value to the measure.</p> <ul style="list-style-type: none"> <li>The Committee accepted the previous evaluation for reliability and validity.</li> </ul>
<p><b>3. Feasibility: H-11; M-1; L-0; I-0</b>  <i>(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/unintended consequences identified 3d. Data collection strategy can be implemented)</i>  <u>Rationale:</u></p> <ul style="list-style-type: none"> <li>The Committee noted the data elements are in electronic records, are in use and are readily available. Without further discussion, the Committee agreed the measure met this criterion.</li> </ul>
<p><b>4. Usability and Use: H-9; M-3; L-0; I-0</b>  <i>(Used and useful to the intended audiences for 4a. Accountability and Transparency; 4b. Improvement; and 4c. Benefits outweigh evidence of unintended consequences)</i>  <u>Rationale</u></p> <ul style="list-style-type: none"> <li>The Committee acknowledges that the measure is in use and is publicly reported in PQRS and MIPS. The Department of Health and Human Services selected this measure as a core HIV indicator. The Committee then agreed that the measure met this criterion.</li> </ul>
<p><b>5. Related and Competing Measures</b></p> <ul style="list-style-type: none"> <li>This measure is related to the following: <ul style="list-style-type: none"> <li>2079 HIV Medical Visit Frequency</li> <li>2080 Gap in HIV Medical Visits</li> <li>2082 HIV Viral Suppression</li> <li>3211 Prescription of HIV Antiretroviral Therapy</li> <li>3210 HIV Viral Suppression</li> <li>3209 HIV Medical Visit Frequency</li> <li>0405 HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis</li> <li>0409 HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</li> </ul> </li> <li>Per the developer, the measure is harmonized with the first six measures listed above. For these six measures the target population is the same (i.e., people living with HIV), however the measure focus is different (gaps in visits, medical visit frequency, and viral load suppression).</li> <li>NQF #0405 and #0409 have been deferred for maintenance of endorsement. There are no additional steps the developer must take since the measure focus is different (HIV patients receiving PCP prophylaxis and those screened for STDs).</li> </ul>
Standing Committee Recommendation for Endorsement: <b>Y-12; N-0</b>
<p><b>6. Public and Member Comment</b></p> <ul style="list-style-type: none"> <li><a href="#">One comment was submitted in support of the Committee's recommendation for endorsement.</a></li> </ul>
<b>7. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X</b>
<b>8. Board of Directors Vote: Y-X; N-X</b>
<b>9. Appeals</b>

## 3211 Prescription of HIV Antiretroviral Therapy

[Submission](#) | [Specifications](#)

**Description:** Percentage of patients, regardless of age, with a diagnosis of HIV prescribed antiretroviral therapy for the treatment of HIV infection during the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.

**Numerator Statement:** Number of patients from the denominator prescribed HIV antiretroviral therapy during the measurement year.

**Denominator Statement:** Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the measurement year.

**Exclusions:** There are no patient exclusions.

**Adjustment/Stratification:** No risk adjustment or risk stratification

**Level of Analysis:** Facility

**Setting of Care:** Clinician Office/Clinic

**Type of Measure:** Process

**Data Source:** Electronic Health Record (Only)

**Measure Steward:** Health Resources and Services Administration - HIV/AIDS Bureau

### STANDING COMMITTEE MEETING [03/14/2017]

#### 1. Importance to Measure and Report: The measure meets the Importance criteria

(1a. Evidence, 1b. Performance Gap)

1a. Evidence: **Applied the vote from #2083**; 1b. Performance Gap: **Applied the vote from #2083**

##### Rationale:

- The Committee acknowledged that this measure shares the same evidence as #2083. Because it is a legacy eMeasure, there is no performance data. The Committee agreed to apply the vote from #2083 to this criterion.

#### 2. Scientific Acceptability of Measure Properties: The measure does meet the Scientific Acceptability criteria

(2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity)

2a. Reliability: **H-0; M-12; L-0; I-0** 2b. Validity: **H-0; M-12; L-0; I-0**

##### Rationale:

- This measure was tested using 34 synthetic cases with the Bonnie tool. Results showed 100% coverage and all 34 cases passed the measure as expected, demonstrating that the measure logic works as constructed. Testing also included specific ways to search for patients that might fall at the edge of the measure specifications.
- The Committee agreed that the measure specifications are consistent with the evidence and noted that a panel of experts looked at each synthetic cases and assigned an outcome which then correlated with the results of the Bonnie testing, which demonstrates that the measure logic works.

#### 3. Feasibility: **H-9; M-3; L-0; I-0**

(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/unintended consequences identified 3d. Data collection strategy can be implemented)

##### Rationale:

- The Committee acknowledged measure feasibility at 98.33% and noted that data elements are in electronic records, are in use and are readily available. The Committee agreed the measure met this criterion.

#### 4. Usability and Use: **H-7; M-5; L-0; I-0**

(Used and useful to the intended audiences for 4a. Accountability and Transparency; 4b. Improvement; and 4c. Benefits outweigh evidence of unintended consequences)

## 3211 Prescription of HIV Antiretroviral Therapy

### Rationale

- The Committee acknowledges that the measure usability is similar to #2083 and without further discussion agreed the measure met this criterion.

### **5. Related and Competing Measures**

- This measure is related to the following:
  - 2079 HIV Medical Visit Frequency
  - 2080 Gap in HIV Medical Visits
  - 2082 HIV Viral Suppression
  - 3210 HIV Viral Suppression
  - 3209 HIV Medical Visit Frequency
  - 0405 HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis
  - 0409 HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis
- Per the developer, the measure is harmonized with the first six measures listed above. For these six measures the target population is the same (i.e., people living with HIV), however the measure focus is different (gaps in visits, medical visit frequency, and viral load suppression).
- NQF #0405 and #0409 have been deferred for maintenance of endorsement. There are no additional steps the developer must take since the measure focus is different (HIV patients receiving PCP prophylaxis and those screened for STDs).

Standing Committee Recommendation for Endorsement: **Y-12; N-0**

### **6. Public and Member Comment**

- [One comment was submitted in support of the Committee's recommendation for endorsement.](#)

**7. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X**

**8. Board of Directors Vote: Y-X; N-X**

**9. Appeals**

## 3215 Adult Inpatient Risk Adjusted Sepsis Mortality

### [Submission](#) | Specifications

**Description:** Annual risk adjusted inpatient mortality rate for adult patients (aged 18 and over) admitted to acute care hospitals with diagnosis of severe sepsis or septic shock. The measure includes patients in acute care hospital settings over one year timeframe who had, either on admission, or during their hospital stay, a clinical diagnosis of severe sepsis (now referred to as 'sepsis') or septic shock using criteria described in the International Sepsis Definitions (Sepsis-2)

Hospitals were required to submit a protocol for early identification and treatment of severe sepsis or septic shock. Subsequent to protocol submission, hospitals were required to submit 100% of their patient cases to a data collection portal using a standardized data dictionary (see relevant sections for details). Numerous data elements including patient demographics and comorbidities among other patient care details were reported. A random sample of the data submissions were validated for accuracy. The full adult data for discharges within calendar year 2015 was used to generate statewide and hospital-specific risk adjusted mortality rates for the calendar year.

**Numerator Statement:** Outcome is risk adjusted inpatient mortality rate for adult patients (18 and over) admitted to an acute care hospital with a diagnosis of severe sepsis or septic shock or who develop severe sepsis or septic shock during their hospital stay.

**Denominator Statement:** All adult patient discharges (18 and over) in a calendar year with a diagnosis of severe sepsis or septic shock on admission or at any time during their hospital stay. This may include multiple admissions of the same patient during the measurement year. Denominator includes all cases identified using any means (administrative, registry, electronic health records, billing data, etc.), either prospectively, retrospectively, or both, that meet the International consensus definition (Sepsis- 2) of severe sepsis or septic shock.

**Exclusions:** Patients with advanced directives in place prior to episode of sepsis which specifically restrict any hospital specific sepsis protocol interventions or who decline (or their proxy declines) treatment for sepsis. Patients who have been transferred from one acute care hospital to another are excluded.

**Adjustment/Stratification:** Multivariate logistic regression model

**Level of Analysis:** Facility

**Setting of Care:** Hospital : Acute Care Facility

**Type of Measure:** Outcome

**Data Source:** Claims (Other), EHRs Hybrid, Laboratory, Management Data, Non-Medical Data, Paper Records, Pharmacy, Registry

**Measure Steward:** New York State Department of Health, Office of Quality and Patient Safety

### STEERING COMMITTEE MEETING [03/14/2017]

#### 1. Importance to Measure and Report: The measure meets the Importance criteria

(1a. Evidence, 1b. Performance Gap)

1a. Evidence: **Y-16; N-0**; 1b. Performance Gap: **H-12; M-3; L-0; I-1**

#### Rationale:

- Mortality is an important outcome for patients with sepsis and according to the developer, mortality rates are high and show significant variability across acute care hospitals unrelated to patient factors.
- The developer suggested that hospitals are able to influence mortality rates using early sepsis detection approaches coupled with rapid delivery of basic resuscitation interventions including the use of adequate intravenous fluids, antibiotics, blood pressure support medications and dynamic clinical monitoring for response.
- The Committee agreed that the developer clearly identified how healthcare facilities and providers influence sepsis mortality outcomes.
- The developer provided the risk-adjusted probability of inpatient sepsis mortality rates from 179 hospitals and 43,204 patients in New York State from January 1, 2015 – December 31, 2015 for this newly developed measure. The mean performance rate in Q1 2015 was 30.4%, 28.9% in Q2 2015,

### 3215 Adult Inpatient Risk Adjusted Sepsis Mortality

28.8% in Q3 2015 and 28.4% in Q5 2015. The performance rates in 2015 ranged from a minimum of 1.0% (Q1-Q4) to a maximum of 95.0% (Q4). The developer also provided the probability of inpatient sepsis mortality rates by population group, which included race/ethnicity, gender, age and insurance/payer. White, non-Hispanics had a rate of 28.2%; Black, non-Hispanics had a rate of 31.5%; and Hispanics had a rate of 26.3%. Rates based on gender were similar with 28.4% for females and 29.8% for males. Patients 70 to 80+ years old had rates from 31.0% to 33.9%. The probability of inpatient sepsis mortality also varied based on insurance/payer. Patients with Medicare had a rate of 30.6%; Medicaid patients had a rate of 26.2%; Private pay and/or HMO patients had a rate of 27.1%; self-pay patients had the highest probability of inpatient sepsis mortality, 34.3%.

- The Committee recognized the variability in coding practices related to sepsis and the potential impact on the denominator of this measure. After a lengthy discussion, the Committee agreed that the data presented by the developer demonstrated significant variation and an opportunity for improvement in inpatient sepsis mortality across hospitals.

#### **2. Scientific Acceptability of Measure Properties: The measure meets the Scientific Acceptability criteria**

(2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity)

2a. Reliability: **H-2; M-8; L-0; I-6** 2b. Validity: **H-4; M-8; L-1; I-3**

##### Rationale:

- The developer conducted data element validity testing, which counted for data element reliability testing as well.
- The dataset included hospitals in New York State that were required to develop and implement early recognition and treatment protocols for sepsis. As part of this statewide initiative, hospitals were required to submit quarterly clinical data to the New York State Department of Health to be evaluated for protocol use, adherence to time interventions and patient outcomes, including mortality.
  - The dataset included 179 hospitals with 43,204 patients diagnosed with severe sepsis and septic shock from January 1, 2015 to December 31, 2015.
  - The dataset used to develop the logistic regression model included:
    - Development sample: 38,884 (90%) patients; 179 hospitals
    - Validation sample: 4,319 (10%) patients; 160 hospitals
- The developer validated the accuracy of the data submission from the hospitals against manual chart abstraction by external auditors (Audit Results), which is considered the gold standard. The developer calculated the percent agreement between the hospital submissions to the chart-abstracted data. Percent agreement from the audit results ranged from 89.9% to 99.1% for the following data elements:
  - Site of infection: 98.9%
  - Lower Respiratory Infection: 98.8%
  - Mechanical Ventilation: 97.8%
  - Age (Date of Birth): 99.2%
  - Thrombocytopenia: 97.7%
  - Septic Shock: 98.4%
  - Serum Lactate (Lactate Level): 93.9%
  - Metastatic Cancer: 97.1%
  - Lymphoma, Leukemia, Multiple Myeloma: 99.1%
  - Square Root of Comorbidity Count (Range of Comorbidities): 89.9% - 99.1%
- The developer noted that the data elements race, ethnicity, payer and admission source were not manually audited but were aligned to state administrative datasets to ensure accuracy.
- The developer did not provide sensitivity, specificity, positive predictive value (PPV) and negative predictive values (NPV) in addition to percent agreement.
- The developer clarified that if a case was initially coded as severe sepsis or septic shock but the manual review found that the case did not meet the clinical definition of severe sepsis or septic shock, the hospital was able to exclude the case.
- A Committee member asked if the developer had compared patients that present to the emergency department (ED) with sepsis vs. patients that develop sepsis while hospitalized. The developer replied

### 3215 Adult Inpatient Risk Adjusted Sepsis Mortality

that they intend to continue researching the differences in outcomes for patients that present to the ED with sepsis vs. patients that acquire sepsis in the hospital.

- The Committee did not express any other concerns with the reliability of the measure and agreed it met the reliability criterion.
- Empirical validity testing of the measure score was assessed by comparing the performance of the risk-adjusted model in the development sample to the validation sample.
  - The developer used the Hosmer-Lemeshow goodness of fit test to assess the observed and expected mortality rates in the development and validation samples.
    - The development dataset was split into group sizes of 10, 100, 500 and 1,000. The p-values were 0.568, 0.972, 0.735, and 0.735, respectively.
    - The validation dataset was split into group sizes of 10, 50, 100 and 150. The p-values were 0.651, 0.977, 0.985, and 0.974, respectively.
- The performance of the risk-adjustment model was similar in the development and validation datasets. The areas under the receiver operating characteristic (ROC) curve (or c-statistic) were **0.770** and **0.773**, respectively. A c-statistic is a model of discrimination statistic. A c-statistic of 0.77 means that 77.0% of all possible pairs of patients – one who died and one who lived – the model correctly assigned a higher probability to those who died. The similar c-statistics indicates good model discrimination.
- A Committee member questioned whether a c-statistic of 0.7 was sufficient. NQF staff responded that although NQF staff does not have specific statistical thresholds, generally, a c-statistic of at least 0.70 is considered acceptable.
- Exclusions include patients with advanced directives in place prior to episode of sepsis which specifically restrict any hospital specific sepsis protocol interventions or who decline (or their proxy declines) treatment for sepsis and patients who have been transferred from one acute care hospital to another. The developer maintained that keeping patients in the dataset that had an advanced directive or declined intervention would bias the results towards higher hospital mortality.
- In the pre-evaluation comments, one of the Committee members questioned why excluding patients with multiple admissions is appropriate because this would artificially increase the mortality rate; however, patients with multiple admissions are not excluded from the denominator. The denominator details state that multiple admissions of the same patient during the measurement year are included.
- This measure is risk-adjusted using a multivariable logistic regression model with 16 variables to estimate the probability of mortality for patients admitted to acute care hospitals with severe sepsis or septic shock. The model was built using the development dataset and starting with all possible covariates in the model. Using an iterative procedure, variables were removed from the model, one by one, if the p-values were not significant at 0.05 level until a parsimonious model was reached.
  - Variables removed during the development procedure were added back into the reduced model if the p-values were significant at the 0.05 level and if model calibration (Hosmer-Lemeshow goodness of fit) was improved through their inclusion.
  - The scale of the three continuous variables (patient age, first serum lactate, and the count of the number of comorbidities) remaining in the model was assessed. Using the method of fractional polynomials patient age was included in the model as a linear term, number of comorbidities was transformed by taking the square root, and first serum lactate was entered into the model as a quadratic expression (linear and a squared term).
  - Model calibration was further improved by adding the following interactions to the model: lower respiratory infection (LRI) by MV severity, patient age by the square root of the number of comorbidities, and first serum lactate by the square root of the number of comorbidities.
  - Age, gender, payer, race and ethnicity were initially included in the model. Gender was the only variable included in the model since its odds ratio and corresponding p-value was 1.0003 and 0.992, respectively. All of the other demographic variables had p-values < 0.001 for at least one of the levels of a specific demographic.
- The developer stated that the intent of this risk model is to estimate the *probability* of mortality due to sepsis not *predict* mortality rates due to sepsis. By estimating the probability, the developer continued, the expected number of events (sepsis mortality) for each hospital is calculated. Variables

### 3215 Adult Inpatient Risk Adjusted Sepsis Mortality

must be clinically and statistically significant to be included in the risk model. The Committee recognized that the developer deliberately excluded hospital characteristics from the variables in the risk model because organizational variables do modify the probability of mortality for patients.

- One of the Committee members stated that they would like to see a statistical analysis, such as a funnel plot, demonstrating outliers (i.e., # of hospitals, hospital size and statistical threshold). A funnel plot is helpful in illustrating real variation among hospitals vs. noise.
- Ultimately, the Committee agreed the measure met the validity criterion.

#### 3. Feasibility: H-1; M-11; L-3; I-1

*(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/unintended consequences identified 3d. Data collection strategy can be implemented)*

##### Rationale:

- The developer noted that some data elements are in defined fields in electronic sources, and that some demographic variables can be extracted electronically and used in a standard format. Other variables are collected manually by hospitals though some hospitals have created electronic data capture avenues.
- During the workgroup call, the Committee discussed the amount of manual chart abstraction that would be required to collect the data for this measure. The Committee acknowledged the challenges related to the feasibility of this measure but agreed that it was not impossible.

#### 4. Usability and Use: H-2; M-10; L-1; I-3

*(Used and useful to the intended audiences for 4a. Accountability and Transparency; 4b. Improvement; and 4c. Benefits outweigh evidence of unintended consequences)*

##### Rationale:

- This measure is in use in New York State. New York State requires the Department of Health to collect and report data regarding the performance of hospitals for patients with sepsis including risk adjusted mortality rates for individual hospitals.
- The Committee agreed that hospitals in other states could use this measure to track sepsis mortality outcomes.

#### 5. Related and Competing Measures

- This measure is related to:
  - #0500: Severe Sepsis and Septic Shock: Management Bundle
- The developer stated the measure specifications are harmonized to the extent possible.

#### Standing Committee Recommendation for Endorsement: Y-11; N-5

#### 6. Public and Member Comment

- [One comment was submitted in support of the Committee's recommendation for endorsement.](#)

#### 7. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X

#### 8. Board of Directors Vote: Y-X; N-X

#### 9. Appeals



## 0500 Severe Sepsis and Septic Shock: Management Bundle

### [Submission](#) | Specifications

**Description:** This measure focuses on adults 18 years and older with a diagnosis of severe sepsis or septic shock. Consistent with Surviving Sepsis Campaign guidelines, the measure contains several elements, including measurement of lactate, obtaining blood cultures, administering broad spectrum antibiotics, fluid resuscitation, vasopressor administration, reassessment of volume status and tissue perfusion, and repeat lactate measurement. As reflected in the data elements and their definitions, these elements should be performed in the early management of severe sepsis and septic shock.

**Numerator Statement:** The number of patients in the denominator who received ALL of the following components (if applicable) for the early management of severe sepsis and septic shock: initial lactate levels, blood cultures, antibiotics, fluid resuscitation, repeat lactate level, vasopressors, and volume status and tissue perfusion reassessment.

- Within 3 hours of presentation of severe sepsis:
  - Measure initial lactate level
  - Draw blood cultures prior to antibiotics
  - Administer broad spectrum or other antibiotics
- Within 6 hours of presentation of severe sepsis:
  - Repeat lactate level (if initial lactate > 2 mmol/L)
- Within 3 hours of presentation of septic shock:
  - Administer 30 ml/kg crystalloid for hypotension or lactate  $\geq 4$  mmol/L
- Within 6 hours of presentation of septic shock:
  - Apply vasopressors (for hypotension that does not respond to initial fluid resuscitation) to maintain a mean arterial pressure (MAP)  $\geq 65$  mm Hg
  - Reassess volume status and tissue perfusion in the event of persistent hypotension (MAP <65 mm Hg) after initial fluid administration or initial lactate level  $\geq 4$  mmol/L
    - The clinician is no longer required to state the method of reassessment used (e.g. physical exam, bedside cardiovascular ultrasound, passive leg raising, CVP, ScVO<sub>2</sub> assessment). The clinician can attest that volume and perfusion reassessment has occurred, even without reference to the method used. This will meet the measure's volume and perfusion reassessment requirement. A provider may also opt to state their chosen method, but this is not required.

**Denominator Statement:** Inpatients age 18 and over with an ICD-10-CM Principal or Other Diagnosis Code of Sepsis, Severe Sepsis, or Septic Shock.

**Exclusions:** The following patients are excluded from the denominator:

- Severe sepsis is not present
- Patients Transferred in from another acute care facility
- Patients receiving IV antibiotics for more than 24 hours prior to presentation of severe sepsis.
- Patients with a Directive for Comfort Care or Palliative Care within 3 hours of presentation of severe sepsis
- Patients with an Administrative Contraindication to Care within 6 hours of presentation of severe sepsis
- Patients with an Administrative Contraindication to Care within 6 hours of presentation of septic shock
- Patients with a Directive for Comfort Care or Palliative Care within 6 hours of presentation of septic shock
- Patients with septic shock who are discharged within 6 hours of presentation
- Patients with severe sepsis who are discharged within 6 hours of presentation
- Patients with a Length of Stay >120 days



## 0500 Severe Sepsis and Septic Shock: Management Bundle

- Patients included in a Clinical Trial

**Adjustment/Stratification:** No risk adjustment or risk stratification

**Level of Analysis:** Facility

**Setting of Care:** Hospital

**Type of Measure:** Composite

**Data Source:** Imaging-Diagnostic, Laboratory, Other, Paper Records, Pharmacy

**Measure Steward:** Henry Ford Hospital

### STEERING COMMITTEE MEETING [03/14/2017]

#### 1. Importance to Measure and Report: The measure meets the Importance criteria

(1a. Evidence, 1b. Performance Gap)

1a. Evidence: **H-4; M-9; L-0; I-2**; 1b. Performance Gap: **H-14; M-0; L-1; I-0**; 1c. Composite – Quality Construct and Rationale: **H-5; M-7; L-1; I-2**

#### Rationale:

- For the 2012 endorsement-maintenance evaluation, the developer provided the 2008 Surviving Sepsis Campaign guidelines with recommendations for initial resuscitation, measuring lactate, obtaining appropriate blood cultures, antibiotic therapy, fluid therapy, vasopressors and monitoring central venous pressure (CVP) and central venous oxygen saturation (ScvO<sub>2</sub>).
- In 2012, concerns were raised about the level of evidence supporting invasive monitoring of CVP and ScvO<sub>2</sub>. The Infectious Disease Steering Committee acknowledged these concerns yet determined that the current evidence at the time was sufficient to warrant endorsement of the full bundle, and the measure was approved as specified. NQF received an appeal and the Consensus Standards Approval Committee (CSAC) upheld the measure's endorsement with the condition that NQF commit to an immediate re-evaluation of the measure upon release of new evidence from several ongoing studies including the Protocolized Care for Early Septic Shock (ProCESS) trial.
- In 2014, the Patient Safety Standing Committee conducted an *ad hoc* review based on a request from the American College of Emergency Physicians (ACEP). The *ad hoc* review focused on the evidence supporting CVP and ScvO<sub>2</sub> and the new data from the ProCESS trial. See [NQF-Endorsed Measures for Patient Safety \(January 30, 2015\)](#) for complete summary.
  - The ProCESS trial demonstrated no difference in mortality outcomes when using an invasive approach to monitoring CVP and ScvO<sub>2</sub> compared to usual care or protocolized care without invasive monitoring. The Committee noted that the new results from the ProCESS trial suggested that a mandate to measure CVP and ScvO<sub>2</sub> with an invasive line might not be necessary in all patients with severe sepsis and septic shock.
  - Experts in support of maintaining these elements in the measure argued additional trials (ARISE and PROMISE) were underway; however, these trials were smaller than the ProCESS trial and not performed in the U.S. In addition, these experts argued that the protocolized care and requirement for CVP and ScvO<sub>2</sub> monitoring was particularly helpful in community hospitals, which were not included in the ProCESS trial.
  - After extensive discussions and negotiations the measure developers, ProCESS trial investigators and specialty societies (including the Society of Critical Care Medicine (SCCM) and the American College of Emergency Physicians (ACEP)) reached a compromise for an evidence-based replacement element – optional measurement of CVP and ScvO<sub>2</sub>, along with reassessment by other means (re-assess volume status and tissue perfusion after initial resuscitation and document findings).
- For the current maintenance of endorsement evaluation, the developer provided the following updated evidence to support the changes to the measure since the last submission:
  - The developer provided a synthesis of the literature for the following updated components, which are based on the [Surviving Sepsis Campaign: International Guidelines for Management](#)

## 0500 Severe Sepsis and Septic Shock: Management Bundle

[of Sepsis and Septic Shock: 2016](#). [Note: Grading of recommendations for the components below are taken from the Sepsis and Septic Shock 2016 Guidelines]

- Measure lactate level; Repeat lactate if initial lactate is elevated [*weak recommendation, low quality of evidence*<sup>1</sup>]
- Obtain cultures prior to antibiotics [*Best practice statement*]
- Administer broad spectrum antibiotics [*strong recommendation, moderate quality of evidence*]
- Administer 30 ml/kg crystalloid for hypotension or lactate  $\geq 4$  mmol/L [*strong recommendation, low quality of evidence*]
- Apply vasopressors (for hypotension that does not respond to initial fluid resuscitation to maintain a mean arterial pressure  $\geq 65$  mmHg) [*strong recommendation, moderate quality of evidence*]
- Reassess volume status and tissue perfusion [*Best practice statement*]. The developer provided a synthesis of the literature for some common practices used when reassessing volume status and tissue perfusion in the event of persistent hypotension (MAP  $< 65$  mm Hg) after initial fluid administration or initial lactate level  $\geq 4$  mmol/L. [Note: Clinician is no longer required to document the method used; attestation is sufficient]
- One of the Committee members questioned the use of lactic levels to diagnose sepsis because other conditions can elevate lactate levels. The developer responded that measuring lactate levels alone does not diagnose sepsis. However, in the presence of a suspected infection, elevated lactate levels are useful in determining illness severity. The Committee pointed out the varying level of evidence for the different components but agreed that overall, the updated evidence is consistent with the 2016 Guidelines for Management of Sepsis and Septic Shock.
- The developer provided the following composite performance rates from CMS' Hospital Inpatient Quality Reporting (IQR) program from October 2015 to June 2016:

	Oct-Dec 2015	Jan-Mar 2016	Apr-Jun 2016
# of hospitals	3,134	3,182	3,193
# of eligible cases	96,516	104,166	101,599
Overall performance rate	34.4	39.5	44.0
10 <sup>th</sup> percentile	5.0	7.7	12.5
25 <sup>th</sup> percentile	17.9	21.6	25.8
Median	31.0	36.1	41.7
75 <sup>th</sup> percentile	45.8	51.3	57.1
90 <sup>th</sup> percentile	60.0	66.7	71.4
Min, Max	0.0, 100.0	0.0, 100.0	0.0, 100.0
Average	32.6	37.1	41.9
Standard Deviation	21.1	21.9	22.9

- The developer also provided the following component rates categorized by 3 and 6 hour elements:

Population Description	Cases	Bundle %
Initial Population Number of Sepsis Cases	325,809	--

<sup>1</sup> Per guideline authors, 'low' grade assigned to quality of evidence (5 RCTs) because 1) all studies were judged to be at high risk of bias due to lack of clarity of the intervention, therefore, we downgraded the quality of evidence by one level for risk of bias; 2) We downgraded the quality of evidence by one level for imprecision, the CI contained small benefit that was lower than the decision threshold; and 3) We assumed a mortality rate for patients with septic shock to be 40%.

## 0500 Severe Sepsis and Septic Shock: Management Bundle

Total Number of Excluded Sepsis Cases	166,520	--
<b>Total Number of Eligible Sepsis Cases</b>	<b>159,289</b>	<b>--</b>
Total Number of Passed Sepsis Cases	64,051	--
Total Number of Failed Sepsis Cases	95,238	--
<b>Severe Sepsis 3 Hour Bundle Eligible Cases</b>	<b>167,114</b>	<b>--</b>
Severe Sepsis 3 Hour Bundle Passes	110,078	65.9%
Severe Sepsis 3 Hour Bundle Failures	54,618	32.7%
Initial Lactate Level Failures	26,503	48.5%
Broad Spectrum or Other Antibiotic Administration Failures	20,951	38.4%
Blood Culture Collection Failures	18,772	34.4%
<b>Severe Sepsis 6 Hour Bundle Eligible Cases</b>	<b>90,385</b>	<b>--</b>
Severe Sepsis 6 Hour Bundle Passes	53,475	59.2%
Severe Sepsis 6 Hour Bundle Failures	36,910	40.8%
Repeat Lactate Level Failures	36,910	
<b>Septic Shock 3 Hour Bundle Eligible Cases</b>	<b>40,989</b>	<b>--</b>
Septic Shock 3 Hour Bundle Passes	22,359	54.5%
Septic Shock 3 Hour Bundle Failures	18,630	45.5%
Crystalloid Fluid Administration Failures	18,630	
<b>Vasopressor Shock 6 Hour Bundle Eligible Cases</b>	<b>8,177</b>	<b>--</b>
Vasopressor Shock 6 Hour Bundle Passes	6,157	75.3%
Vasopressor Shock 6 Hour Bundle Failures	2,020	24.7%
Vasopressor Administration Failures	2,020	
<b>Focus Exam Shock 6 Hour Bundle Eligible Cases</b>	<b>14,630</b>	<b>--</b>
Focus Exam Shock 6 Hour Bundle Passes	3,801	26.0%
Focus Exam Shock 6 Hour Bundle Failures	9,935	67.9%
<b>Hemodynamic Choices Shock 6 Hour Bundle Eligible Cases</b>	<b>10,829</b>	<b>--</b>
Hemodynamic Choices Shock 6 Hour Bundle Passes	894	8.3%
Hemodynamic Choices Shock 6 Hour Bundle Failures	9,935	91.7%

- The developer noted that the 'repeat volume' and 'perfusion assessment' data is broken down into 'focused exam' and 'hemodynamic choice' – data elements that are no longer required. No performance data is yet available on the new attestation strategy.
- The developer also provided the following composite performance rates by ethnicity, gender and Medicare/non-Medicare:

	Oct-Dec 2015	Jan-Mar 2016	Apr-Jun 2016
<b>Hispanic</b>	34.53	39.8	44.23
<b>Non-Hispanic</b>	32.64	36.35	40.82
<b>Females</b>	33.82	39.22	43.28

## 0500 Severe Sepsis and Septic Shock: Management Bundle

<b>Males</b>	34.93	39.86	44.63
<b>Medicare</b>	34.9	40.07	44.57
<b>Non-Medicare</b>	33.32	38.47	42.76
<b>Black</b>	30.64	35.93	40.29
<b>White</b>	34.95	40.08	44.58
<b>Other</b>	34.95	40.01	43.82

- The performance rates for different age categories were similar (~34.0).
- The Committee agreed that the developer presented abundant data demonstrating a performance gap and opportunity for improvement in severe sepsis and septic shock care.
- Another Committee member commented that in addition to a performance gap in care, there is still a lack of implementation of this measure in the clinical setting.
- This all-or-none composite measure requires patients with severe sepsis or septic shock to meet all of the eligible components in the composite: lactate collection, delivery of broad-spectrum antibiotics, obtaining blood cultures, delivering resuscitation fluids, applying vasopressors as needed, reassessing volume and perfusion status and repeating lactate values. All components are equally weighted. The developer noted that the composite ensures a strategy aimed at reducing mortality. The components could not stand alone unless certain preceding conditions have been met. In addition, the components are aggregated in three and 6-hour elements for severe sepsis and septic shock.
- A Committee member suggested that the data elements should be weighted differently based on the level of evidence for each. The Committee member also questioned the two different time periods. The developer responded that severe sepsis and septic shock are two different diagnoses that qualify for this measure. Therefore, the measure is constructed so that there are dependencies, both in time and condition, based on the diagnosis of severe sepsis or septic shock. The severity of the patient's condition determines whether they qualify to receive all of the components in the composite measure and when. The Committee agreed, that overall, the quality construct and rationale for the composite was clearly stated and logical.

### 2. Scientific Acceptability of Measure Properties: The measure meets the Scientific Acceptability criteria

(2a. Reliability - precise specifications, testing; 2b. Validity - testing, threats to validity)

2a. Reliability: **H-7; M-7; L-0; I-1** 2b. Validity: **H-0; M-11; L-1; I-3**

#### Rationale:

- For the 2012 endorsement evaluation, the developer conducted a signal-to-noise analysis of the individual bundle elements and overall bundle reliability and composite measure reliability by site for 165 hospitals and 15,022 patients from January 2005 – March 2008.
- Effective 2013, NQF determined that reliability of the individual component measures was not sufficient; reliability must be demonstrated for the composite measure score.
- For the current evaluation, the developer provided updated reliability testing at the composite score level using a random sample of SEP-1 chart-abstracted data submitted to CMS as part of the Hospital Inpatient Quality Reporting (IQR) program. The sample included 302,281 cases in the denominator (after exclusions) and 119,048 cases in the numerator from 3,134 to 3,193 hospitals (depending on the quarter) from October 2015 to June 2016.
- The developer used a beta-binomial model to assess the signal-to-noise ratio. A reliability of 0.0 implies that all the variability in a measure is attributable to measurement error. A reliability of 1.00 implies that all the variability is attributable to real differences in performance. The higher the reliability score, the greater is the confidence with which one can distinguish the performance of one provider from another. This is an appropriate test for measure score reliability. A reliability of 0.70 is generally considered a minimum threshold for reliability.
  - The median reliability score was calculated including all facilities and facilities with a minimum of 10 eligible cases (more than 86% of reporting facilities).
  - The developer provided the overall reliability score for the composite measure for each quarter:

## 0500 Severe Sepsis and Septic Shock: Management Bundle

- October 2015 – December 2015: 0.92 (CI 0.41 – 1.00)
  - January 2016 – March 2016: 0.93 (CI 0.47 – 1.00)
  - April 2016 – June 2016: 0.93 (CI 0.42 – 1.00)
- In the pre-evaluation comments, the Committee expressed some concerns that the self-reported clinician attestation for the data element ‘volume and perfusion reassessment’ may lead to more subjectivity but lauded the developers in their efforts to reduce documentation and chart abstraction burden.
- During the in-person meeting, the Committee did not voice any additional concerns related to reliability and agreed the reliability testing results were sufficient.
- For the 2012 endorsement evaluation, the developer assessed the validity of the measure score by testing the hypothesis that those with higher scores on the composite performance measure should have a lower score on a risk-adjusted mortality measure. The developer reported that hospital mortality was reduced by 10% for patients that were compliant with the composite measure.
- For the current evaluation, the developer provided updated validity testing at the composite score level. The developer performed a Chi-Square Test of Association and Equal Proportions between two categorical variables: Measure Outcome (Failed or Passed) and Mortality Result (Died or Survived). The Chi-Square Test of Association and Equal Proportions demonstrated a p-value of <.0001, a risk ratio of 1.3856, a lower 95% confidence limit of 1.3616 and an upper 95% confidence limit of 1.4101. A risk ratio higher than 1.0 with a significant p-value, would indicate that there is a higher risk of dying when a case fails the measure compared to when a case passes the measure.
  - The risk ratio can be expressed as an actual ratio and it can be said with 95% confidence, cases that fail the measure have 1.36 to 1.41 times the risk of dying compared to cases that pass the measure; or
  - The risk ratio can also be used as a percentage and be said that with 95% confidence, cases that fail the measure have a 36% to 41% increase in risk of dying compared to cases that pass the measure.
- The developer provided sepsis rate comparisons analyses, which demonstrated a negative association between pass rates and mortality rates from October 2015 to June 2016.
- The developer also included an analysis of pass rates and mortality rates by percentiles and a two-proportions z-test. The z-test determines if there is a statistically significant difference in mortality rates between percentiles. These methods are appropriate for empirically assessing the validity of the composite measure score. The results of the sepsis mortality analysis demonstrated that 30.4% of the total number of ‘Failed Sepsis Cases’ died (at discharge and up to 30 days after discharge) compared to 21.9% of the total number of ‘Passed Sepsis Cases’. The two-proportion z-test demonstrated that four of the percentile comparisons have a statistically significant difference between mortality rates at a significance level of 0.05. Three additional percentile comparisons are fairly close to a statistically significant difference between mortality rates at a significance level of 0.10.
- The developer also provided the mortality rate for patients who received all applicable elements of care for the composite measure (passed sepsis cases) and those who did not (failed sepsis cases) for each quarter – the mortality rate for those who received all applicable elements was on average 8.5% lower compared to patients who did not receive all applicable elements of care.

	Severe Sepsis and Septic Shock Mortality Rate		
Description	2015 Q4	2016 Q1	2016 Q2
Did not Meet Guidelines for SEP-1	29.6%	31.8%	29.7%
Met Guidelines for SEP-1	21.3%	23.0%	21.4%
Absolute Reduction Rate	8.3%	8.8%	8.3%
Relative Reduction Rate	28.04%	27.7%	27.9%
	Potential Preventable Deaths		

## 0500 Severe Sepsis and Septic Shock: Management Bundle

2,783

2,864

2,411

- The Committee discussed the results of the patient-level data element validity testing conducted by CMS. Several Committee members were concerned because out of the 55 data elements tested for validity, 15 data elements (27.27%) had a percent agreement higher than 90%. The remaining 40 data elements (72.73%) had a percent agreement lower than 90%. The developer stated that there have been numerous education and outreach efforts and updates to the measure with the intent of clarifying guidance and decreasing abstractor complexities in an effort to improve successive validation testing. NQF staff noted that this method is not appropriate for composite measures. [NQF composite performance measure evaluation guidance \(2013\)](#) states that validity testing is directed toward the inferences that can be made about accountable entities on the basis of their performance measure scores. For the purposes of endorsing composite performance measures, validity testing of the constructed composite performance measure score is more important than validity testing of the component measures. Even if the individual component measures are valid, the aggregation and weighting rules for constructing the composite could result in a score that is not a true reflection of quality (p. 12-13).
- The developer stated that the number of exclusions was not significant enough to unfairly distort measure performance results and potentially negatively affect the reliability of the measure because the vast majority of the exclusions were cases where severe sepsis was not present (72.34%) and should not be analyzed.
- The developer provided the [table](#) included in performance gap to demonstrate the contribution of each component to the composite score.
- The Committee agreed that the measure met the validity criterion.

### 3. Feasibility: H-1; M-9; L-5; I-0

*(3a. Clinical data generated during care delivery; 3b. Electronic sources; 3c. Susceptibility to inaccuracies/unintended consequences identified 3d. Data collection strategy can be implemented)*

#### Rationale:

- Data elements are abstracted from a record by someone other than the person obtaining the original information (e.g. chart abstraction); some data elements are in defined fields in electronic sources.
- The developer noted that the measure is complex and requires data abstractors to “comb through documentation and interpret” clinician documentation; however, the most recent updates to the measure should lessen documentation and abstractor burden. The developer also stated that the measure has gone through three updates to lessen abstractor burden and address issues related to data availability, missing data, and frequency of data collection.
- The developer also stated that preliminary efforts to convert this measure to an electronic measure within the HQMF/QDM framework were not feasible. Currently, there are no plans to respecify this measure into an eMeasure.
- There are no fees or licenses required to use this measure.
- In the pre-evaluation comments and during the in-person meeting, the Committee noted that the feasibility of this measure was a significant concern. In the pre-evaluation comments one of the Committee members acknowledged that although clinicians routinely document the required data elements for this measure, it is complex and likely results in significant time and costs associated with data collection and reporting. During the in-person meeting, the developer clarified that it is the clinician’s responsibility to document appropriate care, therefore, decreasing the burden on the abstractors.
- The developer and CMS representatives reiterated that they regularly receive feedback from hospitals and their abstractors and are currently monitoring the most recent changes to the specifications and implementation guide.
- The Committee concluded that the feasibility of this measure is challenging but it meets the criteria.

### 4. Usability and Use: H-2; M-6; L-6; I-0 Consensus was not reached on the Usability and Use criteria



## 0500 Severe Sepsis and Septic Shock: Management Bundle

*(Used and useful to the intended audiences for 4a. Accountability and Transparency; 4b. Improvement; and 4c. Benefits outweigh evidence of unintended consequences)*

### Rationale:

- The measure is currently used in CMS' Hospital Inpatient Quality Reporting (IQR) Program for acute care hospitals nation-wide. Across the three quarters of available data, between 3,134 and 3,193 providers submitted data, which represents more than 95% of eligible providers nationwide.
- The measure is not currently publicly reported, but will be added to the Hospital Compare website at a date to be determined. Due to the complexity of the measure specifications, CMS desires to review and analyze the data prior to making it publicly available. There were also several updates to the specifications based on stakeholder feedback, and CMS wants to assure stability of the specifications before public reporting.
- The Committee discussed the unintended consequences associated with diagnosing and treating sepsis. Because sepsis is a combination of symptoms rather than a disease, the Committee noted, patients often receive antibiotics prior to a conclusive diagnosis of sepsis – this is similar to the unintended consequences that occurred with the overuse of antibiotics in the emergency room in an effort to meet the now retired pneumonia measure.
- At the end of the discussion, the Committee did not reach consensus on the usability and use of the measure during the in-person meeting.

### 5. Related and Competing Measures

- This measure is related to:
  - 3215: Adult Sepsis Mortality Outcome Measure (New York State Department of Health)

### Standing Committee Recommendation for Endorsement: Y-10; N-4

### 6. Public and Member Comment

Nine comments were submitted covering the following themes: antibiotic administration, level of evidence, and scientific acceptability.

- In regard to antibiotic administration, commenters noted the three hour time window for antibiotic administration may lead to antibiotic overuse; other comments questioned whether antibiotic administration rather than early goal directed therapy had a larger effect on patient survival rates; and another comment suggested that the developer reduce the antibiotic administration time window from three hours to two hours.
  - Developer Response to Comment ID 6678: We understand that measures can have unintended consequences for patients. We remain convinced that in severe sepsis (please refer to clinical criteria used for the measure) and septic shock the risk of mortality is so high that is critical to provide early and broad antibiotic therapy. The failure to provide a proper antibiotic for a patient with severe sepsis and septic shock carries a much higher risk than the potential harm of providing a single dose of a broad spectrum antibiotic to a patient who turns out not to have severe sepsis or septic shock. This in by no means precludes the clinician's responsibility to judiciously use IV antibiotics in a way which upholds the standards of antibiotic stewardship. The 2016 Surviving Sepsis Guidelines, endorsed by the Infectious Disease Society of America speak to this question: "The rapidity of [antimicrobial] administration is central to the beneficial effect of appropriate antimicrobials. In the presence of severe sepsis or septic shock, each hour delay in administration of appropriate antimicrobials is associated with a measurable increase in mortality." When mortality already approaches 18-40% in shock states, it is unacceptable to suspend antibiotic administration pending further studies. However, in the event an antibiotic is given inappropriately in non-sepsis states, the guidelines also recommend, "Given the potential harm associated with unnecessarily prolonged antimicrobial therapy, daily assessment for de-escalation of antimicrobial therapy is recommended in patients with severe sepsis and septic shock."
  - Developer Response to Comment ID 6680: The PRISM investigators have reported results in the New England Journal of Medicine (NEJM) entirely consistent with the prior Process,

## 0500 Severe Sepsis and Septic Shock: Management Bundle

Promise and Arise trials, also published in NEJM. Little information is imparted by PRISM that was not already known from these previous trials. In fact, PRISM derived all data from the prior trials. We emphasize that SEP-1 is consistent with the conclusions of these trials and does not require an invasive method of patient reassessment. In regards to Dr. Kalil's publication in Critical Care Medicine, "Early Goal-Directed Therapy for Sepsis: A Novel Solution for Discordant Survival Outcomes in Clinical Trials," the major conclusion was that: "[S]urvival discordance was not associated with differences in early goal-directed therapy bundle compliance or hemodynamic goal achievement. Our results suggest that it was associated with faster and more appropriate antibiotic co-intervention in the early goal-directed therapy arm compared with controls in the observational studies but not in the randomized trials." While we may dispute the methods and analysis used to reach this conclusion, we again underscore that SEP-1 does not mandate an invasive reassessment but does require early IV antibiotic administration. Thus SEP-1 is consistent with the Kalil publication in its approach.

- Developer Response to Comment ID 6806: The developers will take the Armstrong Institute's helpful suggestion under advisement and model data to understand how a 2 hours standard would affect the performance characteristics of the measure. We agree that earlier administration of antibiotics is the preferred approach.
- Comments also noted the varying level of evidence for the different components in the measure composite: repeat lactate, fluid reassessment, and physical exam. The comment suggested that these components are not equivalent to antibiotic or IV fluid administration and should not be weighted equally in the composite construct. Another comment recommended a simplified three hour bundle without the repeat lactate and physical exam component.
  - Developer Response to Comment ID 6708: We will take under consideration the suggestion to weight elements in different ways than presently weighted. It is important to note however that as a matter of process for vetting measures, they must be advanced on the basis of accumulated data and evidence. SEP-1 continues to show a high association with reduction in mortality with the individual specified elements as documented in the submission which justified the weighting. As the submission shows, there is a large separation in mortality between those who comply with the elements in total and those who fail any one or more than one of the elements. To understand and model a proposal such as Dr. Doerfler's will require analysis of the measure as a component measure, which necessarily means analyzing the data in a fashion for which it was not designed. We will discuss with CMS Dr. Doerfler's hypothesis regarding preferential weighting of certain data elements.
  - Developer Response to Comment ID 6810: The developers appreciate the opportunity to respond to the High Value Healthcare Collaborative (HVHCC) which has advanced excellent work in quality improvement and care of patients with severe sepsis and septic shock. We note that the HVHCC has indicated that SEP-1 endorses an "all-or-none payment approach." SEP-1 performance does not impact payment to hospitals or providers. As regards concerns that SEP-1's composite construction does not differentiate between importance of antibiotic administration and a physical exam element (cited as skin color), the developers would like to point out that, in the current specification manual and in this NQF submission, SEP-1 does not require documentation of particular physical exam elements. A provider may now indicate simply (within the allotted timeframe) that they have "performed a physical exam" without regard to a means or method, physical exam or otherwise, and pass this data element. In this regard, the developers would suggest that regular reassessment of a patient with septic shock as regards to perfusion status is as important as antibiotic administration and supports the composite construction as advanced in this submission.

As regards the representation that "once a mature care model is in place, compliance with a 3-hr-bundle, had no impact on in-hospital, 30-day, 90-day or 1-year post discharge mortality between those receiving the full bundle vs not. This is consistent with the conclusions from the ProCESS, ARISE, and ProMISE trials," this claim is not an accurate representation of the cited trials. The 3 hour elements of care were required of every patient in the cited trials. All patients received the three



## 0500 Severe Sepsis and Septic Shock: Management Bundle

hour elements (initial lactate, blood culture collection, and broad spectrum antibiotic administration) prior to randomization. Aside from this inaccuracy, it is unclear what the characteristics of a “mature care model” may be in the HVHCC’s remarks, however generalizing that the 3000+ hospitals in the United States subject to SEP-1 have such a model is unsupported by any evidence to properly analyze that claim. Moreover, since HVHCC does endorse a “simplified 3-hour-bundle” it would seem to remain an important element of care. The developers do not believe that clinicians and hospitals may not define “innovative approaches to early sepsis detection” under SEP-1. One method to ascertain time zero that overrides all other methods is a provider’s documentation of the time. In that regard, the HVHCC may use whatever method they prefer to set a time zero as long as their clinicians concur with the HVHCC’s approach. The developers appreciate the HVHCC’s suggestion to proceed with the 3 hour elements in SEP-1 and we assure them that these elements remain in this submission. As regards the representation that there is “no evidence” supporting repeat lactate assessment or a physical exam, the developers repeat that clinicians must only document reassessment of perfusion or volume status by any means of their choosing. This practice, along with repeat lactate assessment do have a supporting evidence base in the 2016 Surviving Sepsis Campaign guidelines. Reassessment is a best practice statement under the GRADE evaluation framework and repeat lactate assessment has a low quality of evidence with a weak recommendation under the same criteria. We will take under advisement that these elements should be examined further in future iterations of the specifications and would welcome the opportunity to work with the HVHCC to model these approaches. We note that the measure does not apply to critical access facilities and is not active at this time in any pay for performance programs.

- Two other comments questioned the percentage agreement rates among the patient level data elements. The commenters also disagreed with the guidance given to the standing committee on evaluation composite measures.
  - Developer Response to Comment ID 6803-6804: The Federation of American Hospitals has raised many questions that were previously discussed in detail in committee. We appreciate the opportunity to summarize these issues. While burden of data collection may be greater than for other measures in healthcare, this is more than counterbalanced by severe sepsis and septic shock’s burden on the healthcare system as the number one cause of inpatient deaths in the United States and highest cost condition for hospital admissions. Evidence that the SEP-1 measure drives quality improvement was provided at NQF. The initial three quarters of data analysis show that hospitals improved their performance from quarter to quarter. In addition, the analysis revealed a statistically significant finding that there is an approximately 8.5% associated reduction in mortality in those who comply with the measure versus those who fail the measure. As measure developers we have no data to comment on the quality of responses provided by Quality Net, but we will share the feedback with the Quality Net team. We cannot address the Federation’s representations regarding the motives of other agencies to utilize the measure, but we have provided substantial data that SEP-1 meets the standards set out in NQF’s measure evaluation framework. As regards validity, the measure met standards for assessing validity at the performance score level, which is the proper level of evaluation for a composite measure. Additionally, it is precisely for the Federations’ argument of the limited sample size (303 cases) that the data element level validity cannot serve as a valid critique of SEP-1. In addition, the measure met all reliability criteria with statistically appropriate analysis using a signal to noise methodology. While the Federation states that the element level validity testing is more important than the performance measure score testing, under the NQF measure evaluation framework, that choice is an improper standard to evaluate a composite measure. We note the Federation misinterprets the Technical Expert Panel’s remarks that “the individual components may not be sufficiently reliable independently, but could contribute to the reliability of the composite performance measure.” First, this quotation refers to reliability whereas the Federation was addressing validity. Secondly, the principle that the individual elements could contribute to the overall

## 0500 Severe Sepsis and Septic Shock: Management Bundle

validity at the level of the performance score is precisely the point of the Technical Expert Panel's comment. This rationale is why element validity is not the criterion for composite measures. Finally, the Federation has advanced no evidence to evaluate the claim that the measure does not meet the validity criteria set out by NQF at the performance measure score level. The Federation has inadvertently misstated the evaluation criteria. Specifically, subcriterion 2d states that "[f]or composite performance measures, empirical analyses support the composite construction approach..." Nothing in the Federation's remarks indicates that the composite construction approach is under question. As regards "missing data," the measure evaluation frame work considers this under subcriterion 2b7, which requires that the measure "analyses and identifies the extent and distribution of missing data (or non-response) and demonstrate that performance results are not biased due to systematic missing data (or differences between responders and nonresponders and how the specified handling of missing data minimizes biases." This requirement is different than the scenario described by the Federation which equates the variation in element level validity testing with missing data. This equivalency is incorrect insofar as the analysis of the complete data shows that some degree of variation actually exists – if the data were "missing" a showing of variation could not be made. In this regard, the developers stand by their submission that all data is present and cannot be missing as part of the reporting requirements met by 99.9% of over 3200 acute care hospitals in three consecutive quarters of analyzed data. To the extent that SEP-1 data elements may be contained in an electronic health record, the developers will take the Federation's excellent suggestion under advisement to consider, if feasible, the measure as an e-measure. The Federation implies that hospitals do not understand why they fail the measure given its composite nature. The developers believe, however, that by analyzing the point of failure in the measure framework, providers know exactly where weak spots are in their clinical care processes. For example, failure of the antibiotic element indicates a process in antibiotic administration that needs attention. In addition, the software provided by major vendors to hospitals to report the measure specifically categorizes the level of the fallout for facilities. The developers strenuously object to the characterization that the measure has limited value in improving patient care: in over 600,000 patients captured by the measure there has been an approximately 8.5% reduction in mortality in those compliant with the measure versus those who were not. For the population of submitted cases, this represents a potential lives saved calculation of over 7,500 patients in the first three quarters of data for the measure.

- Developer Response to Comment ID 6811-6813: The developer's appreciate the opportunity to respond to the remarks of the American Medical Association (AMA) and clarify the operation of SEP-1. As regards to Dr. Pronovost's publication in the American Journal of Medical Quality regarding possible unintended consequences of quality measures, we note that Dr. Pronovost is the Director of the Armstrong Institute for Patient Safety and Quality which has recommended that SEP-1 be re-endorsed with the exception of tightening the antibiotic administration requirement from 3 hours to 2 hours. Please see the submitted comment of Dr. Matt Austin, PHD, on behalf of the Armstrong Institute for Patient Safety and Quality at Johns Hopkins University, which was also received during the post-evaluation comment period. With respect to the specific concern that fluid administration as specified in SEP-1 may be harmful to patients with left ventricular systolic dysfunction, the AMA cites an opinion article authored by an emergency medicine resident. We note that this opinion is not representative of any clinical trial, observational or randomized, controlled or not. The opinion cites another article by Boyd 2011 which indicates that a positive fluid balance and elevated CVP are associated with increased mortality. The developers note that SEP-1 does not advocate for a "positive fluid balance" which refers to volume status over several days of care. SEP-1 is limited to initial resuscitation and the first 6 hours of care after presentation. In addition, SEP-1 does not advocate for CVP measurement, and certainly not an "elevated CVP." Another citation in the resident's article is Pudilo 2012 which reports frequency of myocardial

## 0500 Severe Sepsis and Septic Shock: Management Bundle

dysfunction in severe sepsis and septic shock. The article by Pudilo actually points to a reason for proper volume resuscitation of patients: the well-known presence of severe sepsis induced myocardial dysfunction. The salient finding is not that the patients have intrinsic heart disease, but rather that sepsis has caused impaired myocardial function. In addition, Pudilo does not conclude that a 30 ml/kg initial fluid bolus as an initial resuscitation strategy in septic shock is detrimental to patients. This Pudilo reference does not support any of AMA's criticism of the SEP-1 measure. On the broader concern about the potential risks of fluid resuscitation, we note that there is no published evidence from any randomized controlled trial which indicates that patients with severe sepsis and known congestive heart failure or renal failure who receive a fluid bolus for initial resuscitation do worse than other sepsis patients in terms of mortality. In fact, even the examination of the large trials on septic shock do not support this contention (EGDT 2001, Process 2014, Promise 2014, Arise 2015). In fact the only published evidence on the topic concludes that for patients with intermediate lactate values of 2-4 mmol/L who receive the full fluid bolus of 30 ml/kg with congestive heart failure and renal failure have lower mortality than their counterparts without these co-morbidities. (See: Lui V et al. Fluid Volume, Lactate Values, and Mortality in Sepsis Patients with Intermediate Lactate Values. Ann Am Thorac Soc Vol 10, No 5, pp 466-473, Oct 2013). The thrust of the AMA's comments on this topic regarding LVSD is that physician judgment should be preserved. The developers agree with the AMA that physician judgment is paramount and agree that providers should exercise their best judgment informed by the evidence when caring for sepsis patients. SEP-1 is not a prescriptive recipe for all patients with severe sepsis and septic shock, but rather a measurement strategy for processes of sepsis care. The developers fully expect that practitioners will do what is best in their understanding for each patient, which may result in deviation from SEP-1. Ultimately when sufficient data is amassed, best compliance, which takes into account those necessary deviations, will be known. In that regard, there is no expectation or goal of a 100% compliance with SEP-1. However, we emphasize that all current evidence suggests better mortality with higher compliance. As regards to the concern that a precisely specified measure should account for how unplanned drug shortages could impact an individual hospital's performance and the concern that mortality varied with a shortage of nor-epinephrine, we note that SEP-1 does not require the use of any one particular vasopressor. We also note that SEP-1 is a process measure, not an outcome measure such as mortality. Finally, we note that it is likely that all hospitals would be affected by any shortage in a short period of time. Turning to Dr. Kalil's publication in Critical Care Medicine, "Early Goal-Directed Therapy for Sepsis: A Novel Solution for Discordant Survival Outcomes in Clinical Trials," the major conclusion was that "[s]urvival discordance was not associated with differences in early goal-directed therapy bundle compliance or hemodynamic goal achievement. Our results suggest that it was associated with faster and more appropriate antibiotic co-intervention in the early goal-directed therapy arm compared with controls in the observational studies but not in the randomized trials." While we may dispute the methods and analysis used to reach this conclusion, we again underscore that SEP-1 does not mandate Early Goal Directed Therapy and SEP-1 does require early antibiotic administration. Thus SEP-1 is consistent with the Kalil publication in its approach. The PRISM investigators have reported results in the New England Journal of Medicine (NEJM) entirely consistent with the Process, Promise and Arise trials, also published in NEJM. Little information is imparted by PRISM that was not clearly known from the three other trials, and in fact PRISM derived all data from the prior trials. In any case, SEP-1 as specified is consistent with the conclusions of these trials. AMA has stated concerns with SEP-1's measure performance characteristics. For its validity, the measure met standards for assessing validity at the performance score level, which is the proper level of evaluation for a composite measure. In addition, the measure met reliability criteria with statistically appropriate analysis using a signal to noise methodology. While AMA represents that element level validity testing is more important than the performance measure score testing, this is an

## 0500 Severe Sepsis and Septic Shock: Management Bundle

improper standard to evaluate a composite measure under the NQF measure evaluation framework. Of substantial importance, we note that AMA misinterprets the Technical Expert Panel's remarks that "the individual components may not be sufficiently reliable independently, but could contribute to the reliability of the composite performance measure." First, the quoted principle that the individual elements could contribute to the overall validity at the level of the performance score is precisely the point of the Technical Expert Panel's recommendation not to focus on element level testing for a composite metric. Second, the AMA has advanced no evidence to evaluate the claim that the measure does not meet the validity criteria set out by NQF at the performance measure score level. In the evaluation of a metric, it would be improper to move the goal post mid-evaluation. Regarding their other comments on validity, the AMA incorrectly cites subcriterion 2d. Subcriterion 2d states that "[f]or composite performance measures, empirical analyses support the composite construction approach..." Nothing in the AMA's remarks indicates that the composite construction approach is under question. In regard to "missing data," the measure evaluation frame work actually considers this under subcriterion 2b7, which requires that the measure "analyzes and identifies the extent and distribution of missing data (or non-response) and demonstrate that performance results are not biased due to systematic missing data (or differences between responders and non-responders and how the specified handling of missing data minimizes biases." This requirement is different than the scenario described by AMA which equates the variation in element level validity testing with missing data. This equivalency is incorrect insofar as the analysis of the complete data shows that some degree of variation actually exists – if the data were "missing" a showing of variation could not be made. In this regard, the developers stand by their submission that all data is present and cannot be missing as part of the reporting requirements met by 99.9% of 3225 acute care hospitals in three consecutive quarters of analyzed data. In summary, the developers appreciate the AMA's comments and suggestions. We agree with the AMA that this disease which places an unacceptable burden in terms of deaths and expenditures on the healthcare system deserves a rigorous and robust measure. For this reason we are proud that SEP-1 (which has over 600,000 reported cases since inception) has an associated 8.5% reduction in mortality for those cases that were compliant with the measure versus those which were not. We look forward to tracking the ongoing impact of the SEP-1 measure on sepsis quality improvement and will share our findings with all concerned stakeholders as they become available.

- During the post comment call, the Committee re-iterated the fact that lactate clearance is an important step in evaluating the patient's condition. The Committee also noted that the intent of reassessment is to determine how well the patient is doing and is a best practice in that it drives collaboration among the care team. In regard to antibiotic administration, the Committee recommended that the developer update the rationale in the measure submission form to include that the intent of the measure is to encourage early administration of antibiotics, preferably within one hour diagnosis of sepsis. The developer committed to testing antibiotic administration within one hour of diagnosis of sepsis. NQF also provided additional guidance to the Committee on the scientific acceptability requirements for a composite measure. The Committee then agreed that the measure still met reliability and validity requirements.

**7. Consensus Standards Approval Committee (CSAC) Vote: Y-X; N-X**

**8. Board of Directors Vote: Y-X; N-X**

**9. Appeals**



## Measures Withdrawn from Consideration

Seven measures previously endorsed by NQF have not been re-submitted for maintenance of endorsement. Endorsement for these measures will be removed.

Measure	Reason for withdrawal
0393: Hepatitis C: Confirmation of Hepatitis C Viremia	In order to align the change in endorsement status with CMS's transition from the Physician Quality Reporting System (PQRS) to the Quality Payment Program (QPP).
0395: Paired Measure: Hepatitis C Ribonucleic Acid (RNA) Testing Before Initiating Treatment (paired with 0396)	In order to align the change in endorsement status with CMS's transition from the Physician Quality Reporting System (PQRS) to the Quality Payment Program (QPP).
0396: Paired Measure: Hepatitis C Virus (HCV) Genotype Testing Prior to Treatment (paired with 0395)	In order to align the change in endorsement status with CMS's transition from the Physician Quality Reporting System (PQRS) to the Quality Payment Program (QPP).
0398: Hepatitis C: Hepatitis C Virus (HCV) Ribonucleic Acid (RNA) Testing Between 4-12 Weeks after Initiation of Treatment	In order to align the change in endorsement status with CMS's transition from the Physician Quality Reporting System (PQRS) to the Quality Payment Program (QPP).
0399: Paired Measure: Hepatitis C: Hepatitis A Vaccination (paired with 0400)	In order to align the change in endorsement status with CMS's transition from the Physician Quality Reporting System (PQRS) to the Quality Payment Program (QPP).
0404: HIV/AIDS: CD4 Cell Count or Percentage Performed	Developer is no longer able to support measure. Retired by Developer.
0408: HIV/AIDS: Tuberculosis (TB) Screening	Developer is no longer able to support measure. Retired by Developer.

## **Appendix B: NQF Infectious Disease Portfolio and Related Measures**

*\*Denotes the measure is within the Infectious Disease portfolio*

### ***HIV/AIDS***

2079: HIV Medical Visit Frequency\*

2080: Gap in HIV Medical Visits\*

2083: Prescription of HIV Antiretroviral Therapy\*

0405: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis\*

2082: HIV Viral Load Suppression\*

3086: Population Level HIV Viral Load Suppression

### ***Sexually Transmitted Infections***

0409: Sexually Transmitted Diseases – Screening for Chlamydia, Gonorrhea, and Syphilis\*

0033: Chlamydia Screening in Women

### ***Hepatitis***

3059: One-Time Screening for Hepatitis C Virus (HCV) for Patients at Risk – trial use

3060: Annual Hepatitis C Virus (HCV) Screening for Patients who are Active Injection Drug Users – trial use

3061: Appropriate Screening Follow-up for Patients Identified with Hepatitis C Virus (HCV) Infection – trial use

### ***Vaccinations – Hepatitis B***

0475: Hepatitis B Vaccine Coverage Among All Live Newborn Infants Prior to Hospital or Birthing Facility Discharge

### ***Vaccinations – Childhood Immunizations***

0038: Childhood Immunization Status (CIS)

1407: Immunizations for Adolescents

### ***Vaccinations – HPV***

1959: Human Papillomavirus Vaccine for Female Adolescents (HPV)

### ***Vaccinations – Influenza***

0039: Flu Vaccinations for Adults Ages 18 and Older

0041: Preventive Care and Screening: Influenza Immunization

0431: Influenza Vaccination Coverage Among Healthcare Personnel

0680: Percent of Residents or Patients Who Were Assessed and Appropriately Given the Seasonal Influenza Vaccine (short stay)

0681: Percent of Residents Assessed and Appropriately Given the Seasonal Influenza Vaccine (long stay)

1659: Influenza Immunization

0226: Influenza Immunization in the ESRD Population (Facility Level)

### ***Vaccinations – Pneumococcal***

0043: Pneumococcal Vaccination Status for Older Adults (PNU)

### ***Sepsis***

0500: Severe Sepsis and Septic Shock: Management Bundle\*

3215: Adult Inpatient Risk Adjusted Sepsis Mortality\*

### ***Antimicrobial Use***

2720: National Healthcare Safety Network (NHSN) Antimicrobial Use Measure

### ***ENT***

0654: Acute Otitis Externa: Systemic Antimicrobial Therapy – Avoidance of Inappropriate Use (ENT)

### ***Respiratory***

0058: Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (AAB)\*

0069: Appropriate Treatment for Children with Upper Respiratory Infection (URI)\*

0147: Initial Antibiotic Selection for Community-Acquired Pneumonia (CAP) in Immunocompetent Patients

0096: Community-Acquired Bacterial Pneumonia (CAP): Empiric Antibiotic

2414: Pediatric Lower Respiratory Infection Readmission Measure

### ***Facility-Acquired Infections***

1716: National Healthcare Safety Network (NHSN) Facility-wide Inpatient Hospital-onset Methicillin-resistant Staphylococcus aureus (MRSA) Bacteremia Outcome Measure

1717: National Healthcare Safety Network (NHSN) Facility-wide Inpatient Hospital-onset Clostridium difficile Infection (CDI) Outcome Measure

0138: National Healthcare Safety Network (NHSN) Catheter-associated Urinary Tract Infection (CAUTI) Outcome Measure

0139: National Healthcare Safety Network (NHSN) Central line-associated Bloodstream Infection (CLABSI) Outcome Measure

0684: Percent of Residents with a Urinary Tract Infection (Long-Stay)

3025: Ambulatory Breast Procedure Surgical Site Infection (SSI) Outcome Measure



### ***Perioperative Antibiotics – Selection***

0268: Perioperative Care: Selection of Prophylactic Antibiotic: First OR Second Generation Cephalosporin

0528: Prophylactic Antibiotic Selection for Surgical Patients

0126: Selection of Antibiotic Prophylaxis for Cardiac Surgery Patients

### ***Perioperative Antibiotics – Timing***

0269: Timing of Prophylactic Antibiotics – Administering Physician

0527: Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision

### ***Perioperative Antibiotics – Discontinuation***

0271: Perioperative Care: Discontinuation of Prophylactic Parenteral Antibiotics (Non-Cardiac Procedures)

0529: Prophylactic Antibiotics Discontinued Within 24 Hours After Surgery End Time

0128: Duration of Antibiotic Prophylaxis for Cardiac Surgery Patients

### ***Perinatal***

1746: Intrapartum Antibiotic Prophylaxis for Group B Streptococcus (GBS)

0472: Appropriate Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision – Cesarean Section

0304: Late sepsis or meningitis in Very Low Birth Weight (VLBW) neonates (risk-adjusted)

## Appendix C: Infectious Disease Portfolio—Use in Federal Programs

NQF #	Title	Federal Programs: Finalized as of 2016 - 2017
0500	Severe Sepsis and Septic Shock: Management Bundle	Centers for Medicare & Medicaid Services Hospital Inpatient Quality Reporting Program
2079	HIV Medical Visit Frequency	Centers for Medicare & Medicaid Services Physician Quality Reporting System
2082	HIV Viral Load Suppression	Centers for Medicare & Medicaid Services Physician Quality Reporting System, Medicaid Adult Core Set
2083	Prescription of HIV Antiretroviral Therapy	Centers for Medicare & Medicaid Services Physician Quality Reporting System

## Appendix D: Project Standing Committee and NQF Staff

### STANDING COMMITTEE

**Woody Eisenberg, MD (Co-Chair)**

Senior Vice President of Performance Measurement, PQA  
Springfield, Virginia

**Adam Thompson, B.A (Co-Chair)**

Regional Partner Director, Northeast Caribbean AIDS Education and Training Centers  
Voorhees, New Jersey

**Emily Aaronson, MD**

Fellow, Patient Safety and Quality Improvement, Massachusetts General Hospital  
Boston, Massachusetts

**Amesh Adalja, MD**

Senior Associate, University of Pittsburgh Medical Center for Health Security  
Pittsburgh, Pennsylvania

**Esther Babady Otete, PhD, D (ABMM)**

Director of Microbiology Service Clinical Operations, Memorial Sloan Kettering Cancer Center  
New York, New York

**Nanette Benbow, MA**

Research Assistant Professor, Northwestern University  
Chicago, Illinois

**Kathleen Brady, MD, MSCE**

Medical Director and Medical Epidemiologist, Philadelphia Department of Public Health  
Philadelphia, Pennsylvania

**Laura Evans, MD, MSc**

Medical Director of Critical Care, Bellevue Hospital Center, New York University School of Medicine,  
New York, New York

**Piero Garzaro, MD**

Chair of Chiefs of Infectious Diseases, Kaiser Permanente  
Stockton, California

**Donald Goldman, MD**

Chief Medical and Scientific Officer, Clinical Professor of Pediatrics, Institute for Healthcare  
Improvement  
Cambridge, District of Columbia

**Jeffrey Hart, MS**

Kaiser Permanente

Rockville, Maryland

**Michael Lane, MD, MSc, MPHS, CPPS**

Assistant Professor of Medicine, Washington University School of Medicine  
St Louis, Missouri

**Jeffrey Lewis, BA**

Medical Case Manager, El Rio Community Health Center  
Tucson, Arizona

**Melinda Neuhauser, PharmD, MPH, FCCP, FASHP**

National PBM Clinical Pharmacy Program Manager in Infectious Diseases, Department of Veterans  
Affairs, Hines, Illinois

**Rocco Orlando, MD, FACS**

Senior Vice President and Chief Medical Officer, Hartford Healthcare  
Hartford, Connecticut

**Jamie Roney, DNP, RN-BC, BSHCM, CCRN-K**

Registered Nurse  
Lubbock, Texas

**Pranavi Sreeramoju, MD, MPH, CMQ, FSHEA, FIDSA**

Chief of Infection Prevention, Parkland Health & Hospital System  
Dallas, Texas

**NQF STAFF**

**Helen Burstin, MD, MPH**

Chief Scientific Officer

**Marcia Wilson, PhD, MBA**

Senior Vice President

**Elisa Munthali, MPH**

Vice President

**Melissa Mariñelarena, RN, MPA**

Senior Director

**Christy Skipper, MS**

Project Manager

**Mauricio Menendez, MS**

Project Analyst

## Appendix E: Measure Specifications

	2079 HIV Medical visit frequency
Steward	Health Resources and Services Administration - HIV/AIDS Bureau
Description	Percentage of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between medical visits. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.
Type	Process
Data Source	Paper Records
Level	Facility
Setting	Clinic/Clinician office
Numerator Statement	Number of patients in the denominator who had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period. (Measurement period is a consecutive 24-month period of time.)
Numerator Details	To be included in the numerator, patients must have had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period.
Denominator Statement	Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the first 6 months of the 24-month measurement period.
Denominator Details	To be included in the denominator, patients must meet all of the following conditions/events: <ol style="list-style-type: none"> <li>1. Patients of any age during the measurement period</li> <li>2. Patients without a date of death during the 24-month measurement period</li> <li>3. Patients diagnosed with HIV during the first 3 months of the 24-month measurement period or prior to the measurement period</li> <li>4. Patients who had at least one medical visit in the first 6 months of the 24-month measurement period</li> </ol>
Exclusions	Patients who died at any time during the 24-month measurement period.
Exclusion details	This measure has one exclusions – patient death during the measurement period. Due to constraints, the developer was not able to test the impact of the exclusion on this measure. It is important to note that patient mortality has reduced dramatically over the years primarily in relation to the development and dissemination of HIV antiretroviral therapy. Thus, we do not anticipate a significant number of patients that would be excluded from the measure.
Risk Adjustment	No risk adjustment
Stratification	No stratification
Type Score	Rate/proportion, Better quality = Higher score
Algorithm	1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator: 1.) diagnosed with HIV during the first 3 months of the 24-month measurement period or prior to the 24-month measurement period; 2.) did not have a date of death during the 24-month measurement period; and 3.) had at least one medical visit in the first 6 months of the 24-month measurement period. The individuals who met these three criteria are the denominator population.

	<b>2079 HIV Medical visit frequency</b>
	<p>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: must have had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period.</p> <p>3. Calculate the rate by dividing the numerator population by the denominator population and multiply by 100.</p>
Copyright / Disclaimer	Not applicable

	<b>2080 Gaps in HIV medical visits</b>
Steward	Health Resources and Services Administration-HIV/AIDS Bureau
Description	Percentage of patients, regardless of age, with a diagnosis of HIV who did not have a medical visit in the last 6 months of the measurement year
Type	Process
Data Source	Other, Paper Records
Level	Clinician : Group/Practice, Facility
Setting	Clinician Office/Clinic
Numerator Statement	Number of patients in the denominator who did not have a medical visit in the last 6 months of the measurement year (Measurement year is a consecutive 12-month period of time).
Numerator Details	To be included in the numerator, patients must not have had a medical visit in the last 6 months of the measurement year.
Denominator Statement	Number of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in the first 6 months of the measurement year. (The measurement year can be any consecutive 12-month period.)
Denominator Details	<p>To be included in the denominator, patients must meet all of the following conditions/events:</p> <ol style="list-style-type: none"> <li>1. Patients of any age during the measurement year</li> <li>2. Patients without a date of death during the measurement year</li> <li>3. Patients diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year</li> <li>4. Patients who had at least one medical visit in the first 6 months of the measurement year</li> </ol>
Exclusions	Patients who died at any time during the measurement year.
Exclusion details	Patients with a date of death during the measurement year.
Risk Adjustment	No risk adjustment
Stratification	No risk stratification
Type Score	Rate/Proportion, Better quality = Lower score
Algorithm	<p>1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator:</p> <p>1.) had a HIV diagnosis prior to the measurement year or during the first three months of the measurement year; 2.) did not have a date of death during the measurement year; and</p>

	<b>2080 Gaps in HIV medical visits</b>
	<p>3.) had at least one medical visit in the first 6 months of the measurement year. The individuals who met these three criteria are the denominator population.</p> <p>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: did not have a medical visit in the last 6 months of the measurement year.</p> <p>3. Calculate the percentage by dividing the numerator population by the denominator population and multiply by 100.</p>
Copyright / Disclaimer	Not applicable

	<b>3209 HIV Medical visit frequency</b>
Steward	Health Resources and Services Administration - HIV/AIDS Bureau
Description	Percentage of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in each 6-month period within 24 months with a minimum of 60 days between medical visits. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.
Type	Process
Data Source	Electronic Health Record
Level	Facility
Setting	Clinician Office/Clinic
Numerator Statement	Patients who had at least one medical visit in each 6-month of a consecutive consecutive 24 month period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period.
Numerator Details	<p>HIV medical visits are represented by a QDM variable that is comprised of the below seven different encounter type QDM elements:</p> <ul style="list-style-type: none"> <li>• Encounter, Performed: Face-to-Face Interaction using Face-to-Face Interaction Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1048)</li> <li>• Encounter, Performed: Office Visit using Office Visit Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1001)</li> <li>• Encounter, Performed: Outpatient Consultation using Outpatient Consultation Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1008)</li> <li>• Encounter, Performed: Preventive Care - Established Office Visit, 0 to 17 using Preventive Care - Established Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1024)</li> <li>• Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up using Preventive Care Services - Established Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1025)</li> <li>• Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up using Preventive Care Services-Initial Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1023)</li> <li>• Encounter, Performed: Preventive Care- Initial Office Visit, 0 to 17 using Preventive Care- Initial Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1022)</li> </ul>

	<b>3209 HIV Medical visit frequency</b>
Denominator Statement	Patients, regardless of age, diagnosed with HIV during the first 3 months of the year preceding the measurement period or prior to the measurement period with at least one medical visit in the first 6 months of the year preceding the measurement period.
Denominator Details	<p>The patient's HIV diagnosis is represented by the QDM element "Diagnosis: HIV" using "HIV Grouping Value Set (2.16.840.1.113883.3.464.1003.120.12.1003)". The patient's medical visits are represented by the following QDM elements:</p> <ul style="list-style-type: none"> <li>• Encounter, Performed: Face-to-Face Interaction using Face-to-Face Interaction Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1048)</li> <li>• Encounter, Performed: Office Visit using Office Visit Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1001)</li> <li>• Encounter, Performed: Outpatient Consultation using Outpatient Consultation Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1008)</li> <li>• Encounter, Performed: Preventive Care - Established Office Visit, 0 to 17 using Preventive Care - Established Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1024)</li> <li>• Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up using Preventive Care Services - Established Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1025)</li> <li>• Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up using Preventive Care Services-Initial Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1023)</li> <li>• Encounter, Performed: Preventive Care- Initial Office Visit, 0 to 17 using Preventive Care- Initial Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1022)</li> </ul> <p>The target population is identified by selecting patients based on their diagnosis with HIV.</p>
Exclusions	Patients who died at any time during the measurement period or the 12 months preceding the measurement period.
Exclusion details	This measure has one exclusion – patient death during the measurement period. The developer reports that the exclusion was tested similarly to other criteria using synthetic patients in Bonnie. When the exclusion element was present, the patients were correctly excluded from the measure. In the absence of the exclusion element, cases were not excluded from the measure.
Risk Adjustment	No risk adjustment
Stratification	No risk stratification
Type Score	Rate/proportion
Algorithm	<p>1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator: 1.) diagnosed with HIV during the first 3 months of the 24-month measurement period or prior to the 24-month measurement period; 2.) did not have a date of death during the 24-month measurement period; and 3.) had at least one medical visit in the first 6 months of the 24-month measurement period. The individuals who met these three criteria are the denominator population.</p> <p>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: must have had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period.</p>



	<b>3209 HIV Medical visit frequency</b>
	3. Calculate the rate by dividing the numerator population by the denominator population and multiply by 100.
Copyright / Disclaimer	Not applicable

	<b>2082 HIV Viral load suppression</b>
Steward	Health Resources and Services Administration - HIV/AIDS Bureau
Description	<p>Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year</p> <p>A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.</p>
Type	Outcome
Data Source	Laboratory, Paper records
Level	Facility
Setting	Clinician Office/Clinic
Numerator Statement	Number of patients in the denominator with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year
Numerator Details	To be included in the numerator, patients had a HIV viral load less than 200 copies/mL at the last HIV viral load test during the measurement year
Denominator Statement	Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the measurement year
Denominator Details	<p>To be included in the denominator, patients must meet all of the following conditions/events:</p> <ol style="list-style-type: none"> <li>1. Patients of any age during the measurement year</li> <li>2. Patients diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year</li> <li>3. Patients who had at least one medical visit during the measurement year</li> </ol>
Exclusions	No patient exclusions
Exclusion details	N/A
Risk Adjustment	No risk adjustment
Stratification	No risk stratification
Type Score	Rate/proportion, Better quality = Higher score
Algorithm	<ol style="list-style-type: none"> <li>1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator: 1.) diagnosed with a HIV during the first 3 months of the measurement year or prior to the measurement year; and 2.) had at least one medical visit during the measurement year. The individuals who met these criteria are the denominator population.</li> <li>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: had a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year.</li> <li>3. Calculate the percentage by dividing the numerator population by the denominator population and multiply by 100.</li> </ol>
Copyright / Disclaimer	N/A

	<b>3210 HIV Viral load suppression</b>
Steward	Health Resources and Services Administration - HIV/AIDS Bureau
Description	Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year.
Type	Outcome
Data Source	Electronic Health Record, Other
Level	Facility
Setting	Clinician Office/Clinic
Numerator Statement	Patients with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year. The outcome being measured is HIV viral suppression.
Numerator Details	The viral load suppression laboratory test is represented by the QDM element "Laboratory Test, Performed: HIV Viral Load" using "HIV Viral Load Grouping Value Set (2.16.840.1.113883.3.464.1003.120.12.1002)". The result of the laboratory test is modeled as an attribute of the Viral Load Suppression QDM element and represented as a numerical result associated with copies/mL as the reporting unit.
Denominator Statement	Patients, regardless of age, diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year who had at least one medical visit in the measurement year. The target population for this measure is all people living with HIV.
Denominator Details	<p>The patient's HIV diagnosis is represented by the QDM element "Diagnosis: HIV" using "HIV Grouping Value Set (2.16.840.1.113883.3.464.1003.120.12.1003)". The patient's medical visits are represented by the following QDM elements:</p> <ul style="list-style-type: none"> <li>• "Encounter, Performed: Face-to-Face Interaction" using "Face-to-Face Interaction Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1048)"</li> <li>• "Encounter, Performed: Office Visit" using "Office Visit Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1001)"</li> <li>• "Encounter, Performed: Outpatient Consultation" using "Outpatient Consultation Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1008)"</li> <li>• "Encounter, Performed: Preventive Care - Established Office Visit, 0 to 17" using "Preventive Care - Established Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1024)"</li> <li>• "Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up" using "Preventive Care Services - Established Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1025)"</li> <li>• "Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up" using "Preventive Care Services-Initial Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1023)"</li> <li>• "Encounter, Performed: Preventive Care- Initial Office Visit, 0 to 17" using "Preventive Care- Initial Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1022)"</li> </ul>
Exclusions	No patient exclusions
Exclusion details	Not applicable
Risk Adjustment	No risk adjustment
Stratification	No risk stratification
Type Score	Rate/proportion, Better quality = Higher score
Algorithm	1. Identify patients who meet the initial population criteria as defined by eCQM logic;

	<b>3210 HIV Viral load suppression</b>
	<p>2. Identify and count subset of the initial population that meet denominator criteria as defined by eCQM logic;</p> <p>3. Identify and count subset of patients in the denominator that meet numerator criteria as defined by eCQM logic.</p> <p>4. Calculate the performance measure rate: by dividing the number of patients in the numerator population by the number of patients in the denominator population.</p> <p>Note: the eCQM logic criteria for each population is defined in a computable format in the eCQM specifications provided as an attachment to this submission.</p>
Copyright / Disclaimer	Not applicable

	<b>2083 Prescription of HIV Antiretroviral Therapy</b>
Steward	Health Resources and Services Administration - HIV/AIDS Bureau
Description	Percentage of patients, regardless of age, with a diagnosis of HIV prescribed antiretroviral therapy for the treatment of HIV infection during the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.
Type	Process
Data Source	Paper records, Pharmacy, other
Level	Facility
Setting	Clinician Office/Clinic
Numerator Statement	Number of patients from the denominator prescribed HIV antiretroviral therapy during the measurement year.
Numerator Details	To be included in the numerator, patients were prescribed HIV antiretroviral therapy during the measurement year. HIV antiretroviral therapy at least one HIV antiretroviral medication.
Denominator Statement	Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the measurement year
Denominator Details	To be included in the denominator, patients must meet all of the following conditions/events: <ol style="list-style-type: none"> <li>1. Patients of any age during the measurement year</li> <li>2. Patients diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year</li> <li>3. Patients who had at least one medical visit during the measurement year</li> </ol>
Exclusions	No patient exclusions
Exclusion details	Not applicable
Risk Adjustment	No risk adjustment
Stratification	No risk stratification
Type Score	Rate/proportion, Better quality = Higher score
Algorithm	<ol style="list-style-type: none"> <li>1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator: 1.) diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year; and 2.) had at least one medical visit during the measurement year. The individuals who met these criteria are the denominator population.</li> <li>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: prescribed HIV antiretroviral therapy during the measurement year.</li> <li>3. Calculate the percentage by dividing the numerator population by the denominator population and multiply by 100.</li> </ol>
Copyright / Disclaimer	Not applicable

	<b>3211 Prescription of HIV Antiretroviral Therapy</b>
Steward	Health Resources and Services Administration - HIV/AIDS Bureau
Description	Percentage of patients, regardless of age, with a diagnosis of HIV prescribed antiretroviral therapy for the treatment of HIV infection during the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.
Type	Process
Data Source	Electronic Health Record
Level	Facility
Setting	Clinician Office/Clinic
Numerator Statement	Number of patients from the denominator prescribed HIV antiretroviral therapy during the measurement year.
Numerator Details	The antiretroviral therapy medication order is represented by the QDM element "Medication, Order: FDA Approved HIV Antiretroviral Therapy" using "HIV Antiretroviral Therapy RXNORM Value Set (2.16.840.1.113762.1.4.1032.1)." In order to be included in the numerator, the "Medication, Order: FDA Approved HIV Antiretroviral Therapy" element must start during the measurement period.
Denominator Statement	Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the measurement year
Denominator Details	<p>The patient's HIV diagnosis is represented by the QDM element "Diagnosis: HIV" using "HIV Grouping Value Set (2.16.840.1.113883.3.464.1003.120.12.1003)".</p> <p>The patient's medical visits are represented by the following QDM elements:</p> <ul style="list-style-type: none"> <li>• "Diagnosis: HIV 1" using "HIV 1 Grouping Value Set (2.16.840.1.113883.3.464.1003.120.12.1004)"</li> <li>• "Encounter, Performed: Face-to-Face Interaction" using "Face-to-Face Interaction Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1048)"</li> <li>• "Encounter, Performed: Office Visit" using "Office Visit Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1001)"</li> <li>• "Encounter, Performed: Outpatient Consultation" using "Outpatient Consultation Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1008)"</li> <li>• "Encounter, Performed: Preventive Care - Established Office Visit, 0 to 17" using "Preventive Care - Established Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1024)"</li> <li>• "Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up" using "Preventive Care Services - Established Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1025)"</li> <li>• "Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up" using "Preventive Care Services-Initial Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1023)"</li> <li>• "Encounter, Performed: Preventive Care- Initial Office Visit, 0 to 17" using "Preventive Care- Initial Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1022)"</li> </ul>
Exclusions	No patient exclusions
Exclusion details	Not applicable
Risk Adjustment	No risk adjustment
Stratification	No risk stratification

	<b>3211 Prescription of HIV Antiretroviral Therapy</b>
Type Score	Rate/proportion, Better quality = Higher score
Algorithm	<p>1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator: 1.) diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year; and 2.) had at least one medical visit during the measurement year. The individuals who met these criteria are the denominator population.</p> <p>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: prescribed HIV antiretroviral therapy during the measurement year.</p> <p>3. Calculate the percentage by dividing the numerator population by the denominator population and multiply by 100.</p>
Copyright / Disclaimer	Not applicable

	<b>0500 Severe Sepsis and Septic Shock Management Bundle</b>
Steward	Henry Ford Hospital
Description	This measure focuses on adults 18 years and older with a diagnosis of severe sepsis or septic shock. Consistent with Surviving Sepsis Campaign guidelines, the measure contains several elements, including measurement of lactate, obtaining blood cultures, administering broad spectrum antibiotics, fluid resuscitation, vasopressor administration, reassessment of volume status and tissue perfusion, and repeat lactate measurement. As reflected in the data elements and their definitions, these elements should be performed in the early management of severe sepsis and septic shock.
Type	Composite
Data Source	Imaging-Diagnostic, Laboratory, Other, Paper Records, Pharmacy
Level	Facility
Setting	Hospital
Numerator Statement	<ul style="list-style-type: none"> <li>The number of patients in the denominator who received ALL of the following components (if applicable) for the early management of severe sepsis and septic shock: initial lactate levels, blood cultures, antibiotics, fluid resuscitation, repeat lactate level, vasopressors, and volume status and tissue perfusion reassessment.</li> </ul>
Numerator Details	<p>In addition to the previous information (above) about assessing the numerator population, the following also are part of the numerator details.</p> <ul style="list-style-type: none"> <li>Within 3 hours of presentation of severe sepsis: <ul style="list-style-type: none"> <li>Measure initial lactate level</li> <li>Draw blood cultures prior to antibiotics</li> <li>Administer broad spectrum or other antibiotics</li> </ul> </li> <li>Within 6 hours of presentation of severe sepsis: <ul style="list-style-type: none"> <li>Repeat lactate level (if initial lactate &gt; 2 mmol/L)</li> </ul> </li> <li>Within 3 hours of presentation of septic shock: <ul style="list-style-type: none"> <li>Administer 30 ml/kg crystalloid for hypotension or lactate <math>\geq</math> 4 mmol/L</li> </ul> </li> <li>Within 6 hours of presentation of septic shock: <ul style="list-style-type: none"> <li>Apply vasopressors (for hypotension that does not respond to initial fluid resuscitation) to maintain a mean arterial pressure (MAP) <math>\geq</math> 65 mm Hg</li> <li>Reassess volume status and tissue perfusion in the event of persistent hypotension (MAP &lt;65 mm Hg) after initial fluid administration or initial lactate level <math>\geq</math> 4 mmol/L <ul style="list-style-type: none"> <li>The clinician is no longer required to state the method of reassessment used (e.g. physical exam, bedside cardiovascular ultrasound, passive leg raising, CVP, ScvO2 assessment). The clinician can attest that volume and perfusion reassessment has occurred, even without reference to the method used. This will meet the measure's volume and perfusion reassessment requirement. A provider may also opt to state their chosen method, but this is not required.</li> </ul> </li> </ul> </li> </ul> <p>The full definitions of each of these data elements and additional information are included in the data dictionary, attached to the submission</p>
Denominator Statement	Inpatients age 18 and over with an ICD-10-CM Principal or Other Diagnosis Code of Sepsis, Severe Sepsis, or Septic Shock.



	0500 Severe Sepsis and Septic Shock Management Bundle
Denominator Details	<p>Discharges age 18 and over with an ICD-10-CM Principal or Other Diagnosis Code of Sepsis, Severe Sepsis, or Septic Shock as defined in the table below:</p> <p>ICD-10-CM Code   Code Description</p> <p>A021    Salmonella sepsis</p> <p>A227    Anthrax sepsis</p> <p>A267    Erysipelothrix sepsis</p> <p>A327    Listerial sepsis</p> <p>A400    Sepsis due to streptococcus, group A</p> <p>A401    Sepsis due to streptococcus, group B</p> <p>A403    Sepsis due to Streptococcus pneumoniae</p> <p>A408    Other streptococcal sepsis</p> <p>A409    Streptococcal sepsis, unspecified</p> <p>A4101   Sepsis due to Methicillin susceptible Staphylococcus aureus</p> <p>A4102   Sepsis due to Methicillin resistant Staphylococcus aureus</p> <p>A411    Sepsis due to other specified staphylococcus</p> <p>A412    Sepsis due to unspecified staphylococcus</p> <p>A413    Sepsis due to Hemophilus influenzae</p> <p>A414    Sepsis due to anaerobes</p> <p>A4150   Gram-negative sepsis, unspecified</p> <p>A4151   Sepsis due to Escherichia coli [E. coli]</p> <p>A4152   Sepsis due to Pseudomonas</p> <p>A4153   Sepsis due to Serratia</p> <p>A4159   Other Gram-negative sepsis</p> <p>A4181   Sepsis due to Enterococcus</p> <p>A4189   Other specified sepsis</p> <p>A419    Sepsis, unspecified organism</p> <p>A427    Actinomycotic sepsis</p> <p>A5486   Gonococcal sepsis</p> <p>B377    Candidal sepsis</p> <p>R6520   Severe sepsis without septic shock</p> <p>R6521   Severe sepsis with septic shock</p> <p>Data elements required to calculate the denominator (in alphabetical order):</p> <ul style="list-style-type: none"> <li>• Administrative Contraindication to Care, Septic Shock</li> <li>• Administrative Contraindication to Care, Severe Sepsis</li> <li>• Admission Date</li> <li>• Birthdate</li> <li>• Directive for Comfort Care or Palliative Care, Septic Shock</li> <li>• Directive for Comfort Care or Palliative Care, Severe Sepsis</li> <li>• Discharge Date</li> <li>• Discharge Disposition</li> <li>• Discharge Time</li> <li>• Transfer From Another Hospital or ASC</li> </ul>

	<b>0500 Severe Sepsis and Septic Shock Management Bundle</b>
	The full definitions of each of these data elements and additional information are included in the data dictionary, attached to the submission
Exclusions	<p>The following patients are excluded from the denominator:</p> <ul style="list-style-type: none"> <li>• Severe sepsis is not present</li> <li>• Patients Transferred in from another acute care facility</li> <li>• Patients receiving IV antibiotics for more than 24 hours prior to presentation of severe sepsis.</li> <li>• Patients with a Directive for Comfort Care or Palliative Care within 3 hours of presentation of severe sepsis</li> <li>• Patients with an Administrative Contraindication to Care within 6 hours of presentation of severe sepsis</li> <li>• Patients with an Administrative Contraindication to Care within 6 hours of presentation of septic shock</li> <li>• Patients with a Directive for Comfort Care or Palliative Care within 6 hours of presentation of septic shock</li> <li>• Patients with septic shock who are discharged within 6 hours of presentation</li> <li>• Patients with severe sepsis who are discharged within 6 hours of presentation</li> <li>• Patients with a Length of Stay &gt;120 days</li> <li>• Patients included in a Clinical Trial</li> </ul>
Exclusion details	<p>To determine the length of stay, the admission date and discharge date are entered. If the result of the calculation subtracting the admission date from the discharge date is greater than 120 days, the patient is excluded from the measure.</p> <p>Data Elements required to determine denominator exclusions (in alphabetical order):</p> <p>Administrative Contraindication to Care, Septic Shock</p> <p>Administrative Contraindication to Care, Severe Sepsis</p> <p>Admission Date</p> <p>Birthdate</p> <p>Clinical Trial</p> <p>Directive for Comfort Care or Palliative Care, Septic Shock</p> <p>Directive for Comfort Care or Palliative Care, Severe Sepsis</p> <p>Discharge Date</p> <p>Discharge Disposition</p> <p>Discharge Time</p> <p>Transfer from Another Hospital or ASC</p> <p>The full definitions of each of these data elements and additional information are included in the data dictionary, attached to the submission.</p>
Risk Adjustment	No risk adjustment
Stratification	No risk stratification
Type Score	Rate/proportion, Better quality = Higher score
Algorithm	The detailed measure algorithm for SEP-1 is available in the data dictionary attached to the submission, along with a diagram.

	0500 Severe Sepsis and Septic Shock Management Bundle
	<ol style="list-style-type: none"> <li>1. Find the patients who meet the initial population (ie, the general group of patients that a set of performance measures is designed to address).</li> <li>2. From the patients within the initial population criteria, find the patients who qualify for the denominator (ie, the specific group of patients for inclusion in a specific performance measure based on defined criteria) Note: in some cases the initial population and denominator are identical. Remove any patients that meet the denominator exclusion criteria.</li> <li>3. The following actions are required within 3 hours of presentation of severe sepsis: <ol style="list-style-type: none"> <li>a. Measure initial lactate level</li> <li>b. Draw blood cultures prior to antibiotics</li> <li>c. Administer broad spectrum or other antibiotics</li> </ol> </li> <li>4. The following actions are required within 6 hours of presentation of severe sepsis: <ol style="list-style-type: none"> <li>a. Repeat lactate level (if initial lactate &gt; 2 mmol/L)</li> </ol> </li> <li>5. The following actions are required within 3 hours of presentation of septic shock: <ol style="list-style-type: none"> <li>a. Administer 30 ml/kg crystalloid for hypotension or lactate <math>\geq</math> 4 mmol/L</li> </ol> </li> <li>6. The following actions are required within 6 hours of presentation of septic shock: <ol style="list-style-type: none"> <li>a. Apply vasopressors (for hypotension that does not respond to initial fluid resuscitation) to maintain a mean arterial pressure (MAP) <math>\geq</math> 65 mm Hg</li> <li>b. Reassess volume status and tissue perfusion in the event of persistent hypotension (MAP &lt;65 mm Hg) after initial fluid administration or initial lactate level <math>\geq</math> 4 mmol/L</li> </ol> </li> </ol> <p><i>Note: The clinician is no longer required to state the method of reassessment used (e.g. physical exam, bedside cardiovascular ultrasound, passive leg raising, CVP, ScvO2 assessment). The clinician can attest that volume and perfusion reassessment has occurred, even without reference to the method used. This will meet the measure's volume and perfusion reassessment requirement. A provider may also opt to state their chosen method, but this is not required.</i></p> <ol style="list-style-type: none"> <li>7. All of the above numerator components (as applicable) must be in compliance, otherwise the case is calculated as a 'failed' sepsis case.</li> </ol>
Copyright / Disclaimer	Not applicable

	<b>3215 Adult Inpatient Risk Adjusted Sepsis Mortality</b>
Steward	New York State Department of Health, Office of Quality and Patient Safety
Description	Annual risk adjusted inpatient mortality rate for adult patients (aged 18 and over) admitted to acute care hospitals with diagnosis of severe sepsis or septic shock. The measure includes patients in acute care hospital settings over one year timeframe who had, either on admission, or during their hospital stay, a clinical diagnosis of severe sepsis (now referred to as 'sepsis') or septic shock using criteria described in the International Sepsis Definitions (Sepsis-2)
Type	Outcome
Data Source	Claims (Other), EHRs Hybrid, Laboratory, Management Data, Non-Medical Data, Paper Records, Pharmacy, Registry
Level	Facility
Setting	Hospital
Numerator Statement	Outcome is risk adjusted inpatient mortality rate for adult patients (18 and over) admitted to an acute care hospital with a diagnosis of severe sepsis or septic shock or who develop severe sepsis or septic shock during their hospital stay.
Numerator Details	Inpatient mortality is noted on data submission from hospital. Clinical variables needed for risk adjustment including demographics, co-morbidities, severity, and potential exclusions are reported by hospital as described in the data dictionary.
Denominator Statement	All adult patient discharges (18 and over) in a calendar year with a diagnosis of severe sepsis or septic shock on admission or at any time during their hospital stay. This may include multiple admissions of the same patient during the measurement year. Denominator includes all cases identified using any means (administrative, registry, electronic health records, billing data, etc.), either prospectively, retrospectively, or both, that meet the International consensus definition (Sepsis- 2) of severe sepsis or septic shock.
Denominator Details	All adult patients meeting International consensus definition (Sepsis-2) for Severe Sepsis/Septic shock identified through combination of any relevant hospital clinical and/or administrative databases, prospectively or retrospectively.
Exclusions	Patients with advanced directives in place prior to episode of sepsis which specifically restrict any hospital specific sepsis protocol interventions or who decline (or their proxy declines) treatment for sepsis. Patients who have been transferred from one acute care hospital to another are excluded.
Exclusion details	Patients who have any of the following characteristics, reported on data variables fully described in the data dictionary, are excluded from the calculation of risk adjusted mortality rates for a specific hospital:

	<b>3215 Adult Inpatient Risk Adjusted Sepsis Mortality</b>
	<ol style="list-style-type: none"> <li>1. Advanced Directives in place prior to diagnosis of severe sepsis or septic shock that specifically preclude active treatment according to that hospital's protocol for severe sepsis and septic shock.</li> <li>2. Patient or patient proxy refusal of treatment for severe sepsis or septic shock according to that hospital's protocol for severe sepsis and septic shock.</li> <li>3. Patients who were transferred between acute care hospitals.</li> </ol>
Risk Adjustment	Statistical risk model
Stratification	Not applicable
Type Score	Rate/proportion, Better quality= Lower score
Algorithm	<p>The study objective was to develop a logistic regression model to estimate the probability of hospital mortality among septic patients entering 179 New York State hospitals over the period of January 1, 2015 through December 31, 2015. The a priori analysis plan eliminated any patient with an advanced directive or who declined interventions. When a patient was discharged from a hospital as "transfer to acute care", only the patient's data from the receiving hospital was used in the dataset. If a patient was in the dataset multiple times for sepsis, only the final admission was used. This preserved the outcome of interest (mortality) and observation independence in the data file for developing logistic regression models. This resulted in a database total of 43,204 septic patients. The a priori analysis used only patient demographics, comorbidities, and admission characteristics to estimate the probability of hospital mortality. Specifically treatment variables were not used in the model.</p> <p><b>Septic patients</b></p> <p>All subjects entered into the model met the admitting hospital's criteria for severe sepsis or septic shock. Severe sepsis was defined as a suspected or confirmed infection, at least two systemic manifestations of infection and one or more acute organ dysfunctions. Septic shock was defined as severe sepsis where at least one organ dysfunction with sustained hypotension after a fluid challenge. For this paper, the term sepsis or septic represents the dataset population of severe sepsis and septic shock patients. Mortality is defined as in-hospitals deaths.</p> <p><b>Statistical Methods</b></p> <p>Logistic regression developed a model to estimate the probability of mortality for patients with severe sepsis or septic shock during their hospital stay. A list of the possible predictor variables and definitions are given in Table 1. Maximum likelihood was used to estimate model coefficients and associated standard errors. The hierarchical nature of the data supports random-effects logistic regression use since patients are nested within the 179</p>

	3215 Adult Inpatient Risk Adjusted Sepsis Mortality
	<p>hospitals. However, the 179 random-effect coefficients would have made the resulting model specific only to those 179 New York hospitals and would not be generalizable to patients outside these specific hospitals. A random sample of 10% (N = 4,319) of the observations were set aside and the logistic regression model was developed on the remaining 90% (38,884 observations). The final model was validated on the 10% of observations that were set aside. Patient comorbidities were generated using the list shown in supplemental Table S1. We generated a variable called mechanical ventilation (MV) severity that indicated a severity of illness relating to mechanical ventilation. This dichotomous variable was defined when a patient was admitted to the hospital already mechanically ventilated or requiring mechanical ventilation within 6 hours post admission. Initial serum lactate was not measured in 2,528 (5.9%) patients and was imputed using single imputation. Specifically, truncated linear regression was used during the imputation procedure where the lower limit of left truncation was set at a serum lactate level 0.1 mmol/L (1st percentile) and the upper limit of the right truncation was set at 30.0 mmol/L (99th percentile). A list of predictor variables is shown in supplemental Table S2.</p> <p>A multivariable logistic regression model was built using the developmental dataset and starting with all possible covariates in the model. Using an iterative procedure, variables were removed from the model, one by one, if their p-values were not significant at 0.05 level until a parsimonious model was reached. Variables removed during the development procedure were added back into the model if their p-values were significant at the 0.05 level and if model calibration (Hosmer-Lemeshow goodness of fit) was improved through their inclusion. We then assessed the scale of the 3 continuous variables (patient age, first serum lactate, and the count of the number of comorbidities) remaining in the model. Specifically, we were interested in determining whether these variables had a linear relationship with mortality. Using the method of fractional polynomials patient age was included in the model as a linear term, the number of comorbidities was transformed by taking the square root of the number of comorbidities, and first serum lactate was entered into the model as a quadratic expression (linear and a squared term). Model calibration was further improved by adding the following interactions to the model: lower respiratory infection (LRI) and MV severity, patient age and the square root of the number of comorbidities, and first serum lactate and the square root of the number of comorbidities.</p> <p>Model calibration was assessed using the Hosmer-Lemeshow goodness of fit on both the developmental and the validation datasets. Group sizes of 10, 100, 500, and 1,000 were chosen for the large, developmental, dataset while group sizes of 10, 50, 100, and 150 were chosen for the smaller validation dataset. Model discrimination was assessed using the area</p>

	<b>3215 Adult Inpatient Risk Adjusted Sepsis Mortality</b>
	<p>under the receiver operating characteristic (ROC) curve for both the developmental and validation datasets.</p> <p>The estimated probability of mortality was generated using the model coefficients and the specific patient attributes. If the patient attribute is defined by a categorical variable, then the possible values are either a 0 or 1. If the attribute is defined by a continuous variable, then the specific value is used such as the patient's age. Interaction values are generated by multiplying the values of each of the two individual variables defined by the interaction. The product of the coefficient and the patient's value for all of the variables in the model are generated. Next the logic is defined as the sum of the above products. Finally, the probability of mortality for a specific patient is generated using the follow equation:</p> <p>Probability of mortality= <math>\exp(\text{logit}) / (1 + \exp(\text{logit}))</math></p>
Copyright / Disclaimer	Not applicable

## Appendix F: Related and Competing Measures

### Comparison of NQF #2079, #2080, #2082, and #2083

	2079 HIV Medical Visit Frequency	2080 Gaps in Medical Visits	2082 HIV Viral Load Suppression	2083 Prescription of HIV Antiretroviral Therapy
Steward	Health Resources and Services Administration - HIV/AIDS Bureau	Health Resources and Services Administration-HIV/AIDS Bureau	Health Resources and Services Administration - HIV/AIDS Bureau	Health Resources and Services Administration - HIV/AIDS Bureau
Description	Percentage of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between medical visits. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.	Percentage of patients, regardless of age, with a diagnosis of HIV who did not have a medical visit in the last 6 months of the measurement year	Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.	Percentage of patients, regardless of age, with a diagnosis of HIV prescribed antiretroviral therapy for the treatment of HIV infection during the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.
Type	Process	Process	Outcome	Process
Data Source	Paper Records	Other, Paper Records	Laboratory, Paper records	Paper records, Pharmacy, other
Level	Facility	Clinician : Group/Practice, Facility	Facility	Facility
Setting	Clinic/Clinician office	Clinician Office/Clinic	Clinician Office/Clinic	Clinician Office/Clinic
Numerator Statement	Number of patients in the denominator who had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between first medical visit in the prior 6-month period and the last	Number of patients in the denominator who did not have a medical visit in the last 6 months of the measurement year (Measurement year is a consecutive 12-month period of time).	Number of patients in the denominator with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year	Number of patients from the denominator prescribed HIV antiretroviral therapy during the measurement year.



	2079 HIV Medical Visit Frequency	2080 Gaps in Medical Visits	2082 HIV Viral Load Suppression	2083 Prescription of HIV Antiretroviral Therapy
	medical visit in the subsequent 6-month period. (Measurement period is a consecutive 24-month period of time.)			
Numerator Details	To be included in the numerator, patients must have had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period.	To be included in the numerator, patients must not have had a medical visit in the last 6 months of the measurement year.	To be included in the numerator, patients had a HIV viral load less than 200 copies/mL at the last HIV viral load test during the measurement year	To be included in the numerator, patients were prescribed HIV antiretroviral therapy during the measurement year. HIV antiretroviral therapy at least one HIV antiretroviral medication.
Denominator Statement	Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the first 6 months of the 24-month measurement period.	Number of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in the first 6 months of the measurement year. (The measurement year can be any consecutive 12-month period.)	Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the measurement year	Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the measurement year
Denominator Details	To be included in the denominator, patients must meet all of the following conditions/events: 1. Patients of any age during the measurement period 2. Patients without a date of death during the 24-month measurement period 3. Patients diagnosed with HIV during the first 3 months of the 24-month	To be included in the denominator, patients must meet all of the following conditions/events: 1. Patients of any age during the measurement year 2. Patients without a date of death during the measurement year 3. Patients diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year	To be included in the denominator, patients must meet all of the following conditions/events: 1. Patients of any age during the measurement year 2. Patients diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year 3. Patients who had at least one medical visit during the measurement year	To be included in the denominator, patients must meet all of the following conditions/events: 1. Patients of any age during the measurement year 2. Patients diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year 3. Patients who had at least one medical visit during the measurement year

	2079 HIV Medical Visit Frequency	2080 Gaps in Medical Visits	2082 HIV Viral Load Suppression	2083 Prescription of HIV Antiretroviral Therapy
	measurement period or prior to the measurement period 4. Patients who had at least one medical visit in the first 6 months of the 24-month measurement period	4. Patients who had at least one medical visit in the first 6 months of the measurement year		
Exclusions	Patients who died at any time during the 24-month measurement period.	Patients who died at any time during the measurement year.	No patient exclusions	No patient exclusions
Exclusion Details	This measure has one exclusions – patient death during the measurement period. Due to constraints, the developer was not able to test the impact of the exclusion on this measure. It is important to note that patient mortality has reduced dramatically over the years primarily in relation to the development and dissemination of HIV antiretroviral therapy. Thus, we do not anticipate a significant number of patients that would be excluded from the measure.	Patients with a date of death during the measurement year.	Not Applicable	Not applicable
Risk Adjustment	No risk adjustment	No risk adjustment	No risk adjustment	No risk adjustment
Stratification	No stratification	No risk stratification	No risk stratification	No risk stratification
Type Score	Rate/proportion, Better quality = Higher score	Rate/Proportion, Better quality = Lower score	Rate/proportion, Better quality = Higher score	Rate/proportion, Better quality = Higher score
Algorithm	1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator:	1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator:	1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator: 1.)	1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator: 1.)

	2079 HIV Medical Visit Frequency	2080 Gaps in Medical Visits	2082 HIV Viral Load Suppression	2083 Prescription of HIV Antiretroviral Therapy
	<p>1.) diagnosed with HIV during the first 3 months of the 24-month measurement period or prior to the 24-month measurement period; 2.) did not have a date of death during the 24-month measurement period; and 3.) had at least one medical visit in the first 6 months of the 24-month measurement period. The individuals who met these three criteria are the denominator population.</p> <p>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: must have had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period.</p> <p>3. Calculate the rate by dividing the numerator population by the denominator population and multiply by 100.</p>	<p>1.) had a HIV diagnosis prior to the measurement year or during the first three months of the measurement year; 2.) did not have a date of death during the measurement year; and 3.) had at least one medical visit in the first 6 months of the measurement year. The individuals who met these three criteria are the denominator population.</p> <p>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: did not have a medical visit in the last 6 months of the measurement year.</p> <p>3. Calculate the percentage by dividing the numerator population by the denominator population and multiply by 100.</p>	<p>diagnosed with a HIV during the first 3 months of the measurement year or prior to the measurement year; and 2.) had at least one medical visit during the measurement year. The individuals who met these criteria are the denominator population.</p> <p>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: had a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year.</p> <p>3. Calculate the percentage by dividing the numerator population by the denominator population and multiply by 100.</p>	<p>diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year; and 2.) had at least one medical visit during the measurement year. The individuals who met these criteria are the denominator population.</p> <p>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: prescribed HIV antiretroviral therapy during the measurement year.</p> <p>3. Calculate the percentage by dividing the numerator population by the denominator population and multiply by 100.</p>
Submission items	5.1a. Identified measures 0403 : HIV/AIDS: Medical Visit	5.1a Identified Measures: 0405: HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis	5.1a Identified Measures:	5.1a Identified Measures:

	2079 HIV Medical Visit Frequency	2080 Gaps in Medical Visits	2082 HIV Viral Load Suppression	2083 Prescription of HIV Antiretroviral Therapy
	<p>0405: HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis</p> <p>0409: HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</p> <p>2080: Gap in HIV Medical Visits</p> <p>2082: HIV viral suppression</p> <p>2083: Prescription of HIV Antiretroviral Therapy</p> <p>3211: Prescription of HIV Antiretroviral Therapy</p> <p>3210: HIV viral suppression</p> <p>3010: HIV Medical Visit Frequency</p> <p>5a. Are the measure specifications harmonized to the extent possible?</p> <p>Yes</p> <p>5b. Competing Measures: This measure does not have a competing measure.</p>	<p>0409: HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</p> <p>2079: HIV Medical Visit Frequency</p> <p>2080: HIV Viral Suppression</p> <p>2083: Prescription of HIV Antiretroviral Therapy</p> <p>5a.1 Are specs completely harmonized? Yes</p> <p>5b. Competing Measures: This measure does not have a competing measure.</p>	<p>0407: HIV/AIDS: HIV RNA Control After Six Months of Potent Antiretroviral Therapy</p> <p>0405: HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis</p> <p>0409: HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</p> <p>2079: HIV Medical Visit Frequency</p> <p>2080: Gap in HIV Medical Visits</p> <p>2083: Prescription of HIV Antiretroviral Therapy</p> <p>3211: Prescription of HIV Antiretroviral Therapy</p> <p>3210: HIV Viral Suppression</p> <p>3010: HIV Medical Visit Frequency</p> <p>5a.1 Are specs completely harmonized? No</p> <p>5a.2. If the measure specifications are not completely harmonized, identify the differences, rationale, and impact on interpretability and data collection burden.</p> <p>Harmonized with all measures except 405 and 409. Plans to harmonize with 405 and 409.</p>	<p>0405: HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis</p> <p>0409: HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</p> <p>2079: HIV Medical Visit Frequency</p> <p>2080: Gap in HIV Medical Visits</p> <p>2082: HIV Viral Suppression</p> <p>3211: Prescription of HIV Antiretroviral Therapy</p> <p>3210: HIV viral suppression</p> <p>3010: HIV Medical Visit Frequency</p> <p>3211: Prescription of HIV Antiretroviral Therapy</p> <p>5a.1 Are specs completely harmonized? Yes</p> <p>5a.2. If the measure specifications are not completely harmonized, identify the differences, rationale, and impact on interpretability and data collection burden.</p> <p>Harmonized with all measures except 405 and 409. Plans to harmonize with 405 and 409.</p>

	3209 HIV Medical Visit Frequency eMeasure	3210 HIV Viral Load Suppression eMeasure	3211 Prescription of HIV Antiretroviral Therapy eMeasure
Steward	Health Resources and Services Administration - HIV/AIDS Bureau	Health Resources and Services Administration - HIV/AIDS Bureau	Health Resources and Services Administration - HIV/AIDS Bureau
Description	Percentage of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in each 6-month period within 24 months with a minimum of 60 days between medical visits. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.	Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year.	Percentage of patients, regardless of age, with a diagnosis of HIV prescribed antiretroviral therapy for the treatment of HIV infection during the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.
Type	Process	Outcome	Process
Data Source	Electronic Health Record	Electronic Health Record, Other	Electronic Health Record
Level	Facility	Facility	Facility
Setting	Clinician Office/Clinic	Clinician Office/Clinic	Clinician Office/Clinic
Numerator Statement	Patients who had at least one medical visit in each 6-month of a consecutive consecutive 24 month period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period.	Patients with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year. The outcome being measured is HIV viral suppression.	Number of patients from the denominator prescribed HIV antiretroviral therapy during the measurement year.
Numerator Details	HIV medical visits are represented by a QDM variable that is comprised of the below seven different encounter type QDM elements: <ul style="list-style-type: none"> <li>Encounter, Performed: Face-to-Face Interaction using Face-to-Face Interaction Grouping Value Set</li> </ul>	The viral load suppression laboratory test is represented by the QDM element "Laboratory Test, Performed: HIV Viral Load" using "HIV Viral Load Grouping Value Set (2.16.840.1.113883.3.464.1003.120.12.1002)". The result of the laboratory test is modeled as an attribute of the Viral Load Suppression QDM element and represented as a numerical result associated with copies/mL as the reporting unit.	The antiretroviral therapy medication order is represented by the QDM element "Medication, Order: FDA Approved HIV Antiretroviral Therapy" using "HIV Antiretroviral Therapy RXNORM Value Set (2.16.840.1.113762.1.4.1032.1)." In order to be included in the numerator, the "Medication, Order: FDA Approved HIV

	3209 HIV Medical Visit Frequency eMeasure	3210 HIV Viral Load Suppression eMeasure	3211 Prescription of HIV Antiretroviral Therapy eMeasure
	<p>(2.16.840.1.113883.3.464.1003.101.12.1 048)</p> <ul style="list-style-type: none"> <li>Encounter, Performed: Office Visit using Office Visit Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1 001)</li> <li>Encounter, Performed: Outpatient Consultation using Outpatient Consultation Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1 008)</li> <li>Encounter, Performed: Preventive Care - Established Office Visit, 0 to 17 using Preventive Care - Established Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1 024)</li> <li>Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up using Preventive Care Services - Established Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1 025)</li> <li>Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up using Preventive Care Services-Initial Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1 023)</li> <li>Encounter, Performed: Preventive Care- Initial Office Visit, 0 to 17 using Preventive Care- Initial Office</li> </ul>		<p>Antiretroviral Therapy” element must start during the measurement period.</p>

	3209 HIV Medical Visit Frequency eMeasure	3210 HIV Viral Load Suppression eMeasure	3211 Prescription of HIV Antiretroviral Therapy eMeasure
	Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1022)		
Denominator Statement	Patients, regardless of age, diagnosed with HIV during the first 3 months of the year preceding the measurement period or prior to the measurement period with at least one medical visit in the first 6 months of the year preceding the measurement period.	Patients, regardless of age, diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year who had at least one medical visit in the measurement year. The target population for this measure is all people living with HIV.	Number of patients, regardless of age, with a diagnosis of HIV with at least one medical visit in the measurement year
Denominator Details	<p>The patient's HIV diagnosis is represented by the QDM element "Diagnosis: HIV" using "HIV Grouping Value Set (2.16.840.1.113883.3.464.1003.120.12.1003)". The patient's medical visits are represented by the following QDM elements:</p> <ul style="list-style-type: none"> <li>Encounter, Performed: Face-to-Face Interaction using Face-to-Face Interaction Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1048)</li> <li>Encounter, Performed: Office Visit using Office Visit Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1001)</li> <li>Encounter, Performed: Outpatient Consultation using Outpatient Consultation Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1008)</li> <li>Encounter, Performed: Preventive Care - Established Office Visit, 0 to 17 using Preventive Care -</li> </ul>	<p>The patient's HIV diagnosis is represented by the QDM element "Diagnosis: HIV" using "HIV Grouping Value Set (2.16.840.1.113883.3.464.1003.120.12.1003)". The patient's medical visits are represented by the following QDM elements:</p> <ul style="list-style-type: none"> <li>"Encounter, Performed: Face-to-Face Interaction" using "Face-to-Face Interaction Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1048)"</li> <li>"Encounter, Performed: Office Visit" using "Office Visit Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1001)"</li> <li>"Encounter, Performed: Outpatient Consultation" using "Outpatient Consultation Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1008)"</li> <li>"Encounter, Performed: Preventive Care - Established Office Visit, 0 to 17" using "Preventive Care - Established Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1024)"</li> <li>"Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up" using "Preventive Care Services -</li> </ul>	<p>The patient's HIV diagnosis is represented by the QDM element "Diagnosis: HIV" using "HIV Grouping Value Set (2.16.840.1.113883.3.464.1003.120.12.1003)".</p> <p>The patient's medical visits are represented by the following QDM elements:</p> <ul style="list-style-type: none"> <li>"Diagnosis: HIV 1" using "HIV 1 Grouping Value Set (2.16.840.1.113883.3.464.1003.120.12.1004)"</li> <li>"Encounter, Performed: Face-to-Face Interaction" using "Face-to-Face Interaction Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1048)"</li> <li>"Encounter, Performed: Office Visit" using "Office Visit Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1001)"</li> <li>"Encounter, Performed: Outpatient Consultation" using "Outpatient Consultation Grouping Value Set</li> </ul>

	3209 HIV Medical Visit Frequency eMeasure	3210 HIV Viral Load Suppression eMeasure	3211 Prescription of HIV Antiretroviral Therapy eMeasure
	<p>Established Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1024)</p> <ul style="list-style-type: none"> <li>Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up using Preventive Care Services - Established Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1025)</li> <li>Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up using Preventive Care Services-Initial Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1023)</li> <li>Encounter, Performed: Preventive Care- Initial Office Visit, 0 to 17 using Preventive Care- Initial Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1022)</li> </ul> <p>The target population is identified by selecting patients based on their diagnosis with HIV.</p>	<p>Established Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1025)"</p> <ul style="list-style-type: none"> <li>"Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up" using "Preventive Care Services-Initial Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1023)"</li> <li>"Encounter, Performed: Preventive Care- Initial Office Visit, 0 to 17" using "Preventive Care- Initial Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1022)"</li> </ul>	<p>(2.16.840.1.113883.3.464.1003.101.12.1008)"</p> <ul style="list-style-type: none"> <li>"Encounter, Performed: Preventive Care - Established Office Visit, 0 to 17" using "Preventive Care - Established Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1024)"</li> <li>"Encounter, Performed: Preventive Care Services - Established Office Visit, 18 and Up" using "Preventive Care Services - Established Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1025)"</li> <li>"Encounter, Performed: Preventive Care Services-Initial Office Visit, 18 and Up" using "Preventive Care Services-Initial Office Visit, 18 and Up Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1023)"</li> <li>"Encounter, Performed: Preventive Care- Initial Office Visit, 0 to 17" using "Preventive Care- Initial Office Visit, 0 to 17 Grouping Value Set (2.16.840.1.113883.3.464.1003.101.12.1022)"</li> </ul>
Exclusions	Patients who died at any time during the measurement period or the 12 months preceding the measurement period.	No patient exclusions	No patient exclusions
Exclusion Details	This measure has one exclusion – patient death during the measurement period. The developer reports that the exclusion was tested similarly to other criteria	Not applicable	Not applicable



	3209 HIV Medical Visit Frequency eMeasure	3210 HIV Viral Load Suppression eMeasure	3211 Prescription of HIV Antiretroviral Therapy eMeasure
	using synthetic patients in Bonnie. When the exclusion element was present, the patients were correctly excluded from the measure. In the absence of the exclusion element, cases were not excluded from the measure.		
Risk Adjustment	No risk adjustment	No risk adjustment	No risk adjustment
Stratification	No risk stratification	No risk stratification	No risk stratification
Type Score	Rate/proportion, Better quality = Higher score	Rate/proportion, Better quality = Higher score	Rate/proportion, Better quality = Higher score
Algorithm	<p>1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator: 1.) diagnosed with HIV during the first 3 months of the 24-month measurement period or prior to the 24-month measurement period; 2.) did not have a date of death during the 24-month measurement period; and 3.) had at least one medical visit in the first 6 months of the 24-month measurement period. The individuals who met these three criteria are the denominator population.</p> <p>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: must have had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between first medical visit in the prior 6-month period and the last medical visit in the subsequent 6-month period.</p>	<p>1. Identify patients who meet the initial population criteria as defined by eCQM logic;</p> <p>2. Identify and count subset of the initial population that meet denominator criteria as defined by eCQM logic;</p> <p>3. Identify and count subset of patients in the denominator that meet numerator criteria as defined by eCQM logic.</p> <p>4. Calculate the performance measure rate: by dividing the number of patients in the numerator population by the number of patients in the denominator population.</p> <p>Note: the eCQM logic criteria for each population is defined in a computable format in the eCQM specifications provided as an attachment to this submission.</p>	<p>1. Identify the individuals who satisfy all specific criteria for inclusion in the denominator: 1.) diagnosed with HIV during the first 3 months of the measurement year or prior to the measurement year; and 2.) had at least one medical visit during the measurement year. The individuals who met these criteria are the denominator population.</p> <p>2. Identify the individuals from the denominator population who meet the criterion for inclusion in the numerator: prescribed HIV antiretroviral therapy during the measurement year.</p> <p>3. Calculate the percentage by dividing the numerator population by the denominator population and multiply by 100.</p>

	3209 HIV Medical Visit Frequency eMeasure	3210 HIV Viral Load Suppression eMeasure	3211 Prescription of HIV Antiretroviral Therapy eMeasure
	3. Calculate the rate by dividing the numerator population by the denominator population and multiply by 100.		
Submission items	<p>5.1a Identified Measures:</p> <p>0403: HIV/AIDS: Medical Visit</p> <p>0405: HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis</p> <p>0409: HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</p> <p>2080: Gap in HIV Medical Visits</p> <p>2082: HIV viral suppression</p> <p>2083: Prescription of HIV Antiretroviral Therapy</p> <p>3211: Prescription of HIV Antiretroviral Therapy</p> <p>3210: HIV viral suppression</p> <p>3010: HIV Medical Visit Frequency</p> <p>5a.1 Are specs completely harmonized? Yes</p> <p>5a.2. If the measure specifications are not completely harmonized, identify the differences, rationale, and impact on interpretability and data collection burden.</p> <p>Harmonized with all measures except 405 and 409. Plans to harmonize with 405 and 409.</p>	<p>5.1a Identified Measures:</p> <p>0405: HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis</p> <p>0409: HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</p> <p>2079: HIV Medical Visit Frequency</p> <p>2080: Gap in HIV Medical Visits</p> <p>2082: HIV Viral Suppression</p> <p>2083: Prescription of HIV Antiretroviral Therapy</p> <p>3211: Prescription of HIV Antiretroviral Therapy</p> <p>3010: HIV Medical Visit Frequency</p> <p>5a.1 Are specs completely harmonized? No</p> <p>5a.2. If the measure specifications are not completely harmonized, identify the differences, rationale, and impact on interpretability and data collection burden.</p> <p>Harmonization exists with all measures except 405 and 409. Plan to harmonize with 405 and 409.</p>	<p>5.1a Identified Measures:</p> <p>0405: HIV/AIDS: Pneumocystis Jiroveci Pneumonia (PCP) Prophylaxis</p> <p>0409: HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia, Gonorrhea, and Syphilis</p> <p>2079: HIV Medical Visit Frequency</p> <p>2080: Gap in HIV Medical Visits</p> <p>2082: HIV Viral Suppression</p> <p>3210: HIV Viral Suppression</p> <p>3010: HIV Medical Visit Frequency</p> <p>5a.1 Are specs completely harmonized? Yes</p> <p>5a.2. If the measure specifications are not completely harmonized, identify the differences, rationale, and impact on interpretability and data collection burden.</p> <p>This measure does not have a competing measure.</p>

	0500 Severe Sepsis and Septic Shock Management Bundle	3215 Adult Inpatient Risk Adjusted Sepsis Mortality
Steward	Henry Ford Hospital	New York State Department of Health, Office of Quality and Patient Safety
Description	This measure focuses on adults 18 years and older with a diagnosis of severe sepsis or septic shock. Consistent with Surviving Sepsis Campaign guidelines, the measure contains several elements, including measurement of lactate, obtaining blood cultures, administering broad spectrum antibiotics, fluid resuscitation, vasopressor administration, reassessment of volume status and tissue perfusion, and repeat lactate measurement. As reflected in the data elements and their definitions, these elements should be performed in the early management of severe sepsis and septic shock.	Annual risk adjusted inpatient mortality rate for adult patients (aged 18 and over) admitted to acute care hospitals with diagnosis of severe sepsis or septic shock. The measure includes patients in acute care hospital settings over one year timeframe who had, either on admission, or during their hospital stay, a clinical diagnosis of severe sepsis (now referred to as 'sepsis') or septic shock using criteria described in the International Sepsis Definitions (Sepsis-2)
Type	Composite	Outcome
Data Source	Imaging-Diagnostic, Laboratory, Other, Paper Records, Pharmacy	Claims (Other), EHRs Hybrid, Laboratory, Management Data, Non-Medical Data, Paper Records, Pharmacy, Registry
Level	Facility	Facility
Setting	Hospital	Hospital
Numerator Statement	The number of patients in the denominator who received ALL of the following components (if applicable) for the early management of severe sepsis and septic shock: initial lactate levels, blood cultures, antibiotics, fluid resuscitation, repeat lactate level, vasopressors, and volume status and tissue perfusion reassessment.	Outcome is risk adjusted inpatient mortality rate for adult patients (18 and over) admitted to an acute care hospital with a diagnosis of severe sepsis or septic shock or who develop severe sepsis or septic shock during their hospital stay.
Numerator Details	<p>In addition to the previous information (above) about assessing the numerator population, the following also are part of the numerator details.</p> <ul style="list-style-type: none"> <li>• Within 3 hours of presentation of severe sepsis: <ul style="list-style-type: none"> <li>o Measure initial lactate level</li> <li>o Draw blood cultures prior to antibiotics</li> <li>o Administer broad spectrum or other antibiotics</li> </ul> </li> <li>• Within 6 hours of presentation of severe sepsis:</li> </ul>	Inpatient mortality is noted on data submission from hospital. Clinical variables needed for risk adjustment including demographics, co-morbidities, severity, and potential exclusions are reported by hospital as described in the data dictionary.

	0500 Severe Sepsis and Septic Shock Management Bundle	3215 Adult Inpatient Risk Adjusted Sepsis Mortality
	<ul style="list-style-type: none"> <li>o Repeat lactate level (if initial lactate &gt; 2 mmol/L)</li> <li>• Within 3 hours of presentation of septic shock: <ul style="list-style-type: none"> <li>o Administer 30 ml/kg crystalloid for hypotension or lactate <math>\geq</math> 4 mmol/L</li> </ul> </li> <li>• Within 6 hours of presentation of septic shock: <ul style="list-style-type: none"> <li>o Apply vasopressors (for hypotension that does not respond to initial fluid resuscitation) to maintain a mean arterial pressure (MAP) <math>\geq</math> 65 mm Hg</li> <li>o Reasses volume status and tissue perfusion in the event of persistent hypotension (MAP &lt;65 mm Hg) after initial fluid administration or initial lactate level <math>\geq</math> 4 mmol/L <ul style="list-style-type: none"> <li>▪ The clinician is no longer required to state the method of reassessment used (e.g. physical exam, bedside cardiovascular ultrasound, passive leg raising, CVP, ScVO<sub>2</sub> assessment). The clinician can attest that volume and perfusion reassessment has occurred, even without reference to the method used. This will meet the measure's volume and perfusion reassessment requirement. A provider may also opt to state their chosen method, but this is not required.</li> </ul> </li> </ul> </li> </ul> <p>The full definitions of each of these data elements and additional information are included in the data dictionary, attached to the submission</p>	
Denominator Statement	Inpatients age 18 and over with an ICD-10-CM Principal or Other Diagnosis Code of Sepsis, Severe Sepsis, or Septic Shock.	All adult patient discharges (18 and over) in a calendar year with a diagnosis of severe sepsis or septic shock on admission or at any time during their hospital stay. This may include multiple admissions of the same patient during the measurement year. Denominator includes all cases identified using any means (administrative,

	0500 Severe Sepsis and Septic Shock Management Bundle	3215 Adult Inpatient Risk Adjusted Sepsis Mortality
		registry, electronic health records, billing data, etc.), either prospectively, retrospectively, or both, that meet the International consensus definition (Sepsis- 2) of severe sepsis or septic shock.
Denominator Details	<p>Discharges age 18 and over with an ICD-10-CM Principal or Other Diagnosis Code of Sepsis, Severe Sepsis, or Septic Shock as defined in the table below:</p> <p>ICD-10-CM Code Code Description</p> <p>A021 Salmonella sepsis</p> <p>A227 Anthrax sepsis</p> <p>A267 Erysipelothrix sepsis</p> <p>A327 Listerial sepsis</p> <p>A400 Sepsis due to streptococcus, group A</p> <p>A401 Sepsis due to streptococcus, group B</p> <p>A403 Sepsis due to Streptococcus pneumoniae</p> <p>A408 Other streptococcal sepsis</p> <p>A409 Streptococcal sepsis, unspecified</p> <p>A4101 Sepsis due to Methicillin susceptible Staphylococcus aureus</p> <p>A4102 Sepsis due to Methicillin resistant Staphylococcus aureus</p> <p>A411 Sepsis due to other specified staphylococcus</p> <p>A412 Sepsis due to unspecified staphylococcus</p> <p>A413 Sepsis due to Hemophilus influenzae</p> <p>A414 Sepsis due to anaerobes</p> <p>A4150 Gram-negative sepsis, unspecified</p> <p>A4151 Sepsis due to Escherichia coli [E. coli]</p> <p>A4152 Sepsis due to Pseudomonas</p> <p>A4153 Sepsis due to Serratia</p> <p>A4159 Other Gram-negative sepsis</p> <p>A4181 Sepsis due to Enterococcus</p> <p>A4189 Other specified sepsis</p>	All adult patients meeting International consensus definition (Sepsis-2) for Severe Sepsis/Septic shock identified through combination of any relevant hospital clinical and/or administrative databases, prospectively or retrospectively.

	0500 Severe Sepsis and Septic Shock Management Bundle	3215 Adult Inpatient Risk Adjusted Sepsis Mortality
	<p>A419 Sepsis, unspecified organism  A427 Actinomycotic sepsis  A5486 Gonococcal sepsis  B377 Candidal sepsis  R6520 Severe sepsis without septic shock  R6521 Severe sepsis with septic shock</p> <p>Data elements required to calculate the denominator (in alphabetical order):</p> <ul style="list-style-type: none"> <li>• Administrative Contraindication to Care, Septic Shock</li> <li>• Administrative Contraindication to Care, Severe Sepsis</li> <li>• Admission Date</li> <li>• Birthdate</li> <li>• Directive for Comfort Care or Palliative Care, Septic Shock</li> <li>• Directive for Comfort Care or Palliative Care, Severe Sepsis</li> <li>• Discharge Date</li> <li>• Discharge Disposition</li> <li>• Discharge Time</li> <li>• Transfer From Another Hospital or ASC</li> </ul> <p>The full definitions of each of these data elements and additional information are included in the data dictionary, attached to the submission</p>	
Exclusions	<p>The following patients are excluded from the denominator:</p> <ul style="list-style-type: none"> <li>• Severe sepsis is not present</li> <li>• Patients Transferred in from another acute care facility</li> <li>• Patients receiving IV antibiotics for more than 24 hours prior to presentation of severe sepsis.</li> </ul>	<p>Patients with advanced directives in place prior to episode of sepsis which specifically restrict any hospital specific sepsis protocol interventions or who decline (or their proxy declines) treatment for sepsis. Patients who have been transferred from one acute care hospital to another are excluded.</p>

	0500 Severe Sepsis and Septic Shock Management Bundle	3215 Adult Inpatient Risk Adjusted Sepsis Mortality
	<ul style="list-style-type: none"> <li>• Patients with a Directive for Comfort Care or Palliative Care within 3 hours of presentation of severe sepsis</li> <li>• Patients with an Administrative Contraindication to Care within 6 hours of presentation of severe sepsis</li> <li>• Patients with an Administrative Contraindication to Care within 6 hours of presentation of septic shock</li> <li>• Patients with a Directive for Comfort Care or Palliative Care within 6 hours of presentation of septic shock</li> <li>• Patients with septic shock who are discharged within 6 hours of presentation</li> <li>• Patients with severe sepsis who are discharged within 6 hours of presentation</li> <li>• Patients with a Length of Stay &gt;120 days</li> <li>• Patients included in a Clinical Trial</li> </ul>	
Exclusion Details	<p>To determine the length of stay, the admission date and discharge date are entered. If the result of the calculation subtracting the admission date from the discharge date is greater than 120 days, the patient is excluded from the measure.</p> <p>Data Elements required to determine denominator exclusions (in alphabetical order):</p> <p>Administrative Contraindication to Care, Septic Shock</p> <p>Administrative Contraindication to Care, Severe Sepsis</p> <p>Admission Date</p> <p>Birthdate</p> <p>Clinical Trial</p> <p>Directive for Comfort Care or Palliative Care, Septic Shock</p> <p>Directive for Comfort Care or Palliative Care, Severe Sepsis</p> <p>Discharge Date</p> <p>Discharge Disposition</p> <p>Discharge Time</p> <p>Transfer from Another Hospital or ASC</p>	<p>Patients who have any of the following characteristics, reported on data variables fully described in the data dictionary, are excluded from the calculation of risk adjusted mortality rates for a specific hospital:</p> <ol style="list-style-type: none"> <li>1. Advanced Directives in place prior to diagnosis of severe sepsis or septic shock that specifically preclude active treatment according to that hospital's protocol for severe sepsis and septic shock.</li> <li>2. Patient or patient proxy refusal of treatment for severe sepsis or septic shock according to that hospital's protocol for severe sepsis and septic shock.</li> <li>3. Patients who were transferred between acute care hospitals.</li> </ol>

	0500 Severe Sepsis and Septic Shock Management Bundle	3215 Adult Inpatient Risk Adjusted Sepsis Mortality
	The full definitions of each of these data elements and additional information are included in the data dictionary, attached to the submission.	
Risk Adjustment	No risk adjustment	Statistical risk model
Stratification	No risk stratification	Not applicable
Type Score	Rate/proportion, Better quality = Higher score	Rate/proportion, Better quality=Lower score
Algorithm	<p>The detailed measure algorithm for SEP-1 is available in the data dictionary attached to the submission, along with a diagram.</p> <ol style="list-style-type: none"> <li>Find the patients who meet the initial population (ie, the general group of patients that a set of performance measures is designed to address).</li> <li>From the patients within the initial population criteria, find the patients who qualify for the denominator (ie, the specific group of patients for inclusion in a specific performance measure based on defined criteria) Note: in some cases the initial population and denominator are identical. Remove any patients that meet the denominator exclusion criteria.</li> <li>The following actions are required within 3 hours of presentation of severe sepsis: <ol style="list-style-type: none"> <li>Measure initial lactate level</li> <li>Draw blood cultures prior to antibiotics</li> <li>Administer broad spectrum or other antibiotics</li> </ol> </li> <li>The following actions are required within 6 hours of presentation of severe sepsis: <ol style="list-style-type: none"> <li>Repeat lactate level (if initial lactate &gt; 2 mmol/L)</li> </ol> </li> <li>The following actions are required within 3 hours of presentation of septic shock: <ol style="list-style-type: none"> <li>Administer 30 ml/kg crystalloid for hypotension or lactate <math>\geq</math> 4 mmol/L</li> </ol> </li> </ol>	<p>The study objective was to develop a logistic regression model to estimate the probability of hospital mortality among septic patients entering 179 New York State hospitals over the period of January 1, 2015 through December 31, 2015. The a priori analysis plan eliminated any patient with an advanced directive or who declined interventions. When a patient was discharged from a hospital as “transfer to acute care”, only the patient’s data from the receiving hospital was used in the dataset. If a patient was in the dataset multiple times for sepsis, only the final admission was used. This preserved the outcome of interest (mortality) and observation independence in the data file for developing logistic regression models. This resulted in a database total of 43,204 septic patients. The a priori analysis used only patient demographics, comorbidities, and admission characteristics to estimate the probability of hospital mortality. Specifically treatment variables were not used in the model.</p> <p>Septic patients</p> <p>All subjects entered into the model met the admitting hospital’s criteria for severe sepsis or septic shock. Severe sepsis was defined as a suspected or confirmed infection, at least two systemic manifestations of infection and one or more acute organ dysfunctions. Septic shock was defined as severe sepsis where at least one organ dysfunction with sustained hypotension after a fluid challenge. For this paper, the term sepsis or septic represents the dataset population of severe sepsis and septic shock patients. Mortality is defined as in-hospitals deaths.</p> <p>Statistical Methods</p>



	0500 Severe Sepsis and Septic Shock Management Bundle	3215 Adult Inpatient Risk Adjusted Sepsis Mortality
	<p>6. The following actions are required within 6 hours of presentation of septic shock:</p> <p>a. Apply vasopressors (for hypotension that does not respond to initial fluid resuscitation) to maintain a mean arterial pressure (MAP) <math>\geq</math> 65 mm Hg</p> <p>b. Reassess volume status and tissue perfusion in the event of persistent hypotension (MAP &lt;65 mm Hg) after initial fluid administration or initial lactate level <math>\geq</math> 4 mmol/L</p> <p>Note: The clinician is no longer required to state the method of reassessment used (e.g. physical exam, bedside cardiovascular ultrasound, passive leg raising, CVP, ScVO<sub>2</sub> assessment). The clinician can attest that volume and perfusion reassessment has occurred, even without reference to the method used. This will meet the measure's volume and perfusion reassessment requirement. A provider may also opt to state their chosen method, but this is not required.</p> <p>7. All of the above numerator components (as applicable) must be in compliance, otherwise the case is calculated as a 'failed' sepsis case.</p>	<p>Logistic regression developed a model to estimate the probability of mortality for patients with severe sepsis or septic shock during their hospital stay. A list of the possible predictor variables and definitions are given in Table 1. Maximum likelihood was used to estimate model coefficients and associated standard errors. The hierarchical nature of the data supports random-effects logistic regression use since patients are nested within the 179 hospitals. However, the 179 random-effect coefficients would have made the resulting model specific only to those 179 New York hospitals and would not be generalizable to patients outside these specific hospitals. A random sample of 10% (N = 4,319) of the observations were set aside and the logistic regression model was developed on the remaining 90% (38,884 observations). The final model was validated on the 10% of observations that were set aside. Patient comorbidities were generated using the list shown in supplemental Table S1. We generated a variable called mechanical ventilation (MV) severity that indicated a severity of illness relating to mechanical ventilation. This dichotomous variable was defined when a patient was admitted to the hospital already mechanically ventilated or requiring mechanical ventilation within 6 hours post admission. Initial serum lactate was not measured in 2,528 (5.9%) patients and was imputed using single imputation. Specifically, truncated linear regression was used during the imputation procedure where the lower limit of left truncation was set at a serum lactate level 0.1 mmol/L (1st percentile) and the upper limit of the right truncation was set at 30.0 mmol/L (99th percentile). A list of predictor variables is shown in supplemental Table S2.</p> <p>A multivariable logistic regression model was built using the developmental dataset and starting with all possible covariates in the model. Using an iterative procedure, variables were removed from the model, one by one, if their p-values were not significant at 0.05 level until a parsimonious model was reached. Variables removed during the development procedure were added back into the model if their p-values were significant at the 0.05 level and if model calibration (Hosmer-Lemeshow goodness of fit) was improved through their inclusion. We then assessed the scale of the 3</p>

	0500 Severe Sepsis and Septic Shock Management Bundle	3215 Adult Inpatient Risk Adjusted Sepsis Mortality
		<p>continuous variables (patient age, first serum lactate, and the count of the number of comorbidities) remaining in the model. Specifically, we were interested in determining whether these variables had a linear relationship with mortality. Using the method of fractional polynomials patient age was included in the model as a linear term, the number of comorbidities was transformed by taking the square root of the number of comorbidities, and first serum lactate was entered into the model as a quadratic expression (linear and a squared term). Model calibration was further improved by adding the following interactions to the model: lower respiratory infection (LRI) and MV severity, patient age and the square root of the number of comorbidities, and first serum lactate and the square root of the number of comorbidities.</p> <p>Model calibration was assessed using the Hosmer-Lemeshow goodness of fit on both the developmental and the validation datasets. Group sizes of 10, 100, 500, and 1,000 were chosen for the large, developmental, dataset while group sizes of 10, 50, 100, and 150 were chosen for the smaller validation dataset. Model discrimination was assessed using the area under the receiver operating characteristic (ROC) curve for both the developmental and validation datasets.</p> <p>The estimated probability of mortality was generated using the model coefficients and the specific patient attributes. If the patient attribute is defined by a categorical variable, then the possible values are either a 0 or 1. If the attribute is defined by a continuous variable, then the specific value is used such as the patient's age. Interaction values are generated by multiplying the values of each of the two individual variables defined by the interaction. The product of the coefficient and the patient's value for all of the variables in the model are generated. Next the logit is defined as the sum of the above products. Finally, the probability of mortality for a specific patient is generated using the follow equation:  Probability of mortality= <math>\exp(\text{logit}) / (1 + \exp(\text{logit}))</math></p>
Submission items	5.1a Are there related measures or competing measures: No	5.1a Identified Measures: 0500: Severe Sepsis and Septic Shock: Management Bundle

	0500 Severe Sepsis and Septic Shock Management Bundle	3215 Adult Inpatient Risk Adjusted Sepsis Mortality
		5a.1 Are specs completely harmonized? Yes

## Appendix G: Pre-Evaluation Comments

Comments received as of February 23, 2017

Topic	Commenter	Comment
0500 Severe Sepsis and Septic Shock	Submitted by American Medical Association	The American Medical Association (AMA) appreciates the opportunity to comment prior to the NQF Infectious Disease Standing Committee's review. We strongly support the modifications that the developer made to the measure, #500, Severe Sepsis and Septic Shock: Management Bundle. Specifically, the additional requirement requiring documentation by a clinician to confirm severe sepsis addresses our concern that the previous denominator was too broadly defined and could have unintended negative consequences by including patients inappropriately. We also appreciate the additional enhancements made to the measure to capture individual patient circumstances, including capturing appropriate delays in obtaining blood cultures prior to antibiotic administration and the additional data elements to better identify those patients for whom crystalloid fluid administration is suitable. We would request one clarification in the measure specifications regarding one of the exclusions. Specifically, the denominator exclusions in S.8 include the following element, Patient included in clinical trial; yet, S.3.2, where changes to the measure are outlined, states that this exclusion has been removed. It would be helpful to have the developer clarify this discrepancy prior to releasing the measure for public and member comment. We look forward to monitoring the discussion of the measure and the opportunity to comment during the review and endorsement process.
0500 Severe Sepsis and Septic Shock	Submitted by Federation of American Hospitals	The Federation of American Hospitals ("FAH") appreciates the opportunity to comment prior to the NQF Infectious Disease Standing Committee's review. FAH believes that effective and timely treatment of severe sepsis and septic shock in patients is vital and we support the intent of the measure. FAH appreciates the modification to the measure to allow for administration of targeted antibiotics when the causative organism and susceptibility are known. FAH encourages the developer to continue to monitor the measure to ensure that this additional flexibility addresses the ongoing efforts of effective antibiotic stewardship and minimizing the potential for overuse. The FAH also asks for clarification of one of the

Topic	Commenter	Comment
		<p>measure specifications. Specifically, the denominator exclusions in S.8 include the following element, “Patient included in clinical trial”; yet, S.3.2, where changes to the measure are outlined, states that this exclusion has been removed. The FAH strongly encourages the developer to clarify this discrepancy prior to releasing the measure for public and member comment.</p> <p>The FAH remains extremely concerned that the sepsis measure continues to be overly complex and burdensome to collect, and, therefore, hampers the ability of hospitals to appropriately evaluate their performance on this measure since many of the data elements required for this measure can be captured only through chart abstraction. This complexity is evidenced by the validity testing results where forty data elements achieved less than 90% percent agreement between the data abstracted by hospitals and data abstracted by independent medical abstractors. FAH believes that the validity results demonstrate the inherent problems with implementing a measure with more than 140 data elements. In addition, even though hospitals collected the data for the measure since the last quarter of 2015 and the Centers for Medicare &amp; Medicaid Services (CMS) now has a full year of data, CMS has not yet publicly reported results due to concerns with data quality. The FAH asks that the committee consider these findings during the measure evaluation. While the FAH understands there are limitations to what can be collected in electronic health records currently, we strongly urge the developer to revise the measure to enable electronic data capture. Developing an eMeasure in this area would further ensure that the relevant information is available for use at the point of care and facilitate communication with providers at the next level of care. Continuing to maintain and endorse a measure that requires manual abstraction with this many elements should not be viewed as a long-term solution given the current environment of promoting electronic data capture.</p>
3209 HIV Medical Visit Frequency eMeasure	Submitted by New York State Department of Health AIDS Institute	Please review my comment on NQF#2079 related to the rigidity of frequency measures and their inability to apply to all people with HIV given established practice and clinical guidelines.
2079 HIV Medical Visit Frequency	Submitted by New York State	Representing the HIV Quality of Care Advisory Committee and Consumer Advisory Committee of the NY State

Topic	Commenter	Comment
	Department of Health AIDS Institute	Department of Health AIDS institute we would advise that this measure be dropped based on the variation of expected frequency of visits for patients based on their viral load suppression status. Frequency measures suggest that a rigid spacing of intervals of visits can be universally applied which is no longer the global standard of care - even in resource limited settings where differentiated care models are promoted by WHO and the Global Fund. The measure is not as useful at clinic level for improvement as missed visit measures, based on extensive research led by Mugavero among others.
2080 Gap in HIV Medical Visits	Submitted by New York State Department of Health AIDS Institute	Gaps in care should be focused on minimum standards applying to all patients as absence of a clinical visit within a 12 month period.
Comments on Standards not recommended	Submitted by New York State Department of Health AIDS Institute	In the proposed measures list, measures for CD4 monitoring and PCP prophylaxis still appear. The former is not a useful measure given the role of VL monitoring and suppression and the inability to see change easily over time. Although PCP prophylaxis is obviously of critical importance the number of eligible patients for intervention has diminished dramatically to make it of limited value as a quality measure.

National Quality Forum  
1030 15th St NW, Suite 800  
Washington, DC 20005  
<http://www.qualityforum.org>

ISBN [Assigned by Communications]  
©2015 National Quality Forum