



Infectious Disease 2016-2017: Maintenance Measures

Click the measure numbers to read more about the measure on QPS!

Measure Number	Title	Description	Measure Steward
0058	Avoidance of Antibiotic Treatment in Adults With Acute Bronchitis (AAB)	The percentage of adults 18–64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription.	National Committee for Quality Assurance
0069	Appropriate Treatment for Children With Upper Respiratory Infection (URI)	Percentage of children 3 months to 18 years of age with a diagnosis of upper respiratory infection (URI) who were not dispensed an antibiotic medication.	National Committee for Quality Assurance
0393	Hepatitis C: Confirmation of Hepatitis C Viremia	Percentage of patients aged 18 years and older who are hepatitis C antibody positive seen for an initial evaluation for whom hepatitis C virus (HCV) RNA testing was ordered or previously performed	American Gastroenterological Association
0395	Paired Measure: Hepatitis C Ribonucleic Acid (RNA) Testing Before Initiating Treatment (paired with 0396)	Percentage of patients aged 18 years and older with a diagnosis of chronic hepatitis C who started antiviral treatment within the 12 month reporting period for whom quantitative hepatitis C virus (HCV) ribonucleic acid (RNA) testing was performed within 12 months prior to initiation of antiviral treatment	American Gastroenterological Association
0396	Paired Measure: Hepatitis C Virus (HCV) Genotype Testing Prior to Treatment (paired with 0395)	Percentage of patients aged 18 years and older with a diagnosis of chronic hepatitis C who started antiviral treatment within the 12 month reporting period for whom hepatitis C virus (HCV) genotype testing was performed within 12 months prior to initiation of antiviral treatment	American Gastroenterological Association
0398	Hepatitis C: Hepatitis C Virus (HCV) Ribonucleic Acid (RNA) Testing Between 4-12 Weeks after Initiation of Treatment	Percentage of patients aged 18 years and older with a diagnosis of chronic hepatitis C who are receiving antiviral treatment for whom quantitative hepatitis C Virus (HCV) ribonucleic acid (RNA) testing was performed between 4-12 weeks after the initiation of antiviral treatment	American Gastroenterological Association

Measure Number	Title	Description	Measure Steward
0399	Paired Measure: Hepatitis C: Hepatitis A Vaccination (paired with 0400)	Percentage of patients aged 18 years and older with a diagnosis of hepatitis C who have received at least one injection of hepatitis A vaccine, or who have documented immunity to hepatitis A	American Gastroenterological Association
0404	HIV/AIDS: CD4 Cell Count or Percentage Performed	Percentage of patients aged six months and older with a diagnosis of HIV/AIDS, with at least two CD4 cell counts or percentages performed during the measurement year at least 3 months apart	National Committee for Quality Assurance
0405	HIV/AIDS: Pneumocystis jiroveci pneumonia (PCP) Prophylaxis	Percentage of patients aged 6 weeks or older with a diagnosis of HIV/AIDS, who were prescribed Pneumocystis jiroveci pneumonia (PCP) prophylaxis	National Committee for Quality Assurance
0408	HIV/AIDS: Tuberculosis (TB) Screening	Percentage of patients aged 3 months and older with a diagnosis of HIV/AIDS, for whom there was documentation that a tuberculosis (TB) screening test was performed and results interpreted (for tuberculin skin tests) at least once since the diagnosis of HIV infection.	National Committee for Quality Assurance
0409	HIV/AIDS: Sexually Transmitted Diseases – Screening for Chlamydia, Gonorrhea, and Syphilis	Percentage of patients aged 13 years and older with a diagnosis of HIV/AIDS, who have received chlamydia, gonorrhea, and syphilis screenings at least once since the diagnosis of HIV infection	National Committee for Quality Assurance
0500	Severe Sepsis and Septic Shock: Management Bundle	This measure will focus on patients aged 18 years and older who present with symptoms of severe sepsis or septic shock. These patients will be eligible for the 3 hour (severe sepsis) and/or 6 hour (septic shock) early management bundle.	Henry Ford Hospital
2079	HIV medical visit frequency	Percentage of patients, regardless of age, with a diagnosis of HIV who had at least one medical visit in each 6-month period of the 24-month measurement period with a minimum of 60 days between medical visits. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.	Health Resources and Services Administration - HIV/AIDS Bureau

Measure Number	Title	Description	Measure Steward
2080	Gap in HIV medical visits	Percentage of patients, regardless of age, with a diagnosis of HIV who did not have a medical visit in the last 6 months of the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.	Health Resources and Services Administration- HIV/AIDS Bureau
2082	HIV viral load suppression	Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.	Health Resources and Services Administration - HIV/AIDS Bureau
2083	Prescription of HIV Antiretroviral Therapy	Percentage of patients, regardless of age, with a diagnosis of HIV prescribed antiretroviral therapy for the treatment of HIV infection during the measurement year. A medical visit is any visit in an outpatient/ambulatory care setting with a nurse practitioner, physician, and/or a physician assistant who provides comprehensive HIV care.	Health Resources and Services Administration - HIV/AIDS Bureau