

NATIONAL QUALITY FORUM

National Voluntary Consensus Standards for Nursing Homes 2010

Measure Number/Title: NH-020-10: Percent of Residents Who Have/Had a Catheter Inserted and Left in Their Bladder (Long Stay)

Description: This measure updates CMS' current QM on catheter insertions. It is based on data from Minimum Data Set (MDS) 3.0 assessments of long-stay nursing home residents (those whose stay is longer than 100 days). This measure captures the percentage of long-stay residents who have had an indwelling catheter in the last 5 days noted on the most recent MDS 3.0 assessment, which may be annual, quarterly, significant change or significant correction during the selected quarter (3-month period). Long-stay residents are those residents who have been in nursing care at least 100 days. The measure is restricted to this population, which has long-term care needs, rather than the short stay population who are discharged within 100 days of admission.

Numerator Statement: The numerator statement refers to a catheter that was inserted and left in the bladder by the facility during the assessment period. During MDS 3.0 field testing, look-back periods were highlighted as a significant issue across the assessment tool. For clinical assessment items, longer look-back periods served to increase the amount of record review, increasing assessment burden and leading to more opportunities for error. During national testing of look-back periods for the MDS 3.0 proposed items, the 5-day look-back period performed well and likely contributed to the improved reliability of this item. The numerator is the number of long-stay residents who have/had a urinary catheter in the last 5 days (H0100A is checked).

Denominator Statement: The denominator is the total of all long-stay residents in the nursing home who have been assessed with an annual, quarterly, significant change or significant correction MDS 3.0 assessment during the quarter (3-month period) and who do not meet the exclusion criteria.

Level of Analysis: Facility/Agency

Data Source: Electronic clinical data

Measure developer: Research Triangle Institute International

Type of Endorsement (full or time-limited): Full

Attachments: Catheter Table

NATIONAL QUALITY FORUM

Measure Evaluation 4.1 December 2009

This form contains the measure information submitted by stewards. Blank fields indicate no information was provided. Attachments also may have been submitted and are provided to reviewers. The subcriteria and most of the footnotes from the evaluation criteria are provided in Word comments within the form and will appear if your cursor is over the highlighted area. Hyperlinks to the evaluation criteria and ratings are provided in each section.

TAP/Workgroup (if utilized): Complete all **yellow highlighted** areas of the form. Evaluate the extent to which each subcriterion is met. Based on your evaluation, summarize the strengths and weaknesses in each section.

Note: *If there is no TAP or workgroup, the SC also evaluates the subcriteria (yellow highlighted areas).*

Steering Committee: Complete all **pink** highlighted areas of the form. Review the workgroup/TAP assessment of the subcriteria, noting any areas of disagreement; then evaluate the extent to which each major criterion is met; and finally, indicate your recommendation for the endorsement. Provide the rationale for your ratings.

Evaluation ratings of the extent to which the criteria are met

- C = Completely (unquestionably demonstrated to meet the criterion)
- P = Partially (demonstrated to partially meet the criterion)
- M = Minimally (addressed BUT demonstrated to only minimally meet the criterion)
- N = Not at all (NOT addressed; OR incorrectly addressed; OR demonstrated to NOT meet the criterion)
- NA = Not applicable (only an option for a few subcriteria as indicated)

(for NQF staff use) NQF Review #: NH-020-10 NQF Project: Nursing Homes 2010
MEASURE DESCRIPTIVE INFORMATION
De.1 Measure Title: Percent of Residents Who Have/Had a Catheter Inserted and Left in Their Bladder (Long Stay)
De.2 Brief description of measure: This measure updates CMS' current QM on catheter insertions. It is based on data from Minimum Data Set (MDS) 3.0 assessments of long-stay nursing home residents (those whose stay is longer than 100 days). This measure captures the percentage of long-stay residents who have had an indwelling catheter in the last 5 days noted on the most recent MDS 3.0 assessment, which may be annual, quarterly, significant change or significant correction during the selected quarter (3-month period).
Long-stay residents are those residents who have been in nursing care at least 100 days. The measure is restricted to this population, which has long-term care needs, rather than the short stay population who are discharged within 100 days of admission.
1.1-2 Type of Measure: Process
De.3 If included in a composite or paired with another measure, please identify composite or paired measure The recommendation of the Steering Committee [to pair this measure with measure NH-019-10:Percent of Low Risk Residents Who Lose Control of Their Bowel or Bladder (Long Stay)] is noted and will be communicated to the business owner component of CMS.
De.4 National Priority Partners Priority Area: Care coordination
De.5 IOM Quality Domain: Patient-centered
De.6 Consumer Care Need:

CONDITIONS FOR CONSIDERATION BY NQF	
Four conditions must be met before proposed measures may be considered and evaluated for suitability as voluntary consensus standards:	NQF Staff
A. The measure is in the public domain or an intellectual property (measure steward agreement) is signed. <i>Public domain only applies to governmental organizations. All non-government organizations must sign a</i>	A Y <input type="checkbox"/>

<p><i>measure steward agreement even if measures are made publicly and freely available.</i></p> <p>A.1 Do you attest that the measure steward holds intellectual property rights to the measure and the right to use aspects of the measure owned by another entity (e.g., risk model, code set)? Yes</p> <p>A.2 Indicate if Proprietary Measure (as defined in measure steward agreement):</p> <p>A.3 Measure Steward Agreement: Government entity and in the public domain - no agreement necessary</p> <p>A.4 Measure Steward Agreement attached:</p>	<p>N <input type="checkbox"/></p>
<p>B. The measure owner/steward verifies there is an identified responsible entity and process to maintain and update the measure on a schedule that is commensurate with the rate of clinical innovation, but at least every 3 years. Yes, information provided in contact section</p>	<p>B Y <input type="checkbox"/> N <input type="checkbox"/></p>
<p>C. The intended use of the measure includes both public reporting and quality improvement. ► Purpose: Public reporting, Internal quality improvement</p>	<p>C Y <input type="checkbox"/> N <input type="checkbox"/></p>
<p>D. The requested measure submission information is complete. Generally, measures should be fully developed and tested so that all the evaluation criteria have been addressed and information needed to evaluate the measure is provided. Measures that have not been tested are only potentially eligible for a time-limited endorsement and in that case, measure owners must verify that testing will be completed within 12 months of endorsement.</p> <p>D.1 Testing: Yes, fully developed and tested</p> <p>D.2 Have NQF-endorsed measures been reviewed to identify if there are similar or related measures? Yes</p>	<p>D Y <input type="checkbox"/> N <input type="checkbox"/></p>
<p>(for NQF staff use) Have all conditions for consideration been met? Staff Notes to Steward (if submission returned):</p>	<p>Met Y <input type="checkbox"/> N <input type="checkbox"/></p>
<p>Staff Notes to Reviewers (issues or questions regarding any criteria):</p>	
<p>Staff Reviewer Name(s):</p>	

TAP/Workgroup Reviewer Name:	
Steering Committee Reviewer Name:	
1. IMPORTANCE TO MEASURE AND REPORT	
<p>Extent to which the specific measure focus is important to making significant gains in health care quality (safety, timeliness, effectiveness, efficiency, equity, patient-centeredness) and improving health outcomes for a specific high impact aspect of healthcare where there is variation in or overall poor performance. Measures must be judged to be important to measure and report in order to be evaluated against the remaining criteria. (evaluation criteria)</p> <p>1a. High Impact _____</p> <p>(for NQF staff use) Specific NPP goal:</p> <p>1a.1 Demonstrated High Impact Aspect of Healthcare: Affects large numbers, Patient/societal consequences of poor quality</p> <p>1a.2</p> <p>1a.3 Summary of Evidence of High Impact: At any given time, more than 100,000 residents in American nursing facilities have urethral catheters in place. (1) Catheters are commonly used for urinary retention, wound management, and in some circumstances, patient comfort. When not properly maintained and monitored, indwelling catheters can cause chronic pain or infections leading to a greater functional decline and decreased quality of life for the resident. (2) A thorough assessment of the resident and evaluation of the medical need for the catheter can sometimes decrease or prevent the use of catheters.</p> <p>The indwelling catheter quality measure can potentially serve as a reminder to facilities of the importance of limiting catheter use. (3) Overuse of catheters to manage incontinence, other than for short-term periods, is a</p>	<p>Ev al Rat ing</p> <p>1a C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>

Comment [KP1]: 1a. The measure focus addresses:

- a specific national health goal/priority identified by NQF's National Priorities Partners; OR
- a demonstrated high impact aspect of healthcare (e.g., affects large numbers, leading cause of morbidity/mortality, high resource use (current and/or future), severity of illness, and patient/societal consequences of poor quality).

potential sign of suboptimal care and an indication that further assessment and alternative treatment could be offered. (4) Among nursing facility residents, there is evidence that institutional policies and educational programs strongly impact care provider practices.

There are clear benefits to nursing homes conducting a thorough evaluation of the medical need for the catheterization of their residents. A determination regarding continued use or removal should be completed as soon as possible following admission. Nursing facilities need to assess the frequency of urinary catheterization practices to ensure that policies reflect current practice standards, and increase compliance with Centers for Disease Control guidelines for prevention of infection related to catheter use.(1)

Using MDS 2.0 data for April-June 2008, the national prevalence of indwelling catheters in nursing facilities was 7.7%, with a range from an average of 5.2% in Rhode Island to a high of an average of 11.3% in North Dakota.(5) National measure results have been stable over time, ranging from 5.7% in 2003 to 5.8% in 2008.(6) The current indwelling catheter quality measure is currently one of the 19 publicly reported quality measures for nursing facilities on the Centers for Medicare & Medicaid Services (CMS) Nursing Home Compare Web site.

1a.4 Citations for Evidence of High Impact: 1. Nursing Home Quality Initiative. Fast facts: urinary catheters overview. MedQIC. 2004. Available from <http://www.qualitynet.org/dcs/ContentServer?cid=1109274857368&pagename=Medqic%2FOtherResource%2FOtherResourcesTemplate&c=OtherResource>.

2. Quality Measures Management Information System (QMIS). Measure details. November 12, 2002. Available from <https://www.qualitynet.org/qmis/measureDetailView.htm?measureId=10176&viewType=0>.

3. Brega A, Hittle D, Goodrich G, Kramer A, Conway K, Levy C. Empirical review of publicly reported nursing home quality measures. Denver: Division of Health Care Policy and Research University of Colorado at Denver; Abt Associates, Inc, 2007.

4. Georgiou A, Potter J, Brocklehurst JC, Lowe D, Pearson M. Measuring the quality of urinary continence care in long-term care facilities: An analysis of outcome indicators. Age Aging. 2001;30:63-6

5. Centers for Medicare & Medicaid Services. MDS quality measure/indicator report. Available from http://www.cms.hhs.gov/MDSPubQandResRep/02_qmreport.asp?isSubmitted=qm3&group=13&qtr=14.

6. American Health Care Association. Trends in publicly reported nursing facility quality measures. July 2009. Available from http://www.ahcancal.org/research_data/trends_statistics/Documents/trends_nursing_facilities_quality_measures.pdf.

1b. Opportunity for Improvement

1b.1 Benefits (improvements in quality) envisioned by use of this measure: Facilities can use information from this measure to determine whether they may be overusing catheters for their long stay residents. Reduced use of urinary catheters, and associated problems with catheter use including pain, infections and functional decline, are the expected benefits envisioned by use of this measure.

1b.2 Summary of data demonstrating performance gap (variation or overall poor performance) across providers:

A version of the current quality measure has been in use by CMS since 2002, drawing on data from a similar but less detailed MDS 2.0 item. An analysis by the Division of Health Care Policy and Research at the University of Colorado at Denver found that the measure demonstrated very limited variability across facilities. The quality measure varied from 2.9% at the 25th percentile, to 7.7% at the 75th percentile; having an interquartile range (the 75th percentile minus the 25th percentile) of less than 5 percentage points.(1)

See attached Table 1: Measure Variability Across Facilities.

In a study to measure the quality of urinary continence care in long-term care facilities, catheterization rates were approximately 10% in nursing facilities, ranging from 0%-44% among fourteen nursing homes where data was collected on the outcome measure.(2) Thus, there was great variability in this quality measure within

1b
C
P
M
N

Comment [KP2]: 1b. Demonstration of quality problems and opportunity for improvement, i.e., data demonstrating considerable variation, or overall poor performance, in the quality of care across providers and/or population groups (disparities in care).

Comment [k3]: 1 Examples of data on opportunity for improvement include, but are not limited to: prior studies, epidemiologic data, measure data from pilot testing or implementation. If data are not available, the measure focus is systematically assessed (e.g., expert panel rating) and judged to be a quality problem.

settings. The authors were cautious to note that interpretation of the outcome results required more precise details on case-mix and the definition of outcome measures. In another study looking at state variation in indicators of quality of care in nursing facilities, limited variation among states was observed for urinary catheterization. However, among the risk-adjusted quality scores, the authors observed the most variation for urinary catheterization (an approximately twofold difference). (3)

1b.3 Citations for data on performance gap:

1. Brega A, Hittle D, Goodrich G, Kramer A, Conway K, Levy C. Empirical review of publicly reported nursing home quality measures. Denver: Division of Health Care Policy and Research University of Colorado at Denver; Abt Associates, Inc, 2007.
2. Georgiou A, Potter J, Brocklehurst JC, Lowe D, Pearson M. Measuring the quality of urinary continence care in long-term care facilities: An analysis of outcome indicators. *Age Aging*. 2001;30:63-6
3. Castle N, Degenholtz H, Engberg J. State variability in indicators of quality of care in nursing facilities. *J Gerontol*. 2005;60A(9):1173-9.

1b.4 Summary of Data on disparities by population group:

Racial segregation between nursing facilities has been shown to be a major factor in racial disparities in the nursing facility population, primarily for African Americans. In 2000, a study drawing on national MDS and Online Survey, Certification, and Reporting (OSCAR) data found that two-thirds of all black residents were living in just 10% of all facilities. (1) A 2002 survey of a stratified sample of 39 nursing facilities and 181 residential care/assisted living facilities in four states had similar findings. (2) Facilities serving African Americans have demonstrated a lower level of quality care than those serving whites with lower staff to resident ratios and higher deficiency ratings. (3) Minority groups in general and African Americans in particular have also had more limited access to nursing facility care than whites. (4)

Although research suggests racial disparities in quality of care in nursing facilities between African Americans and whites (1, 2, 3, 4), no analyses have been conducted specifically examining racial disparities in catheterization use. No other research has been conducted on other types of disparities (e.g., ethnicity, rural/urban, or income) for this measure.

1b.5 Citations for data on Disparities:

1. Smith D, Feng Z, Fennell M, Zinn J, Mor V. Separate and unequal: racial segregation and disparities in quality across U.S. nursing homes. *Health Aff (Millwood)*. 2007;26(5):1448-558.
2. Howard D, Sloane P, Zimmerman S, Eckert J, Walsh J, Buie V, Taylor P, Koch G. Distribution of African Americans in residential care/assisted living and nursing homes: more evidence of racial disparity? *Am J Public Health*. 2002;92(8):1272-7.
3. Grabowski D. The admission of blacks to high-deficiency nursing homes. *Med Care*. 2004;42(5):456-64.
4. National Center for Health Statistics (NCHS). *Health, United States, 1996-97, and injury chartbook*. Hyattsville, MD: NCHS, 1997.

1c. Outcome or Evidence to Support Measure Focus

1c.1 Relationship to Outcomes (For non-outcome measures, briefly describe the relationship to desired outcome. For outcomes, describe why it is relevant to the target population): The benefits of limiting catheter use in nursing facilities are well documented in the literature. Catheters are commonly used for urinary retention, wound management, and in some circumstances, patient comfort. When not properly maintained and monitored, indwelling catheters can cause chronic pain or infections leading to a greater functional decline and decreased quality of life for the resident. (1) Indwelling urinary catheterization can frequently causes bacteremia, or in many cases, urinary tract infections, in the elderly. Catheterization causes bacteremia to occur at a rate of 3 to 10 percent of patients per day; a single in and out catheterization may cause bacteremia in as many as 20 percent of patients (2). At least 40% of all infections seen in the nursing homes are in the urinary tract system; of those infections, 80% are due to urinary tract catheterization and instrumentation (3).

1c
 C
 P
 M
 N

Comment [k4]: 1c. The measure focus is:
 •an outcome (e.g., morbidity, mortality, function, health-related quality of life) that is relevant to, or associated with, a national health goal/priority, the condition, population, and/or care being addressed;
 OR
 •if an intermediate outcome, process, structure, etc., there is evidence that supports the specific measure focus as follows:
 oIntermediate outcome - evidence that the measured intermediate outcome (e.g., blood pressure, Hba1c) leads to improved health/avoidance of harm or cost/benefit.
 oProcess - evidence that the measured clinical or administrative process leads to improved health/avoidance of harm and if the measure focus is on one step in a multi-step care process, it measures the step that has the greatest effect on improving the specified desired outcome(s).
 oStructure - evidence that the measured structure supports the consistent delivery of effective processes or access that lead to improved health/avoidance of harm or cost/benefit.
 oPatient experience - evidence that an association exists between the measure of patient experience of health care and the outcomes, values and preferences of individuals/ the public.
 oAccess - evidence that an association exists between access to a health service and the outcomes of, or experience with, care.
 oEfficiency - demonstration of an association between the measured resource use and level of performance with respect to one or more of the other five IOM aims of quality.

Comment [k5]: 4 Clinical care processes typically include multiple steps: assess → identify problem/potential problem → choose/plan intervention (with patient input) → provide intervention → evaluate impact on health status. If the measure focus is one step in such a multi-step process, the step with the greatest effect on the desired outcome should be selected as the focus of measurement. For example, although assessment of immunization status and recommending immunization are necessary steps, they are not sufficient to achieve the desired impact on health status - patients must be vaccinated to achieve immunity. This does not preclude consideration of measures of preventive screening interventions where there is a strong link with desired outcomes (e.g., mammography) or measures for multiple care processes that affect a single outcome.

Many times residents are admitted to a nursing facility from hospitals with catheters in place, and the facility must make a determination whether or not to continue use of the device. A thorough assessment of the resident and evaluation of the medical need for the catheter can sometimes decrease or prevent the use of catheters and the risks associated with their use.

1. Gammack JK. Use of management of chronic urinary catheters in long-term care: much controversy, little consensus. *J Am Med Dir Assoc.* 2003;4(2 Supp):S52-9.

2. 12: Kamel HK. Managing urinary tract infections in the nursing home: Myths, mysteries and realities. *Int J Geriatr Gerontol.* 2004;1(2). Available from http://www.ispub.com/journal/the_internet_journal_of_geriatrics_and_gerontology/volume_1_number_2_21/article/managing_urinary_tract_infections_in_the_nursing_home_myths_mysteries_and_realities.html

3. Newman DK, Fader M, Bliss DZ. Managing incontinence using technology, devices and products. *Nursing Research.* 2004; 53(6 Suppl): 42-48.

4. Quality Measures Management Information System (QMIS). Measure details. November 12, 2002. Available from <https://www.qualitynet.org/qmis/measureDetailView.htm?measureId=10176&viewType=0>.

1c.2-3. Type of Evidence: Evidence-based guideline

1c.4 Summary of Evidence (as described in the criteria; for outcomes, summarize any evidence that healthcare services/care processes influence the outcome):

There are clear benefits to nursing homes conducting a thorough evaluation of the medical need for the catheterization of their residents. A determination regarding continued use or removal should be completed as soon as possible following admission. Nursing facilities need to assess the frequency of urinary catheterization practices to ensure that policies reflect current practice standards, and increase compliance with Centers for Disease Control guidelines for prevention of infection related to catheter use.(1)

1. Quality Measures Management Information System (QMIS). Measure details. November 12, 2002. Available from <https://www.qualitynet.org/qmis/measureDetailView.htm?measureId=10176&viewType=0>.

1c.5 Rating of strength/quality of evidence (also provide narrative description of the rating and by whom):
The evidence was not rated.

1c.6 Method for rating evidence:

1c.7 Summary of Controversy/Contradictory Evidence: No contradictory evidence has been identified.

1c.8 Citations for Evidence (other than guidelines):

1c.9 Quote the Specific guideline recommendation (including guideline number and/or page number):

1. Guideline for Prevention of Catheter-Associated Urinary Tract Infections 2009. Centers for Disease Control and Prevention. http://www.cdc.gov/ncidod/dhqp/pdf/guidelines/CAUTI_Guideline2009final.pdf
2. Urinary Incontinence Clinical Practice Guideline. Contains a table for appropriate indications for chronic indwelling catheters. Purchase information available at: <http://www.amda.com/tools/cpg/incontinence.cfm>

1c.10 Clinical Practice Guideline Citation: 1. Guideline for Prevention of Catheter-Associated Urinary Tract Infections 2009 http://www.cdc.gov/ncidod/dhqp/pdf/guidelines/CAUTI_Guideline2009final.pdf

2. Urinary Incontinence Clinical Practice Guideline. <http://www.amda.com/tools/cpg/incontinence.cfm>

1c.11 National Guideline Clearinghouse or other URL:

<http://www.ncbi.nlm.nih.gov/bookshelf/br.fcgi?book=hsarchive&part=A9995> and

1c.12 Rating of strength of recommendation (also provide narrative description of the rating and by whom):

The HICPAC (Healthcare Infection Control Practices Advisory Committee) reviewed the evidence supporting the guideline recommendations using an adapted GRADE system and rated the evidence underlying the specific recommendations from 1A (the highest level of evidence) to 2.

Comment [k6]: 3 The strength of the body of evidence for the specific measure focus should be systematically assessed and rated (e.g., USPSTF grading system

<http://www.ahrq.gov/clinic/uspstf07/methods/benefit.htm>). If the USPSTF grading system was not used, the grading system is explained including how it relates to the USPSTF grades or why it does not. However, evidence is not limited to quantitative studies and the best type of evidence depends upon the question being studied (e.g., randomized controlled trials appropriate for studying drug efficacy are not well suited for complex system changes). When qualitative studies are used, appropriate qualitative research criteria are used to judge the strength of the evidence.

Comment [k7]: USPSTF grading system <http://www.ahrq.gov/clinic/uspstf/grades.htm>: A - The USPSTF recommends the service. There is high certainty that the net benefit is substantial. B - The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial. C - The USPSTF recommends against routinely providing the service. There may be considerations that support providing the service in an individual patient. There is at least moderate certainty that the net benefit is small. Offer or provide this service only if other considerations support the offering or providing the service in an individual patient. D - The USPSTF recommends against the service. There is moderate or high certainty that the service has no net benefit or that the harms outweigh the benefits. I - The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined.

<p>1c.13 Method for rating strength of recommendation (<i>If different from USPSTF system, also describe rating and how it relates to USPSTF</i>): HICPAC used an adapted version of the GRADE Working group system.(1)</p> <p>1. Atkins D, Best D, Briss PA, et al. Grading quality of evidence and strength of recommendations. <i>BMJ</i>. 2004;336(7652):1049-51.</p> <p>1c.14 Rationale for using this guideline over others: No contradictory evidence has been identified.</p>	
<p>TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for <i>Importance to Measure and Report</i>?</p>	1
<p>Steering Committee: Was the threshold criterion, <i>Importance to Measure and Report</i>, met? Rationale:</p>	<p>1 Y <input type="checkbox"/> N <input type="checkbox"/></p>
<p style="text-align: center;">2. SCIENTIFIC ACCEPTABILITY OF MEASURE PROPERTIES</p>	
<p>Extent to which the measure, <u>as specified</u>, produces consistent (reliable) and credible (valid) results about the quality of care when implemented. (evaluation criteria)</p>	<p style="text-align: center;">Ev al Rat ing</p>
<p style="text-align: center;">2a. MEASURE SPECIFICATIONS</p>	
<p>S.1 Do you have a web page where current detailed measure specifications can be obtained? S.2 If yes, provide web page URL:</p> <p>2a. Precisely Specified </p>	
<p>2a.1 Numerator Statement (<i>Brief, text description of the numerator - what is being measured about the target population, e.g. target condition, event, or outcome</i>): The numerator statement refers to a catheter that was inserted and left in the bladder by the facility during the assessment period.</p> <p>During MDS 3.0 field testing, look-back periods were highlighted as a significant issue across the assessment tool. For clinical assessment items, longer look-back periods served to increase the amount of record review, increasing assessment burden and leading to more opportunities for error. During national testing of look-back periods for the MDS 3.0 proposed items, the 5-day look-back period performed well and likely contributed to the improved reliability of this item.(1)</p> <p>1. Saliba D, Buchanan J. Development and Validation of a Revised Nursing Home Assessment Tool: MDS 3.0. Contract No. 500-00-0027/Task Order #2. Santa Monica, CA: Rand Corporation, Apr 2008. Available from http://www.cms.hhs.gov/NursingHomeQualityInits/Downloads/MDS30FinalReport.pdf.</p> <p>The numerator is the number of long-stay residents who have/had a urinary catheter in the last 5 days (H0100A is checked).</p> <p>2a.2 Numerator Time Window (<i>The time period in which cases are eligible for inclusion in the numerator</i>): Numerator data come from MDS 3.0 annual, quarterly, significant change or significant correction assessment conducted during each quarter (3-month period).</p> <p>2a.3 Numerator Details (<i>All information required to collect/calculate the numerator, including all codes, logic, and definitions</i>): Residents are counted if they are long-stay residents, defined as residents whose length of stay is greater than 100 days. Residents who return to the nursing home following a hospital discharge will not have their stay reset to zero. The numerator includes residents who have indwelling catheters (H0100A is checked) on the</p>	<p>2a- spe cs C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> <input type="checkbox"/></p>

Comment [KP8]: 2a. The measure is well defined and precisely specified so that it can be implemented consistently within and across organizations and allow for comparability. The required data elements are of high quality as defined by NQF's Health Information Technology Expert Panel (HITEP) .

most recent MDS 3.0 assessment (which may be an annual, quarterly, significant change or significant correction assessment). Exclusions are assessments where data for the urinary catheter item (H0100) is missing. Also, residents with diagnoses of neurogenic bladder (item I1550) or obstructive uropathy (item I1650) are excluded because these are conditions in which the person is unable to empty the bladder voluntarily or effectively, putting the person at risk or complications, such as overflow incontinence, recurrent infection, vesicoureteral reflux, or autonomic dysreflexia. 2a.8. (denominator details). Residents are counted if they are long-stay residents defined as residents whose length of stay is greater than 100 days. Residents who return to the nursing home following a hospital discharge will not have their day count reset to zero. The target population includes all long-stay residents who have had an annual, quarterly, significant change or significant correction MDS 3.0 assessment (A0130.A= 02,03,04,05 or 06) during the selected quarter, except for those who meet the exclusion criteria or have missing data in the responses to the relevant items in the MDS.

2a.4 Denominator Statement (*Brief, text description of the denominator - target population being measured*):

The denominator is the total of all long-stay residents in the nursing home who have been assessed with an annual, quarterly, significant change or significant correction MDS 3.0 assessment during the quarter (3-month period) and who do not meet the exclusion criteria.

2a.5 Target population gender: Male, Female

2a.6 Target population age range: The target population includes all long-stay residents of any age residing in the nursing facility.

2a.7 Denominator Time Window (*The time period in which cases are eligible for inclusion in the denominator*):

Denominator data come from MDS 3.0 annual, quarterly, significant change or significant correction assessment conducted during each quarter (3-month period).

2a.8 Denominator Details (*All information required to collect/calculate the denominator - the target population being measured - including all codes, logic, and definitions*):

Residents are counted if they are long-stay residents defined as residents whose length of stay is greater than 100 days. Residents who return to the nursing home following a hospital discharge will not have their day count reset to zero. The target population includes all long-stay residents who have had an annual, quarterly, significant change or significant correction MDS 3.0 assessment (A0130.A= 02,03,04,05 or 06) during the selected quarter, except for those who meet the exclusion criteria or have missing data in the responses to the relevant items in the MDS.

2a.9 Denominator Exclusions (*Brief text description of exclusions from the target population*): A resident is excluded from the denominator if the MDS assessment was conducted within 14 days of admission or if there is missing data in the responses to the relevant questions in the MDS assessment. Other exclusions include residents with neurogenic bladder or obstructive uropathy. Residents with diagnoses of neurogenic bladder (item I1550) or obstructive uropathy (item I1650) are excluded because these are conditions in which the person is unable to empty the bladder voluntarily or effectively, putting the person at risk of complications, such as overflow incontinence, recurrent infection, vesicoureteral reflux, or autonomic dysreflexia.

Facilities are excluded from public reporting if they have fewer than 30 residents due to small sample size.

2a.10 Denominator Exclusion Details (*All information required to collect exclusions to the denominator, including all codes, logic, and definitions*):

1. The target assessment is an OBRA admission assessment (item A0310A = 01).
2. There is missing data on indwelling catheter (item H0100A).
3. Residents with neurogenic bladder (item I1550) or obstructive uropathy (item I1650). Residents with diagnoses of neurogenic bladder (item I1550 on the MDS 3.0) and obstructive uropathy (item I1650 on the MDS 3.0) are excluded because these are conditions in which the person is unable to empty the bladder voluntarily or effectively, putting the person at risk of complications, such as overflow incontinence, recurrent infection, vesicoureteral reflux, or autonomic dysreflexia.

2a.11 Stratification Details/Variables (*All information required to stratify the measure including the stratification variables, all codes, logic, and definitions*):
This is not applicable.

Comment [k9]: 11 Risk factors that influence outcomes should not be specified as exclusions.
12 Patient preference is not a clinical exception to eligibility and can be influenced by provider interventions.

2a.12-13 Risk Adjustment Type:

2a.14 Risk Adjustment Methodology/Variables (*List risk adjustment variables and describe conceptual models, statistical models, or other aspects of model or method*):

Resident-level limited covariate risk adjustment for residents who are bowel incontinent on prior MDS (item H0400 = 2 or 3), or had pressure sores at stage 2, 3, or 4 on prior MDS (M0300B1 > 0 or M0300C1 > 0 or M0300D1 > 0).

2a.15-17 Detailed risk model available Web page URL or attachment: URL

<http://www.cms.hhs.gov/NursingHomeQualityInits/downloads/NHQIQMUsersManual.pdf>

2a.18-19 Type of Score: Ratio

2a.20 Interpretation of Score:

2a.21 Calculation Algorithm (*Describe the calculation of the measure as a flowchart or series of steps*):

For each facility, the number of long-stay residents meeting the numerator criteria and the number of (non-excluded) residents meeting the denominator criteria are counted. The facility observed score for the measure is a prevalence score calculated as the number of residents in the facility in the numerator divided by all non-excluded residents in the denominator. The number of long-stay residents meeting the numerator criteria and the number of residents meeting the denominator criteria are also counted for the covariate measure, which is bowel incontinence or presence of pressure sores, as reported on the resident's prior MDS assessment.

The covariate scores are then entered into a logistic regression equation, and the result is an expected score for the resident for that quality measure (QM). The logistic regression equations are of the form: where e is the base of natural logarithms and x is a linear combination of the logistic regression coefficients and the covariate scores of the form:

$C0 + C1*COVA + C2*COVB + \dots$ where C0 is the logistic regression constant, C1 is the logistic regression coefficient for the first covariate (where applicable), COVA is the resident-level score for the first covariate, C2 is the logistic regression coefficient for the second covariate, and COVB is the resident-level score for the second covariate (where applicable), etc. The regression constant and regression coefficients are numbers obtained through statistical logistic regression analysis.(1)

The expected score for the measure is then calculated as the expected number of residents in the facility meeting the numerator criteria divided by all non-excluded residents in the denominator.

1. Abt Associates, Inc. National nursing home quality measures: user's manual. (2004). Cambridge, MA: Abt Associates, Inc., 2004.

2a.22 Describe the method for discriminating performance (*e.g., significance testing*):

Because the computed scores are not estimates, but include all residents who meet the measure criteria, in terms of discriminating performance, the computed scores can be used to make valid comparisons.

2a.23 Sampling (Survey) Methodology *If measure is based on a sample (or survey), provide instructions for obtaining the sample, conducting the survey and guidance on minimum sample size (response rate):*

This is not applicable.

2a.24 Data Source (*Check the source(s) for which the measure is specified and tested*)

Electronic clinical data

2a.25 Data source/data collection instrument (*Identify the specific data source/data collection instrument, e.g. name of database, clinical registry, collection instrument, etc.*):

The proposed data source is the Nursing Home MDS 3.0.

2a.26-28 Data source/data collection instrument reference web page URL or attachment: URL

http://www.cms.hhs.gov/NursingHomeQualityInits/25_NHQIMDS30.asp#TopOfPage

2a.29-31 Data dictionary/code table web page URL or attachment: URL

http://www.cms.hhs.gov/NursingHomeQualityInits/25_NHQIMDS30.asp#TopOfPage

2a.32-35 Level of Measurement/Analysis (*Check the level(s) for which the measure is specified and tested*)

downloaded from MDS assessment records. Analyses were based on complete MDS data from January 2005 through March 2006, as well as nearly complete data for April 2006 and partial data for May and June 2006. (2)

The national test of MDS 3.0 items examined agreement between assessors (reliability); validity of new cognitive, depression, and behavior items; response rates for interview items; user satisfaction and feedback on changes; and time to complete the assessment. The network of Quality Improvement Organizations (QIOs) was employed to identify gold-standard (research) nurses and recruit community nursing facilities to participate in the national evaluation, including a representative sample of for-profit and not-for-profit facilities and hospital-based and freestanding facilities. The gold-standard nurses were trained in the MDS 3.0 instrument and, in turn, trained a facility nurse from each participating nursing facility in their home states.

1. Abt Associates, Inc.; Stepwise Systems, Inc.; Qualidigm. Data Assessment and Verification (DAVE 2) project—MDS two-stage discrepancy findings, April-December 2006. Cambridge, MA: Abt Associates, Inc, 2007.
2. Brega A, Hittle D, Goodrich G, Kramer A, Conway K, Levy C. Empirical review of publicly reported nursing home quality measures. Denver: Division of Health Care Policy and Research University of Colorado at Denver; Abt Associates, Inc, 2007.
3. Saliba D, Buchanan J. Development and validation of a revised nursing home assessment tool: MDS 3.0. Contract No. 500-00-0027/Task Order #2. Santa Monica, CA: Rand Corporation, Apr 2008. Available from <http://www.cms.hhs.gov/NursingHomeQualityInits/Downloads/MDS30FinalReport.pdf>.

2b.3 Testing Results (*reliability statistics, assessment of adequacy in the context of norms for the test conducted*):

As part of the DAVE 2 project, Abt Associates assessed the reliability of the MDS 2.0 quality measures. (1) For each MDS data element, the rate of discrepancies between the reconciled and original facility assessments has been reported. For catheter use, the two-stage review discrepancy rate was 0.0%, which the University of Colorado deemed performed well on the indicator of reliability. (2)

Second, in terms of measure stability, the University of Colorado examined the percentage of facilities that had a change in ranking from one quarter to the next of at least three deciles. (2) They found that facility catheter rates for this measure were unstable over time: 18.9% of facilities had a three-decile-or-more change from one quarter to the next quarter.

Third, in the national analysis of assessing the reliability of the MDS 3.0 conducted by the RAND Corporation, agreement between MDS 3.0 assessors on bladder and bowel items, including catheter use, was excellent. The average kappa for the gold-standard nurse to gold-standard nurse agreement was 0.949, and the average kappa for the gold-standard nurse to facility nurse agreement was 0.945. (3)

1. Abt Associates, Inc.; Stepwise Systems, Inc.; Qualidigm. Data Assessment and Verification (DAVE 2) project—MDS two-stage discrepancy findings, April-December 2006. Cambridge, MA: Abt Associates, Inc, 2007.
2. Brega A, Hittle D, Goodrich G, Kramer A, Conway K, Levy C. Empirical review of publicly reported nursing home quality measures. Denver: Division of Health Care Policy and Research University of Colorado at Denver; Abt Associates, Inc, 2007.
3. Saliba D, Buchanan J. Development and validation of a revised nursing home assessment tool: MDS 3.0. Contract No. 500-00-0027/Task Order #2. Santa Monica, CA: Rand Corporation, Apr 2008. Available from <http://www.cms.hhs.gov/NursingHomeQualityInits/Downloads/MDS30FinalReport.pdf>.

2c. Validity testing

2c.1 Data/sample (*description of data/sample and size*): The data came from two sources: national facility-level quality measure data from 2003 Q3 through 2006 Q3 came from the QIES MDS Express Reports on the CMS intranet; OSCAR data related to facility characteristics (e.g., state, resident census, number of beds, staffing) and certification survey results were downloaded from QIES Workbench. A 10% random sample of all Medicare-certified nursing facilities was also downloaded from MDS assessment records. Analyses were based on complete MDS data from January 2005 through March 2006, as well as nearly complete data for April 2006 and partial data for May and June 2006.

2c.2 Analytic Method (*type of validity & rationale, method for testing*):

The analysis evaluated measure validity in a number of ways: to examine the expected positive influence of public reporting on quality of care, an assessment of the degree to which quality measure triggering rates

2c
C
P
M
N

Comment [KP12]: 2c. Validity testing demonstrates that the measure reflects the quality of care provided, adequately distinguishing good and poor quality. If face validity is the only validity addressed, it is systematically assessed.

Comment [k13]: 9 Examples of validity testing include, but are not limited to: determining if measure scores adequately distinguish between providers known to have good or poor quality assessed by another valid method; correlation of measure scores with another valid indicator of quality for the specific topic; ability of measure scores to predict scores on some other related valid measure; content validity for multi-item scales/tests. Face validity is a subjective assessment by experts of whether the measure reflects the quality of care (e.g., whether the proportion of patients with BP < 140/90 is a marker of quality). If face validity is the only validity addressed, it is systematically assessed (e.g., ratings by relevant stakeholders) and the measure is judged to represent quality care for the specific topic and that the measure focus is the most important aspect of quality for the specific topic.

have improved over time; to evaluate convergent validity, an assessment of the correlation of the quality measure with all other measures; and to determine if the quality measure triggering rate was influenced by factors that are unrelated to facility quality, an evaluation of seasonal variations in triggering rates across the 13 quarters of data. The analysis also computed descriptive statistics and conducted a one-way analysis of variance (ANOVA) for the measure to examine the amount of variance in triggering rates explained by the state in which a facility was located.

2c.3 Testing Results (*statistical results, assessment of adequacy in the context of norms for the test conducted*):

The trend shows a seasonal fluctuation in the rate for indwelling catheter over time, but from one year's quarter to the next, there is essentially no difference. There is a distinct peak in the first quarter of each year, followed by a relatively flat rate trend in the other three quarters. This pattern is similar to seasonal variation in hospital and skilled nursing facility utilization, indicating that it may reflect seasonal variations in the general health of the nursing home population rather than seasonal variation in nursing home quality.(1)

See attached Table 2: Measure Trends Over Time.

In a clinical review conducted by the University of Colorado, participants expressed concern about the inability to exclude residents for whom an indwelling catheter is a necessary component of high-quality medical care.(2) To ensure that facilities are not penalized for having a large population of residents who meet this criteria, participants recommended the exclusion of residents with obstructive uropathy or neurogenic bladder. The proposed MDS 3.0 measure has been revised based on this recommendation.

1. Brega A, Hittle D, Goodrich G, Kramer A, Conway K, Levy C. Empirical review of publicly reported nursing home quality measures. Denver: Division of Health Care Policy and Research University of Colorado at Denver; Abt Associates, Inc, 2007.
2. Brega A, Levy C, Kramer A, Eilertsen T, Hittle D, Goodrich G. Limited clinical review of publicly reported nursing home quality measures. Denver: Division of Health Care Policy and Research University of Colorado at Denver, 2007.

2d. Exclusions Justified

2d.1 Summary of Evidence supporting exclusion(s):

All long-stay residents for whom complete data exists are included. Post-acute care residents are not included because they are likely to have had an indwelling catheter prior to their nursing facility stay in the hospital or other acute setting. A Technical Expert Panel (TEP) convened in January 2009 expressed the need to exclude residents with medical conditions requiring catheters. There are some cases in which catheter use is warranted because of medical conditions that may be untreatable in a nursing facility setting. Therefore, residents with diagnoses of neurogenic bladder and obstructive uropathy are excluded because these are conditions in which the person is unable to empty the bladder voluntarily or effectively, putting the person at risk of complications, such as overflow incontinence, recurrent infection, vesicoureteral reflux, or autonomic dysreflexia.

Excluding missing data for existing quality measures is standard practice and was initially endorsed by NQF. Missing data is excluded from the calculation of the quality measures for several reasons. 1) There are legitimate reasons for facility staff not to select a 'dash' rather than a response; for example, if a resident is discharged or transferred abruptly, the staff may not be able to complete all items, however, an assessment is required for payment. The intent of the 'dash' is to allow the facility to submit an assessment when the staff are unable to complete the entire assessment. 2) Historically there has been very little missing data. For example, the current quality measure "Percent of residents who were physically restrained", is based on three fields on the MDS 3.0. For all of the non-admission target assessments for calendar year 2009, there were 5,242,022 such assessments and 629 assessments (0.012%) had a dash for one or more of the three fields for the physical restraint measure. 3) We remain concerned about a change in measure definition that may result in incentivizing the facility staff to fill in a response to avoid a missing item. We believe that the result will lead to decreased validity and usefulness of the measure.

2d.2 Citations for Evidence:

This is not applicable.

2d
C
P
M
N
NA

Comment [KP14]: 2d. Clinically necessary measure exclusions are identified and must be:
 •supported by evidence of sufficient frequency of occurrence so that results are distorted without the exclusion;
 AND

•a clinically appropriate exception (e.g., contraindication) to eligibility for the measure focus;
 AND
 •precisely defined and specified:
 –if there is substantial variability in exclusions across providers, the measure is specified so that exclusions are computable and the effect on the measure is transparent (i.e., impact clearly delineated, such as number of cases excluded, exclusion rates by type of exclusion);

if patient preference (e.g., informed decision-making) is a basis for exclusion, there must be evidence that it strongly impacts performance on the measure and the measure must be specified so that the information about patient preference and the effect on the measure is transparent (e.g., numerator category computed separately, denominator exclusion category computed separately).

Comment [k15]: 10 Examples of evidence that an exclusion distorts measure results include, but are not limited to: frequency of occurrence, sensitivity analyses with and without the exclusion, and variability of exclusions across providers.

<p>2d.3 Data/sample (<i>description of data/sample and size</i>): This is not applicable.</p> <p>2d.4 Analytic Method (<i>type analysis & rationale</i>): This is not applicable.</p> <p>2d.5 Testing Results (<i>e.g., frequency, variability, sensitivity analyses</i>): This is not applicable.</p>	
<p>2e. Risk Adjustment for Outcomes/ Resource Use Measures</p> <p>2e.1 Data/sample (<i>description of data/sample and size</i>): The data came from two sources: national facility-level quality measure data from 2003 Q3 through 2006 Q3 came from the QIES MDS Express Reports on the CMS intranet; OSCAR data related to facility characteristics (e.g., state, resident census, number of beds, staffing) and certification survey results were downloaded from QIES Workbench. A 10% random sample of all Medicare-certified nursing facilities was also downloaded from MDS assessment records. Analyses were based on complete MDS data from January 2005 through March 2006, as well as nearly complete data for April 2006 and partial data for May and June 2006.</p> <p>2e.2 Analytic Method (<i>type of risk adjustment, analysis, & rationale</i>): Bowel incontinence and pressure ulcers on prior MDS are risk factors for this measure. The University of Colorado attempted to improve risk adjustment for this measure. They found bowel incontinence to be a valuable covariate and recommended retaining it as a risk adjuster and further evaluating the use of pressure ulcers as a risk adjuster to indwelling catheter use.(1)</p> <p>1. Brega A, Hittle D, Goodrich G, Kramer A, Conway K, Levy C. Empirical review of publicly reported nursing home quality measures. Denver: Division of Health Care Policy and Research University of Colorado at Denver; Abt Associates, Inc, 2007.</p> <p>2e.3 Testing Results (<i>risk model performance metrics</i>): Although the U. of Colorado's risk adjustment model had better fit statistics than the unadjusted measure, the improvement was insufficient for meeting their criteria for predictive performance (C = 0.6960, R2 = 0.0790). A C-statistic equal to or greater than 0.70 met the University of Colorado's criteria for risk-adjustment adequacy, but they also required an R2 value of 0.10 or greater. Thus, while the C-statistic was close to an acceptable threshold for risk adjustment adequacy the R2 value was not.</p> <p>2e.4 If outcome or resource use measure is not risk adjusted, provide rationale:</p>	<p>2e C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/></p>
<p>2f. Identification of Meaningful Differences in Performance</p> <p>2f.1 Data/sample from Testing or Current Use (<i>description of data/sample and size</i>): The data came from two sources: national facility-level quality measure data from 2003 Q3 through 2006 Q3 came from the QIES MDS Express Reports on the CMS intranet; OSCAR data related to facility characteristics (e.g., state, resident census, number of beds, staffing) and certification survey results were downloaded from QIES Workbench. A 10% random sample of all Medicare-certified nursing facilities was also downloaded from MDS assessment records. Analyses were based on complete MDS data from January 2005 through March 2006, as well as nearly complete data for April 2006 and partial data for May and June 2006.</p> <p>2f.2 Methods to identify statistically significant and practically/meaningfully differences in performance (<i>type of analysis & rationale</i>): Because the computed scores are not estimates, but include all residents who meet the measure criteria, in terms of discriminating performance, the computed scores can be used to make valid comparisons.</p> <p>2f.3 Provide Measure Scores from Testing or Current Use (<i>description of scores, e.g., distribution by quartile, mean, median, SD, etc.; identification of statistically significant and meaningfully differences in performance</i>): An analytic team at the University of Colorado Health Sciences Center examined the rates for the measure at the facility level. Below are the measure scores from testing or current use (Description of scores, e.g., distribution by quartile, mean, median, standard deviation, etc.; identification of statistically significant and meaningfully differences in performance). For 11,928 facilities, the mean rate was 5.6% with a standard</p>	<p>2f C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>

Comment [KP16]: 2e. For outcome measures and other measures (e.g., resource use) when indicated:
•an evidence-based risk-adjustment strategy (e.g., risk models, risk stratification) is specified and is based on patient clinical factors that influence the measured outcome (but not disparities in care) and are present at start of care; OR rationale/data support no risk adjustment.

Comment [k17]: 13 Risk models should not obscure disparities in care for populations by including factors that are associated with differences/inequalities in care such as race, socioeconomic status, gender (e.g., poorer treatment outcomes of African American men with prostate cancer, inequalities in treatment for CVD risk factors between men and women). It is preferable to stratify measures by race and socioeconomic status rather than adjusting out differences.

Comment [KP18]: 2f. Data analysis demonstrates that methods for scoring and analysis of the specified measure allow for identification of statistically significant and practically/clinically meaningful differences in performance.

Comment [k19]: 14 With large enough sample sizes, small differences that are statistically significant may or may not be practically or clinically meaningful. The substantive question may be, for example, whether a statistically significant difference of one percentage point in the percentage of patients who received smoking cessation counseling (e.g., 74% v. 75%) is clinically meaningful; or whether a statistically significant difference of \$25 in cost for an episode of care (e.g., \$5,000 v. \$5,025) is practically meaningful. Measures with overall poor performance may not demonstrate much variability across providers.

deviation of 4.0%.(1) See attached Table 1: Measure Variability Across Facilities. 1. Brega A, Hittle D, Goodrich G, Kramer A, Conway K, Levy C. Empirical review of publicly reported nursing home quality measures. Denver: Division of Health Care Policy and Research University of Colorado at Denver; Abt Associates, Inc, 2007.	
2g. Comparability of Multiple Data Sources/Methods 2g.1 Data/sample (description of data/sample and size): This is not applicable. 2g.2 Analytic Method (type of analysis & rationale): This is not applicable. 2g.3 Testing Results (e.g., correlation statistics, comparison of rankings): This is not applicable.	2g C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/>
2h. Disparities in Care 2h.1 If measure is stratified, provide stratified results (scores by stratified categories/cohorts): The measure is not stratified. 2h.2 If disparities have been reported/identified, but measure is not specified to detect disparities, provide follow-up plans: While MDS 3.0 collects data on the resident's race there are no current plans to stratify the measure by race because facilities tend to be homogenous by race, making disparities generally evident in the rating of the facility.(1) 1. Smith D, Feng Z, Zinn J, Mor V. Racial disparities in access to long-term care: the illusive pursuit of equity. J Health Polit Policy Law. 2008;33(5):861-81.	2h C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/>
TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for <i>Scientific Acceptability of Measure Properties</i> ?	2
Steering Committee: Overall, to what extent was the criterion, <i>Scientific Acceptability of Measure Properties</i> , met? Rationale:	2 C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
3. USABILITY	
Extent to which intended audiences (e.g., consumers, purchasers, providers, policy makers) can understand the results of the measure and are likely to find them useful for decision making. (evaluation criteria)	Ev al Rat ing
3a. Meaningful, Understandable, and Useful Information 3a.1 Current Use: In use 3a.2 Use in a public reporting initiative (disclosure of performance results to the public at large) (If used in a public reporting initiative, provide name of initiative(s), locations, Web page URL(s). If not publicly reported, state the plans to achieve public reporting within 3 years): Nursing Home Compare http://www.medicare.gov/NHCompare/Include/DataSection/Questions/SearchCriteriaNEW.asp?version=default&t&brower=IE%7C6%7CWInXP&language=English&defaultststatus=0&pagelist=Home&CookiesEnabledStatus=True	3a C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>

Comment [KP20]: 2g. If multiple data sources/methods are allowed, there is demonstration they produce comparable results.

Comment [KP21]: 2h. If disparities in care have been identified, measure specifications, scoring, and analysis allow for identification of disparities through stratification of results (e.g., by race, ethnicity, socioeconomic status, gender);OR rationale/data justifies why stratification is not necessary or not feasible.

Comment [KP22]: 3a. Demonstration that information produced by the measure is meaningful, understandable, and useful to the intended audience(s) for both public reporting (e.g., focus group, cognitive testing) and informing quality improvement (e.g., quality improvement initiatives). An important outcome that may not have an identified improvement strategy still can be useful for informing quality improvement by identifying the need for and stimulating new approaches to improvement.

<p>3a.3 If used in other programs/initiatives (If used in quality improvement or other programs/initiatives, name of initiative(s), locations, Web page URL(s). If not used for QI, state the plans to achieve use for QI within 3 years): CMS expects that the quality measure will be used by nursing facilities as a tool to evaluate their own performance and develop quality improvement programs.</p> <p>Testing of Interpretability (Testing that demonstrates the results are understood by the potential users for public reporting and quality improvement)</p> <p>3a.4 Data/sample (description of data/sample and size): A recent study examined whether consumers could accurately interpret the quality information given for all the measures reported by Nursing Home Compare. Data were collected from 4,754 family members of nursing home residents.(1)</p> <p>1. Castle N. The Nursing Home Compare report card: consumers' use and understanding. J Aging Soc Policy. 2009;21(2), 187-208.</p> <p>3a.5 Methods (e.g., focus group, survey, QI project): A comprehension index was used to examine whether the information contained in Nursing Home Compare for each quality measure was understood by family members.</p> <p>3a.6 Results (qualitative and/or quantitative results and conclusions): The study found that 31% of the consumers used the Internet in choosing a nursing home, 12% recalled using Nursing Home Compare, and in general, the consumers' comprehension index scores were high, indicating good understanding. However, this specific measure was not evaluated.</p>	
<p>3b/3c. Relation to other NQF-endorsed measures</p> <p>3b.1 NQF # and Title of similar or related measures: The proposed measure is intended to replace the currently endorsed NQF measure #0184, Residents who have a catheter in their bladder at any time during the 14 day assessment period because the data source has changed; the MDS 2.0 is being replaced with the MDS 3.0. NQF #0138-Urinary catheter-associated urinary tract infection for intensive care unit (ICU) patients. NQF #0453 -Urinary catheter removed on Postoperative Day 1 (POD1) or Postoperative Day 2 (POD2) with day of surgery being day zero. NQF #0183- Low-risk residents who frequently lose control of their bowel or bladder. (This quality measure is paired with the current measure.) NQF#0184 Home health care: percentage of patients with improvement in urinary incontinence.</p>	
<p>(for NQF staff use) Notes on similar/related endorsed or submitted measures:</p>	
<p>3b. Harmonization</p> <p>If this measure is related to measure(s) already endorsed by NQF (e.g., same topic, but different target population/setting/data source or different topic but same target population):</p> <p>3b.2 Are the measure specifications harmonized? If not, why? This measure specifically addresses urinary catheter use in the last 5 days for the long-stay nursing facility population while other the measures listed focus on either UTIs associated with catheter use in ICU patients, catheter removal during day 1 or 2 for post-op acute care patients, or catheter use associated with urinary incontinence for home health patients.</p>	<p>3b</p> <p>C <input type="checkbox"/></p> <p>P <input type="checkbox"/></p> <p>M <input type="checkbox"/></p> <p>N <input type="checkbox"/></p> <p>NA <input type="checkbox"/></p>
<p>3c. Distinctive or Additive Value</p> <p>3c.1 Describe the distinctive, improved, or additive value this measure provides to existing NQF-endorsed measures: A new measure is needed based on the MDS 3.0 assessment specifications as it is replacing the current data source, the MDS 2.0. The measure has also been improved by excluding residents with medical conditions requiring catheterization (neurogenic bladder and obstructive uropathy).</p> <p>5.1 If this measure is similar to measure(s) already endorsed by NQF (i.e., on the same topic and the same target population), Describe why it is a more valid or efficient way to measure quality:</p>	<p>3c</p> <p>C <input type="checkbox"/></p> <p>P <input type="checkbox"/></p> <p>M <input type="checkbox"/></p> <p>N <input type="checkbox"/></p> <p>NA <input type="checkbox"/></p>
<p>TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for Usability?</p>	<p>3</p>

Comment [KP23]: 3b. The measure specifications are harmonized with other measures, and are applicable to multiple levels and settings.

Comment [k24]: 16 Measure harmonization refers to the standardization of specifications for similar measures on the same topic (e.g., influenza immunization of patients in hospitals or nursing homes), or related measures for the same target population (e.g., eye exam and HbA1c for patients with diabetes), or definitions applicable to many measures (e.g., age designation for children) so that they are uniform or compatible, unless differences are dictated by the evidence. The dimensions of harmonization can include numerator, denominator, exclusions, and data source and collection instructions. The extent of harmonization depends on the relationship of the measures, the evidence for the specific measure focus, and differences in data sources.

Comment [KP25]: 3c. Review of existing endorsed measures and measure sets demonstrates that the measure provides a distinctive or additive value to existing NQF-endorsed measures (e.g., provides a more complete picture of quality for a particular condition or aspect of healthcare, is a more valid or efficient way to measure).

Steering Committee: Overall, to what extent was the criterion, <i>Usability</i> , met? Rationale:		3 C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
4. FEASIBILITY		
Extent to which the required data are readily available, retrievable without undue burden, and can be implemented for performance measurement. (evaluation criteria)		Ev al Rat ing
4a. Data Generated as a Byproduct of Care Processes		4a C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
4a.1-2 How are the data elements that are needed to compute measure scores generated? Data generated as byproduct of care processes during care delivery (Data are generated and used by healthcare personnel during the provision of care, e.g., blood pressure, lab value, medical condition), Coding/abstraction performed by someone other than person obtaining original information (E.g., DRG, ICD-9 codes on claims, chart abstraction for quality measure or registry)		
4b. Electronic Sources		4b C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
4b.1 Are all the data elements available electronically? (elements that are needed to compute measure scores are in defined, computer-readable fields, e.g., electronic health record, electronic claims) No		
4b.2 If not, specify the near-term path to achieve electronic capture by most providers.		
4c. Exclusions		4c C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/>
4c.1 Do the specified exclusions require additional data sources beyond what is required for the numerator and denominator specifications? No		
4c.2 If yes, provide justification.		
4d. Susceptibility to Inaccuracies, Errors, or Unintended Consequences		4d C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
4d.1 Identify susceptibility to inaccuracies, errors, or unintended consequences of the measure and describe how these potential problems could be audited. If audited, provide results. A technical expert panel convened in January 2009 to make recommendations to retain, retire, or revise the quality measures as they transitioned from MDS 2.0 to MDS 3.0. TEP members recommended revising the measure to exclude residents with medical conditions requiring catheterization (neurogenic bladder and obstructive neuropathy).		
4e. Data Collection Strategy/Implementation		4e C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
4e.1 Describe what you have learned/modified as a result of testing and/or operational use of the measure regarding data collection, availability of data/missing data, timing/frequency of data collection, patient confidentiality, time/cost of data collection, other feasibility/ implementation issues: The data collection method is already in operational use, and there are no issues with these areas.		
4e.2 Costs to implement the measure (costs of data collection, fees associated with proprietary measures): Data are collected as part of an existing process with no additional cost.		

Comment [KP26]: 4a. For clinical measures, required data elements are routinely generated concurrent with and as a byproduct of care processes during care delivery. (e.g., BP recorded in the electronic record, not abstracted from the record later by other personnel; patient self-assessment tools, e.g., depression scale; lab values, meds, etc.)

Comment [KP27]: 4b. The required data elements are available in electronic sources. If the required data are not in existing electronic sources, a credible, near-term path to electronic collection by most providers is specified and clinical data elements are specified for transition to the electronic health record.

Comment [KP28]: 4c. Exclusions should not require additional data sources beyond what is required for scoring the measure (e.g., numerator and denominator) unless justified as supporting measure validity.

Comment [KP29]: 4d. Susceptibility to inaccuracies, errors, or unintended consequences and the ability to audit the data items to detect such problems are identified.

Comment [KP30]: 4e. Demonstration that the data collection strategy (e.g., source, timing, frequency, sampling, patient confidentiality, etc.) can be implemented (e.g., already in operational use, or testing demonstrates that it is ready to put into operational use).

<p>4e.3 Evidence for costs: This is not applicable.</p> <p>4e.4 Business case documentation: The proposed measure relies on data from the MDS 3.0. As there is no change in the data collection method for the MDS 3.0 as compared with its predecessor, the MDS 2.0, we do not anticipate any additional burden to nursing facilities. MDS 2.0, and soon to be MDS 3.0, data are collected as part of an existing, federally mandated process used for payment and quality monitoring purposes.</p>	
TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for <i>Feasibility</i>?	4
<p>Steering Committee: Overall, to what extent was the criterion, <i>Feasibility</i>, met? Rationale:</p>	<p>4 C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> A <input type="checkbox"/></p>
RECOMMENDATION	
(for NQF staff use) Check if measure is untested and only eligible for time-limited endorsement.	Time-limited <input type="checkbox"/>
<p>Steering Committee: Do you recommend for endorsement? Comments:</p>	<p>Y <input type="checkbox"/> N <input type="checkbox"/> A <input type="checkbox"/></p>
CONTACT INFORMATION	
<p>Co.1 Measure Steward (Intellectual Property Owner) Co.1 Organization Centers for Medicare & Medicaid Services, 7500 Security Boulevard, Mail Stop S3-02-01, Baltimore, Maryland, 21244-1850</p> <p>Co.2 Point of Contact Judith, Tobin, PT, MBA, Judith.Tobin@cms.hhs.gov, 410-786-6892-</p>	
<p>Measure Developer If different from Measure Steward Co.3 Organization RTI International, 1440 Main Street, Suite 310, Waltham, Massachusetts, 02451-1623</p> <p>Co.4 Point of Contact Robertta, Constantine, RN, MBA, PhD, rconstantine@rti.org, 781-434-1711-</p>	
<p>Co.5 Submitter If different from Measure Steward POC Robertta, Constantine, RN, MBA, PhD, rconstantine@rti.org, 781-434-1711-, RTI International</p>	
<p>Co.6 Additional organizations that sponsored/participated in measure development</p>	
ADDITIONAL INFORMATION	
<p>Workgroup/Expert Panel involved in measure development Ad.1 Provide a list of sponsoring organizations and workgroup/panel members' names and organizations. Describe the members' role in measure development. See Table 3: Nursing Home Quality Measures Technical Expert Panel (January 2009). This TEP met over 2 days in January 2009 to review the environmental scan of the current quality measures and make recommendations regarding their transition from MDS 2.0 to MDS 3.0.</p>	

<p>Ad.2 If adapted, provide name of original measure: This measure was adapted from the measure of the same name derived from MDS 2.0 data.</p> <p>Ad.3-5 If adapted, provide original specifications URL or attachment MedQIC resource manual - http://www.qualitynet.org/dcs/ContentServer?cid=1138050766910&pagename=Medqic%2FOtherResource%2FOtherResourcesTemplate&c=OtherResource</p>
<p>Measure Developer/Steward Updates and Ongoing Maintenance</p> <p>Ad.6 Year the measure was first released: 2002</p> <p>Ad.7 Month and Year of most recent revision: 02, 2010</p> <p>Ad.8 What is your frequency for review/update of this measure? Every 3 years</p> <p>Ad.9 When is the next scheduled review/update for this measure? 02, 2013</p>
<p>Ad.10 Copyright statement/disclaimers:</p>
<p>Ad.11 -13 Additional Information web page URL or attachment: Attachment Catheter tables_FINAL-634045010523392500.doc</p>
<p>Date of Submission (MM/DD/YY): 07/12/2010</p>

Project Name: NQF Nursing Home Project

Measure Title: Percent of Residents Who Have/Had a Catheter Inserted and Left in Their Bladder (Long Stay)

Planned Date of Measure Submission: March 19, 2010

Steward Name:

Point of Contact

Judith C. Tobin, PT, MBA
Centers for Medicare & Medicaid Services
7500 Security Boulevard
Mail Stop S3-02-01
Baltimore, MD 21244-1850
410-786-6892
Judith.Tobin@cms.hhs.gov

Developer/Submitter Name:

RTI International
Roberta Constantine, RN, MBA, PhD
1440 Main Street, Suite 310
Waltham, MA 02451-1623
781-434-1711
rconstantine@rti.org

Table 1. Measure Variability Across Facilities

Quality Measure (QM)	N of Facilities ¹	Std Mean	Std Dev	10 th Percentile	25 th Percentile	50 th Percentile	75 th Percentile	90 th Percentile	Facilities with QM = 0%
Indwelling Catheter	13,722	5.6%	4.0%	1.4%	2.9%	5.0%	7.7%	10.6%	6.2%

Table 2. Measure Trends Over Time

Quality Measure (QM)	Mean of Facility Triggering Rates (%)												
	Q3, 2003	Q4, 2003	Q1, 2004	Q2, 2004	Q3, 2004	Q4, 2004	Q1, 2005	Q2, 2005	Q3, 2005	Q4, 2005	Q1, 2006	Q2, 2006	Q3, 2006
Indwelling Catheter	5.7	5.8	6.1	5.9	5.9	5.9	6.3	6.1	5.9	5.8	6.0	5.8	5.6

Table 3. Nursing Home Quality Measures Technical Expert Panel (January 2009)

Name	Title	Affiliation
Barbara Anglin, RN	Program Services Consultant	American Association of Nurse Assessment Coordinators (AANAC)
Bonnie Burak-Danielson, MSM, EXP, LPTA	Rehab Manager of Reimbursement	Spaulding Rehab Network
Sarah Burger, MPH, RN	Senior Advisor and Coordinator	Coalition of Geriatric Nursing Organizations The John A. Hartford Institute for Geriatric Nursing
Diane Carter, MSN, RN, CS	President	AANAC
Kate Dennison, RN, RAC-MT	Minimum Data Set (MDS) Coordinator	The Cedars
Mary Ellard, RN, MPA/H, RAC-CT	Clinical Assessment Specialist	Five Star Quality Care, Inc.
Sandy Fitzler, RN	Senior Director of Clinical Services	American Health Care Association
David F. Hittle, PhD	Assistant Professor	Division of Health Care Policy and Research University of Colorado Denver, School of Medicine
Steve Levenson, MD, CMD	Multi-Facility Medical Director, Baltimore, MD	
Carol Maher, RN-BC, RAC-CT	Director of Clinical Reimbursement	Ensign Facilities Services
Barbara Manard, PhD	Vice President, Long Term Care/Health Strategies	American Association of Homes and Services for the Aging

Debra Saliba, MD, MPH	Anna and Harry Borun Chair in Geriatrics and Gerontology at UCLA Research Physician VA GLAHS GRECC Director of UCLA/JHA Borun Center for Gerontological Research Senior Natural Scientist RAND Health	University of California, Los Angeles (UCLA), Veterans Affairs (VA), RAND Corporation
Eric Tangalos, MD	Professor of Medicine	Mayo Clinic
Jacqueline Vance, RNC, CDONA/LTC	Director of Clinical Affairs	(American Medical Directors Association) AMDA
Mary Van de Kamp, MS/CCC-SLP	Vice President, Clinical Rehabilitation	Peoplefirst Rehabilitation
Charlene Harrington, PhD, RN, FAAN*	Professor Emeritus	University of California, San Francisco Fellow in the American Academy of Nursing

Measure #/Title/Steward

NH-020-10: Percent of Residents Who Have/Had a Catheter Inserted and Left in Their Bladder (Long Stay)
(Centers for Medicare & Medicaid Services)

Description: This measure updates CMS' current QM on catheter insertions. It is based on data from Minimum Data Set (MDS) 3.0 assessments of long-stay nursing home residents (those whose stay is longer than 100 days). This measure captures the percentage of long-stay residents who have had an indwelling catheter in the last 5 days noted on the most recent MDS 3.0 assessment, which may be annual, quarterly, significant change or significant correction during the selected quarter (3-month period). Long-stay residents are those residents who have been in nursing care at least 100 days. The measure is restricted to this population, which has long-term care needs, rather than the short stay population who are discharged within 100 days of admission.

Initial In-Person Vote:

Recommended for endorsement with conditions – 19
Not present - 1

<u>Steering Committee Questions/Conditions for Measure Developer:</u>	<u>Response from Measure Developer</u>
<ul style="list-style-type: none">• Do not exclude patients with missing data	<ul style="list-style-type: none">• The developer agreed to conduct an analysis of missing data as part of the validation analysis using the MDS 3.0 examining the extent of missing items, the impact of missing items on facility score and the impact of including vs. excluding patients with missing data on the facility score for the measure.
<ul style="list-style-type: none">• The definition of long-stay residents needs to be clarified	<ul style="list-style-type: none">• Long-stay residents are defined as those greater than 100 days. Residents who return to the nursing home following a hospital discharge will not have their stay reset to zero.