

NATIONAL QUALITY FORUM

National Voluntary Consensus Standards for Pediatric Cardiac Surgery Measures

Measure Number: PCS-001-09

Measure Title: Participation in a national database for pediatric and congenital heart surgery

Description: Participation in at least one multi-center, standardized data collection and feedback program that provides benchmarking of the physician’s data relative to national and regional programs and uses process and outcome measures. Participation is defined as submission of all congenital and pediatric operations performed to the database.

Numerator Statement: Whether or not there is participation in at least one multi-center data collection and feedback program.

Denominator Statement: N/A

Level of Analysis: Group of clinicians, Facility, Integrated delivery system, health plan, community/population

Data Source: Electronic Health/Medical Record, Electronic Clinical Database (The Society of Thoracic Surgeons Congenital Heart Surgery Database), Electronic Clinical Registry (The Society of Thoracic Surgeons Congenital Heart Surgery Database), Electronic Claims, Paper Medical Record

Measure Developer: Society of Thoracic Surgeons

Type of Endorsement: Recommended for Time-Limited Endorsement (Steering Committee Vote, Yes-9, No-0, Abstain-0)

Attachments: “STS Attachment: STS Procedure Code Definitions”

Meas# / Title/ (Owner)	Steering Committee Evaluation and Recommendation
PCS-001-09 Participation in a national database for pediatric and congenital heart surgery (Society of Thoracic Surgeons)	<p>Recommendation: Time-Limited Endorsement Yes-9; No-0; Abstain-0</p> <p>Final Measure Evaluation Ratings: I: Y-9; N-0 S: H-4; M-4; L-1 U: H-9; M-0; L-0 E: H-7; M-2; L-0</p> <p>Discussion: I: The Steering Committee agreed that this measure is important to measure and report. By reporting through a database, it is possible to identify potential quality issues and provide benchmarks. The measure identifies participation in a National Database for Pediatric and Congenital Heart Surgery. Participation in such a national database allows for program evaluation, as well as means for clinical improvement. Given the volume of pediatric surgeries performed, it is important to track them via database outcomes and collect feedback as to what types of interventions increased the likelihood of positive outcomes and also gives the ability to identify opportunities. Research has shown that</p>

NATIONAL QUALITY FORUM

National Voluntary Consensus Standards for Pediatric Cardiac Surgery Measures

Meas# / Title/ (Owner)	Steering Committee Evaluation and Recommendation
	<p>participation in multi-institutional databases/registries improves patient outcomes.</p> <p>S: The Steering Committee was divided in regards to the ratings for scientific acceptability; they ranged from high to moderate. Another similar measure was previously endorsed for adult cardiac surgery (NQF measure #0113).</p> <p>U: The Steering Committee agreed that this measure rated highly for usability. While the measure developers stated that this measure of participation in a database is not limited to the STS database, the STS database is used by a large number of programs already and retains over 90% of active programs in the United States.</p> <p>F: Members of the Steering Committee rated this measure highly overall for feasibility. It is easy to implement, as so many programs use this database and the mechanism for data procurement. Susceptibilities to errors have been addressed by the measure developers. Although this information is probably kept in the institutions, it is noted that it may be more difficult for smaller institutions to adhere to this measure. There were concerns raised regarding how submission of data to a registry would, as well as this being an expensive endeavor that requires high-level administrative commitment in order to implement. In response to a question about whether the measure requires participation in the STS database, the developer confirmed that the measure did not specify the STS registry.</p>

THE NATIONAL QUALITY FORUM

MEASURE SUBMISSION FORM VERSION 3.1

March 2009

The measure information you submit will be shared with NQF’s Steering Committees and Technical Advisory Panels to evaluate measures against the NQF criteria of importance to measure and report, scientific acceptability of measure properties, usability, and feasibility. Four conditions (as indicated below) must be met before proposed measures may be considered and evaluated for suitability as voluntary consensus standards. Not all acceptable measures will be strong—or equally strong—among each set of criteria. The assessment of each criterion is a matter of degree; however, all measures must be judged to have met the first criterion, *importance to measure and report*, in order to be evaluated against the remaining criteria. References to the specific measure evaluation criteria are provided in parentheses following the item numbers. Please refer to the *Measure Evaluation Criteria* for more information at www.qualityforum.org under Core Documents. Additional guidance is being developed and when available will be posted on the NQF website.

Use the tab or arrow (↓→) keys to move the cursor to the next field (or back ←↑). There are three types of response fields:

- drop-down menus - select one response;
- check boxes - check as many as apply; and
- text fields - you can copy and paste text into these fields or enter text; these fields are not limited in size, but in most cases, we ask that you summarize the requested information.

Please note that URL hyperlinks do not work in the form; you will need to type them into your web browser.

Be sure to answer all questions. Fields that are left blank will be interpreted as no or none. **Information must be provided in this form.** Attachments are not allowed except to provide additional detail or source documents for information that is summarized in this form. If you have important information that is not addressed by the questions, they can be entered into item #46 near the end of the form.

For questions about this form, please contact the NQF Project Director listed in the corresponding call for measures.

CONDITIONS FOR CONSIDERATION BY NQF	
	<i>Four conditions must be met before proposed measures may be considered and evaluated for suitability as voluntary consensus standards.</i>
A (A)	<i>Public domain or Measure Steward Agreement signed: Agreement signed and submitted (If no, do not submit) Template for the Measure Steward Agreement is available at www.qualityforum.org under Core Documents.</i>
B (B)	<i>Measure steward/maintenance: Is there an identified responsible entity and process to maintain and update the measure on a schedule commensurate with clinical innovation, but at least every 3 years? Yes, information provided in contact section (If no, do not submit)</i>
C (C)	<i>Intended use: Does the intended use of the measure include BOTH public reporting AND quality improvement? Yes (If no, do not submit)</i>
D (D)	<i>Fully developed and tested: Is the measure fully developed AND tested? No, testing will be completed within 24 months (If not tested and no plans for testing within 24 months, do not submit)</i>

THE NATIONAL QUALITY FORUM

MEASURE SUBMISSION FORM VERSION 3.1

March 2009

	(for NQF staff use) NQF Review #: PCS-001-09	NQF Project: Pediatric Cardiac Surgery
MEASURE SPECIFICATIONS & DESCRIPTIVE INFORMATION		
1	Information current as of (date- MM/DD/YY): 7/21/10	
2	Title of Measure: Participation in a National Database for Pediatric and Congenital Heart Surgery	
3	Brief description of measure ¹ : Participation in at least one multi-center, standardized data collection and feedback program that provides benchmarking of the physician's data relative to national and regional programs and uses process and outcome measures.	
4 (2a)	<p>Numerator Statement: Whether or not there is participation in at least one multi-center data collection and feedback program for pediatric and congenital heart surgery.</p> <p>Time Window: One year (12 months) and 4 years (48 months)</p> <p>Numerator Details (Definitions, codes with description): Participation is defined as submission of all congenital and pediatric operations performed to the database. Cardiac operations are defined as operations that are of operation types "CPB" or "No CPB Cardiovascular" (CPB is cardiopulmonary bypass.) [1].</p> <p>The following are STS procedure codes for pediatric and congenital cardiac operations per the STS Congenital Heart Surgery Database Version 3.0 Data Specifications. Analysis should include any index operation performed with any of the following component procedures on a patient with pediatric and/or congenital cardiac disease:</p> <p>10, 20, 30, 40, 2110, 50, 60, 70, 80, 85, 100, 110, 120, 130, 140, 150, 170, 180, 190, 2300, 2250, 2230, 210, 220, 230, 240, 2290, 250, 2220, 260, 270, 2120, 280, 2200, 290, 300, 310, 330, 340, 350, 360, 370, 380, 390, 400, 420, 430, 440, 450, 460, 2280, 465, 470, 480, 490, 500, 510, 520, 530, 540, 550, 570, 590, 2270, 600, 630, 640, 650, 610, 620, 1774, 1772, 580, 660, 2240, 2310, 2320, 670, 680, 690, 700, 715, 720, 730, 735, 740, 750, 760, 770, 780, 2100, 790, 800, 810, 820, 830, 2260, 840, 850, 860, 870, 880, 2160, 2170, 2180, 2140, 2150, 890, 900, 910, 920, 930, 940, 950, 960, 970, 980, 1000, 1010, 1025, 1030, 2340, 1035, 1050, 1060, 1070, 1080, 1090, 1110, 1120, 1123, 1125, 1130, 1140, 1145, 1150, 1160, 2190, 2210, 1180, 1200, 1210, 1220, 1230, 1240, 1250, 1260, 1275, 1280, 1285, 1290, 1291, 1300, 1310, 1320, 1330, 1340, 1360, 1365, 1370, 1380, 1390, 1410, 1450, 1460, 2350, 1470, 1480, 1490, 1500, 1590, 1600, 1610, 1630, 2095, 1640, 1650, 1660, 1670, 1680, 1690, 1700, 2330, 2130, 1720, 1730, 1740, 1760, 1780, 1790, 1802, 1804, 1830, 1860</p> <p>**Please find data definitions in STS Attachment 2 (of 2) - STS Procedure Code Definitions.</p> <p>Pediatric heart surgery is heart surgery on patients <18 years of age to treat congenital or acquired cardiac disease. Congenital heart surgery is heart surgery on patients of any age to treat congenital cardiac disease.</p> <p>Our measures apply to both pediatric heart surgery and congenital heart surgery, thus applying to the following operations:</p> <ol style="list-style-type: none"> 1. heart surgery on patients less than 18 years of age to treat congenital or acquired cardiac disease 2. heart surgery on patients of any age to treat congenital cardiac disease <p>References: 1. Jacobs JP, Mavroudis C, Jacobs ML, Maruszewski B, Tchervenkov CI, Lacour-Gayet FG, Clarke DR, Yeh T,</p>	

¹ Example of measure description: Percentage of adult patients with diabetes aged 18-75 years receiving one or more A1c test(s) per year.
 NQF Measure Submission Form, V3.1

	Walters HL 3rd, Kurosawa H, Stellin G, Ebels T, Elliott MJ. What is Operative Mortality? Defining Death in a Surgical Registry Database: A Report from the STS Congenital Database Task Force and the Joint EACTS-STC Congenital Database Committee. The Annals of Thoracic Surgery, 81(5):1937-41, May 2006.
5 (2a)	Denominator Statement: NA Time Window: Denominator Details (Definitions, codes with description):
6 (2a, 2d)	Denominator Exclusions: Measure Exclusions: Any operation that is not a pediatric or congenital Cardiac Operation. Cardiac operations are defined as operations that are of operation types of "CPB" or "No CPB Cardiovascular" (CPB is cardiopulmonary bypass.) [1]. 1. Jacobs JP, Mavroudis C, Jacobs ML, Maruszewski B, Tchervenkov CI, Lacour-Gayet FG, Clarke DR, Yeh T, Walters HL 3rd, Kurosawa H, Stellin G, Ebels T, Elliott MJ. What is Operative Mortality? Defining Death in a Surgical Registry Database: A Report from the STS Congenital Database Task Force and the Joint EACTS-STC Congenital Database Committee. The Annals of Thoracic Surgery, 81(5):1937-41, May 2006. Denominator Exclusion Details (Definitions, codes with description):
7 (2a, 2h)	Stratification Do the measure specifications require the results to be stratified? No ▶ If "other" describe: Identification of stratification variable(s): Stratification Details (Definitions, codes with description):
8 (2a, 2e)	Risk Adjustment Does the measure require risk adjustment to account for differences in patient severity before the onset of care? No ▶ If yes, (select one) ▶ Is there a separate proprietary owner of the risk model? (select one) Identify Risk Adjustment Variables: Detailed risk model: attached <input type="checkbox"/> OR Web page URL:
9 (2a)	Type of Score: Dichotomous Calculation Algorithm: attached <input type="checkbox"/> OR Web page URL: Interpretation of Score (<i>Classifies interpretation of score according to whether better quality is associated with a higher score, a lower score, a score falling within a defined interval, or a passing score</i>) Other ▶ If "Other", please describe: Answer "yes" indicates higher score. Just monitoring this measure should improve quality.
10 (2a, 4a, 4b)	Identify the required data elements(e.g., primary diagnosis, lab values, vital signs): operation Data dictionary/code table attached <input type="checkbox"/> OR Web page URL: Data Quality (2a) <i>Check all that apply</i> <input checked="" type="checkbox"/> Data are captured from an authoritative/accurate source (e.g., lab values from laboratory personnel) <input checked="" type="checkbox"/> Data are coded using recognized data standards <input checked="" type="checkbox"/> Method of capturing data electronically fits the workflow of the authoritative source <input checked="" type="checkbox"/> Data are available in EHRs <input checked="" type="checkbox"/> Data are auditable
11 (2a, 4b)	Data Source and Data Collection Methods <i>Identifies the data source(s) necessary to implement the measure specifications. Check all that apply</i> <input checked="" type="checkbox"/> Electronic Health/Medical Record <input checked="" type="checkbox"/> Paper Medical Record <input checked="" type="checkbox"/> Electronic Clinical Database, Name: The Society of Thoracic Surgeons Congenital Heart Surgery Database <input type="checkbox"/> Standardized clinical instrument, Name: <input type="checkbox"/> Standardized patient survey, Name: <input type="checkbox"/> Standardized clinician survey, Name: <input checked="" type="checkbox"/> Electronic Clinical Registry, Name: The Society <input type="checkbox"/> Other, Describe:

	<p>of Thoracic Surgeons Congenital Heart Surgery Database</p> <p>Instrument/survey attached <input type="checkbox"/> OR Web page URL:</p> <p><input checked="" type="checkbox"/> Electronic Claims</p> <p><input type="checkbox"/> Electronic Pharmacy data</p> <p><input type="checkbox"/> Electronic Lab data</p> <p><input type="checkbox"/> Electronic source - other, Describe:</p>
12 (2a)	<p>Sampling <i>If measure is based on a sample, provide instructions and guidance on sample size.</i></p> <p>Minimum sample size:</p> <p>Instructions:</p>
13 (2a)	<p>Type of Measure: Structure ▶ If "Other", please describe:</p> <p>▶ If part of a composite or paired with another measure, please identify composite or paired measure</p>
14 (2a)	<p>Unit of Measurement/Analysis (<i>Who or what is being measured</i>) <i>Check all that apply.</i></p> <p><input type="checkbox"/> Can be measured at all levels</p> <p><input type="checkbox"/> Individual clinician (e.g., physician, nurse)</p> <p><input checked="" type="checkbox"/> Group of clinicians (e.g., facility department/unit, group practice)</p> <p><input checked="" type="checkbox"/> Facility (e.g., hospital, nursing home)</p> <p><input checked="" type="checkbox"/> Integrated delivery system</p> <p><input checked="" type="checkbox"/> Health plan</p> <p><input checked="" type="checkbox"/> Community/Population</p> <p><input type="checkbox"/> Other (<i>Please describe</i>):</p>
15 (2a)	<p>Applicable Care Settings <i>Check all that apply</i></p> <p><input type="checkbox"/> Can be used in all healthcare settings</p> <p><input type="checkbox"/> Ambulatory Care (office/clinic)</p> <p><input type="checkbox"/> Behavioral Healthcare</p> <p><input type="checkbox"/> Community Healthcare</p> <p><input type="checkbox"/> Dialysis Facility</p> <p><input type="checkbox"/> Emergency Department</p> <p><input type="checkbox"/> EMS emergency medical services</p> <p><input type="checkbox"/> Health Plan</p> <p><input type="checkbox"/> Home Health</p> <p><input type="checkbox"/> Hospice</p> <p><input checked="" type="checkbox"/> Hospital</p> <p><input type="checkbox"/> Long term acute care hospital</p> <p><input type="checkbox"/> Nursing home/ Skilled Nursing Facility (SNF)</p> <p><input type="checkbox"/> Prescription Drug Plan</p> <p><input type="checkbox"/> Rehabilitation Facility</p> <p><input type="checkbox"/> Substance Use Treatment Program/Center</p> <p><input type="checkbox"/> Other (<i>Please describe</i>):</p>
<p>IMPORTANCE TO MEASURE AND REPORT</p>	
<p>Note: This is a threshold criterion. If a measure is not judged to be sufficiently important to measure and report, it will not be evaluated against the remaining criteria.</p>	
16 (1a)	<p>Is measure related to a National Priority Partners priority area? Safety reliability</p> <p>(for NQF staff use) Does measure address a <u>specific</u> NPP goal? (www.qualityforum.org/about/NPP/): Safety</p>
17 (1a)	<p>Does the measure address a high impact aspect of healthcare high resource use</p> <p>Summary of Evidence: Congenital heart disease is a common birth defect that affects approximately 1 in 125 live births [1]. Pediatric and Congenital Heart Surgery is a subspecialty of high resource utilization that has the potential to repair or palliate the majority of patients with pediatric and congenital cardiac disease.</p> <p>Citations² for Evidence: 1. Tchervenkov CI, Jacobs JP, Bernier P-L, Stellin G, Kurosawa H, Mavroudis C, Jonas RA, Cicek SM, Al-Halees Z, J. Elliott MJ, Jatene MB, Kinsley RH, Kreutzer C, Leon-Wyss J, Liu J, Maruszewski B, Nunn GR, Ramirez-Marroquin S, Sandoval N, Sano S, Sarris GE, Sharma R, Shoeb A, Spray TL, Ungerleider RM, Yangni-Angate H, Ziemer G. The improvement of care for paediatric and congenital cardiac disease across the World: a challenge for the World Society for Pediatric and Congenital Heart Surgery. In: 2008 Supplement to Cardiology in the Young: Databases and The Assessment of Complications associated with The Treatment of Patients with Congenital Cardiac Disease, Prepared by: The Multi-Societal Database Committee for Pediatric and Congenital Heart Disease, Jeffrey P. Jacobs, MD (editor). Cardiology in the Young, Volume</p>


² Citations can include, but are not limited to journal articles, reports, web pages (URLs).
NQF Measure Submission Form, V3.1

	<p>18, Issue S2 (Suppl. 2), pp 63-69, December 9, 2008.</p>
18	<p>Opportunity for Improvement <i>Provide evidence that demonstrates considerable variation, or overall poor performance, across providers.</i></p> <p>(1b) Summary of Evidence: The Report of the 2005 STS Congenital Heart Surgery Practice and Manpower Survey, undertaken by the STS Workforce on Congenital Heart Surgery, documented that 122 centers in the United States of America perform pediatric and congenital heart surgery [1]. The Spring 2009 STS Congenital Heart Surgery Database Report [2, 3] contains data from 75 of these 122 centers from the United States of America. Therefore, the number of centers that are involved in a systemic multi-institutional database continues to improve however over 35% of institutions still performing congenital heart surgery do not participate in the STS Congenital Heart Surgery Database the largest and most comprehensive database for congenital heart centers in the US. Therefore, there remains significant variability in this structure measurement.</p> <p>Citations for Evidence: 1. Jacobs ML, Mavroudis C, Jacobs JP, Tchervenkov CI, Pelletier GJ. Report of the 2005 STS Congenital Heart Surgery Practice and Manpower Survey: A Report from The STS Work Force on Congenital Heart Surgery. The Annals of Thoracic Surgery, 82(3):1152-8, 1159e1-5; discussion 1158-9, September 2006.</p> <p>2. www.sts.org</p> <p>3. Jacobs JP, Jacobs ML, Mavroudis C, Lacour-Gayet FG, Tchervenkov CI. Executive Summary: The Society of Thoracic Surgeons Congenital Heart Surgery Database - Tenth Harvest - (January 1, 2005 - December 31, 2008). The Society of Thoracic Surgeons (STS) and Duke Clinical Research Institute (DCRI), Duke University Medical Center, Durham, North Carolina, United States, Spring 2009 Harvest.</p> <p>4. Jacobs JP, Jacobs ML, Mavroudis C, Backer CL, Lacour-Gayet FG, Tchervenkov CI, Franklin RCG, Béland MJ, Jenkins KJ, Walters III H, Bacha EA, Maruszewski B, Kurosawa H, Clarke DR, Gaynor JW, Spray TL, Stellin G, Ebels T, Krogmann ON, Aiello VD, Colan SD, Weinberg P, Giroud JM, Everett A, Wernovsky G, Martin J. Elliott MJ, Edwards FH. Nomenclature and databases for the surgical treatment of congenital cardiac disease - an updated primer and an analysis of opportunities for improvement. In: 2008 Supplement to Cardiology in the Young: Databases and The Assessment of Complications associated with The Treatment of Patients with Congenital Cardiac Disease, Prepared by: The Multi-Societal Database Committee for Pediatric and Congenital Heart Disease, Jeffrey P. Jacobs, MD (editor). Cardiology in the Young, Volume 18, Issue S2 (Suppl. 2), pp 38-62, December 9, 2008.</p>
19	<p>Disparities <i>Provide evidence that demonstrates disparity in care/outcomes related to the measure focus among populations.</i></p> <p>(1b) Summary of Evidence: Children undergoing congenital heart surgery are cared for by many specialties (i.e. anesthesia, critical care, cardiology, surgery). The major organizations (i.e. The Congenital Cardiac Anesthesia Society, Pediatric Cardiac Intensive Care Society, and The American College of Cardiology) from each of these specialties have written their opinions in a manuscript why creation and participation in a national database is essential to improve the quality of care and outcomes of children with congenital heart disease.</p> <p>No formal testing of disparities has been done. Disparities and trends could be tested for many of these metrics using the STS Database.</p> <p>Citations for evidence: 1. Jacobs JP, Wernovsky G, Elliott MJ. Analysis of outcomes for congenital cardiac disease: can we do better? <i>Cardiol Young</i>. 2007 Sep;17 Suppl 2:145-58.</p> <p>2. LaRovere JM, Jeffries HE, Sachdeva RC, Rice TB, Wetzell RC, Cooper DS, Bird GL, Ghanayem NS, Checchia PA, Chang AC, Wessel DL. Databases for assessing the outcomes of the treatment of patients with congenital and paediatric cardiac disease--the perspective of critical care. <i>Cardiol Young</i>. 2008 Dec;18 Suppl 2:130-6.</p> <p>3. Jacobs ML, Jacobs JP, Franklin RC, Mavroudis C, Lacour-Gayet F, Tchervenkov CI, Walters H, Bacha EA, Clarke DR, William Gaynor J, Spray TL, Stellin G, Ebels T, Maruszewski B, Tobota Z, Kurosawa H, Elliott M. Databases for assessing the outcomes of the treatment of patients with congenital and paediatric cardiac disease--the perspective of cardiac surgery. <i>Cardiol Young</i>. 2008 Dec;18 Suppl 2:101-15.</p>

	<p>4. Jenkins KJ, Beekman Iii RH, Bergersen LJ, Everett AD, Forbes TJ, Franklin RC, Klitzner TS, Krogman ON, Martin GR, Webb CL. Databases for assessing the outcomes of the treatment of patients with congenital and paediatric cardiac disease--the perspective of cardiology. <i>Cardiol Young</i>. 2008 Dec;18 Suppl 2:116-23.</p>						
<p>20 (1c)</p>	<p>If measuring an Outcome Describe relevance to the national health goal/priority, condition, population, and/or care being addressed:</p> <p>If not measuring an outcome, provide evidence supporting this measure topic and grade the strength of the evidence <i>Summarize the evidence (including citations to source) supporting the focus of the measure as follows:</i></p> <ul style="list-style-type: none"> • Intermediate outcome - evidence that the measured intermediate outcome (e.g., blood pressure, Hba1c) leads to improved health/avoidance of harm or cost/benefit. • Process - evidence that the measured clinical or administrative process leads to improved health/avoidance of harm and if the measure focus is on one step in a multi-step care process, it measures the step that has the greatest effect on improving the specified desired outcome(s). • Structure - evidence that the measured structure supports the consistent delivery of effective processes or access that lead to improved health/avoidance of harm or cost/benefit. • Patient experience - evidence that an association exists between the measure of patient experience of health care and the outcomes, values and preferences of individuals/ the public. • Access - evidence that an association exists between access to a health service and the outcomes of, or experience with, care. • Efficiency- demonstration of an association between the measured resource use and level of performance with respect to one or more of the other five IOM aims of quality. <p>Type of Evidence Check all that apply</p> <table border="0"> <tr> <td><input type="checkbox"/> Evidence-based guideline</td> <td><input checked="" type="checkbox"/> Quantitative research studies</td> </tr> <tr> <td><input type="checkbox"/> Meta-analysis</td> <td><input checked="" type="checkbox"/> Qualitative research studies</td> </tr> <tr> <td><input type="checkbox"/> Systematic synthesis of research</td> <td><input type="checkbox"/> Other (<i>Please describe</i>):</td> </tr> </table> <p>Overall Grade for Strength of the Evidence³ (<i>Use the USPSTF system, or if different, also describe how it relates to the USPSTF system</i>):</p> <p>Summary of Evidence (<i>provide guideline information below</i>): This measure is a Structure Measure. The importance of how participation in a national database is essential for the optimization of patient care, the institution of government programs or policies, and the economics around the care of congenital heart disease has been extensively expressed by multiple specialties taking care of these children. Over 50 manuscript have been written using the STS congenital heart surgery database that have helped create risk stratification mechanisms and other standards and tools that have improved the care in children with Congenital Heart Disease.</p> <p>Citations for Evidence: 1. Jacobs JP, Wernovsky G, Elliott MJ. Analysis of outcomes for congenital cardiac disease: can we do better? <i>Cardiol Young</i>. 2007 Sep;17 Suppl 2:145-58.</p> <p>2. LaRovere JM, Jeffries HE, Sachdeva RC, Rice TB, Wetzell RC, Cooper DS, Bird GL, Ghanayem NS, Checchia PA, Chang AC, Wessel DL. Databases for assessing the outcomes of the treatment of patients with congenital and paediatric cardiac disease--the perspective of critical care. <i>Cardiol Young</i>. 2008 Dec;18 Suppl 2:130-6.</p> <p>3. Jacobs ML, Jacobs JP, Franklin RC, Mavroudis C, Lacour-Gayet F, Tchervenkov CI, Walters H, Bacha EA,</p>	<input type="checkbox"/> Evidence-based guideline	<input checked="" type="checkbox"/> Quantitative research studies	<input type="checkbox"/> Meta-analysis	<input checked="" type="checkbox"/> Qualitative research studies	<input type="checkbox"/> Systematic synthesis of research	<input type="checkbox"/> Other (<i>Please describe</i>):
<input type="checkbox"/> Evidence-based guideline	<input checked="" type="checkbox"/> Quantitative research studies						
<input type="checkbox"/> Meta-analysis	<input checked="" type="checkbox"/> Qualitative research studies						
<input type="checkbox"/> Systematic synthesis of research	<input type="checkbox"/> Other (<i>Please describe</i>):						

³The strength of the body of evidence for the specific measure focus should be systematically assessed and rated, e.g., USPSTF grading system www.ahrq.gov/clinic/uspstmeth.htm: A - The USPSTF recommends the service. There is high certainty that the net benefit is substantial. B - The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial. C - The USPSTF recommends against routinely providing the service. There may be considerations that support providing the service in an individual patient. There is at least moderate certainty that the net benefit is small. Offer or provide this service only if other considerations support the offering or providing the service in an individual patient. D - The USPSTF recommends against the service. There is moderate or high certainty that the service has no net benefit or that the harms outweigh the benefits. I - The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined.

	<p>Clarke DR, William Gaynor J, Spray TL, Stellin G, Ebels T, Maruszewski B, Tobota Z, Kurosawa H, Elliott M. Databases for assessing the outcomes of the treatment of patients with congenital and paediatric cardiac disease--the perspective of cardiac surgery. <i>Cardiol Young</i>. 2008 Dec;18 Suppl 2:101-15.</p> <p>4. Jenkins KJ, Beekman lii RH, Bergersen LJ, Everett AD, Forbes TJ, Franklin RC, Klitzner TS, Krogman ON, Martin GR, Webb CL. Databases for assessing the outcomes of the treatment of patients with congenital and paediatric cardiac disease--the perspective of cardiology. <i>Cardiol Young</i>. 2008 Dec;18 Suppl 2:116-23.</p> <p>5. Welke KF, O'Brien SM, Peterson ED, Ungerleider RM, Jacobs ML, Jacobs JP. The complex relationship between pediatric cardiac surgical case volumes and mortality rates in a aational clinical database. <i>The Journal of Thoracic and Cardiovascular Surgery</i>, . 2009 May;137(5):1133-40. Epub 2009 Mar 17, PMID: 19379979, May, 2009.</p> <p>6. Sachdeva RC. Electronic healthcare data collection and pay-for-performance: translating theory into practice. <i>Ann Health Law</i>. 2007 Summer;16(2):291-311</p>
<p>21 (1c)</p>	<p>Clinical Practice Guideline <i>Cite the guideline reference; quote the specific guideline recommendation related to the measure and the guideline author's assessment of the strength of the evidence; and summarize the rationale for using this guideline over others.</i></p> <p>Guideline Citation: At the current time no uniform practice guidelines are in place for pediatric and congenital cardiac surgery. Clinical care rationale mainly depends on the consensus of a panel of experts in the field. In lieu of guideline support for the measures, published consensus opinion and supporting clinical data from the STS Congenital Heart Surgery Database will be used.</p> <p>Specific guideline recommendation:</p> <p>Guideline author's rating of strength of evidence <i>(If different from USPSTF, also describe it and how it relates to USPSTF):</i></p> <p>Rationale for using this guideline over others:</p>
<p>22 (1c)</p>	<p>Controversy/Contradictory Evidence <i>Summarize any areas of controversy, contradictory evidence, or contradictory guidelines and provide citations.</i></p> <p>Summary: The only controversy that arises regarding the participation in a multi-institutional database has centered around cost in dollars and effort to collect the data and submit it. However, the argument that one should not collect their outcomes and understand their own institution's result in comparison to national standards seems to be below the acceptable norms for caring and operating on children with congenital heart disease as stated in multiple manuscripts and by our panel of experts.</p> <p>Citations: 1. Jacobs JP, Wernovsky G, Elliott MJ. Analysis of outcomes for congenital cardiac disease: can we do better? <i>Cardiol Young</i>. 2007 Sep;17 Suppl 2:145-58.</p> <p>2. LaRovere JM, Jeffries HE, Sachdeva RC, Rice TB, Wetzel RC, Cooper DS, Bird GL, Ghanayem NS, Checchia PA, Chang AC, Wessel DL. Databases for assessing the outcomes of the treatment of patients with congenital and paediatric cardiac disease--the perspective of critical care. <i>Cardiol Young</i>. 2008 Dec;18 Suppl 2:130-6.</p> <p>3. Jacobs ML, Jacobs JP, Franklin RC, Mavroudis C, Lacour-Gayet F, Tchervenkov CI, Walters H, Bacha EA, Clarke DR, William Gaynor J, Spray TL, Stellin G, Ebels T, Maruszewski B, Tobota Z, Kurosawa H, Elliott M. Databases for assessing the outcomes of the treatment of patients with congenital and paediatric cardiac disease--the perspective of cardiac surgery. <i>Cardiol Young</i>. 2008 Dec;18 Suppl 2:101-15.</p> <p>4. Jenkins KJ, Beekman lii RH, Bergersen LJ, Everett AD, Forbes TJ, Franklin RC, Klitzner TS, Krogman ON, Martin GR, Webb CL. Databases for assessing the outcomes of the treatment of patients with congenital and paediatric cardiac disease--the perspective of cardiology. <i>Cardiol Young</i>. 2008 Dec;18 Suppl 2:116-23.</p>
<p>23 (1)</p>	<p>Briefly describe how this measure (as specified) will facilitate significant gains in healthcare quality related to the specific priority goals and quality problems identified above: The decision for a center to participate in a multi-institutional database states their commitment to patient</p>

	<p>care because they are helping to provide the infrastructure, spanning geographical and subspecialty boundaries, for collaboration between health care professionals interested in the analysis of outcomes of treatments provided to patients with congenital cardiac disease, with the ultimate aim of improvement in the quality of care provided to these patients. The participation of the center regardless of their size makes the database stronger and a more accurate representation of congenital heart surgery so that the processes and patient care recommendations based on the database are useful to the participating programs.</p> <p>Over the past decade, mortality after pediatric cardiac surgery has been declining and currently stands at 4%. This declining mortality and the limited number of pediatric cardiac surgical cases have made the assessment of mortality, as a single measure to evaluate the quality of care, less meaningful. The assessment of morbidity has become more important [1].</p> <p>In order to properly assess morbidity, one must be able to track performance using a resource such as a data registry that enables health providers to get a firm grasp on the details of pediatric and congenital cardiac surgery performed at a center over both 1 year and 4 year time intervals.</p> <p>1. Jacobs JP, Jacobs ML, et. al. What is Operative Morbidity? Defining Complications in a Surgical Registry Database. <i>Annals of Thoracic Surgery</i> 2007;87:1416-1421.</p>
SCIENTIFIC ACCEPTABILITY OF MEASURE PROPERTIES	
<p>Note: Testing and results should be summarized in this form. However, additional detail and reports may be submitted as supplemental information or provided as a web page URL. If a measure has not been tested, it is only potentially eligible for time-limited endorsement.</p>	
24	<p>Supplemental Testing Information: attached <input type="checkbox"/> OR Web page URL: </p>
25 (2b)	<p>Reliability Testing</p> <p>Data/sample: In 2007, The Society of Thoracic Surgeons (STS) Task Force to develop National Quality Forum (NQF) Indicators for Pediatric and Congenital Cardiac Surgery held a series of focused phone conferences involving multiple STS Congenital and Pediatric Heart Surgery leaders. This generated a proposal for National Quality Forum Indicators for Pediatric and Congenital Cardiac Surgery.</p> <p>The attached proposal has been approved unanimously by the STS Task Force to develop National Quality Forum Indicators for Pediatric and Congenital Cardiac Surgery. The proposal has been reviewed and approved by the following 6 STS Committees:</p> <ol style="list-style-type: none"> 1. The STS Task Force to develop NQF Indicators for Pediatric and Congenital Cardiac Surgery 2. The STS Congenital Database Task Force 3. The STS Congenital Heart Surgery Work Force 4. The STS National Database Work Force 5. The STS Council on Quality, Research & Patient Safety 6. The STS Executive Committee <p>Formal reliability/validity testing on these metrics has not ever been done. In fact, formal reliability/validity testing has never been done on any quality metrics for pediatric and congenital heart surgery.</p> <p>Analytic Method:</p> <p>Testing Results:</p>
26 (2c)	<p>Validity Testing</p> <p>Data/sample: The measure validity was confirmed by The Society of Thoracic Surgeons (STS) Task Force to develop National Quality Forum (NQF) Indicators for Pediatric and Congenital Cardiac Surgery after a comprehensive literature review. The measure validity was confirmed further by:</p> <ul style="list-style-type: none"> - The STS Congenital Database Task Force

	<ul style="list-style-type: none"> - The STS Congenital Heart Surgery Work Force - The STS National Database Work Force - The STS Council on Quality, Research & Patient Safety - The STS Executive Committee <p>Formal reliability/validity testing on these metrics has not ever been done. In fact, formal reliability/validity testing has never been done on any quality metrics for pediatric and congenital heart surgery.</p> <p>Analytic Method:</p> <p>Testing Results:</p>
<p>27 (2d)</p>	<p>Measure Exclusions <i>Provide evidence to justify exclusion(s) and analysis of impact on measure results during testing.</i></p> <p>Summary of Evidence supporting exclusion(s): We will exclude any operation that is not a pediatric or congenital Cardiac Operation. Cardiac operations are defined as operations that are of operation types of "CPB" or "No CPB Cardiovascular" (CPB is cardiopulmonary bypass.) [1]. Excluding centers not performing congenital heart surgery will not impact a multi-institutional congenital heart surgery database.</p> <p>Furthermore, it is important to understand that the Society of Thoracic Surgeons advocates utilization of a systematic multi-institutional clinical database (registry) for the analysis of cardiac surgical outcomes and the assessment of quality. Evidence from three recent investigations suggests that the validity of coding of lesions seen in the congenitally malformed heart via the International Classification of Diseases as used in Administrative Databases is likely to be poor[2, 3, 4]. First, in a series of 373 infants with congenital cardiac defects at Children’s Hospital of Wisconsin, investigators report that only 52% of the cardiac diagnoses in the medical records had a corresponding code from the International Classification of Diseases in the hospital discharge database [2]. Second, the Hennepin County Medical Center discharge database in Minnesota identified all infants born during 2001 with a code for congenital cardiac disease using the International Classification of Diseases. A review of these 66 medical records by physicians was able to confirm only 41% of the codes contained in the administrative database from the International Classification of Diseases [3]. Third, the Metropolitan Atlanta Congenital Defect Program of the Birth Defect Branch of the Centers for Disease Control and Prevention of the federal government of the United States of America carried out surveillance of infants and fetuses with cardiac defects delivered to mothers residing in Atlanta during the years 1988 through 2003 [4]. These records were reviewed and classified using both administrative coding and the clinical nomenclature used in the Society of Thoracic Surgeons Congenital Heart Surgery Database. This study concluded that analyses based on the codes available in the International Classification of Diseases are likely to “have substantial misclassification” of congenital cardiac disease.</p> <p>Several potential reasons can explain the poor diagnostic accuracy of Administrative Databases and codes from the International Classification of Diseases:</p> <ol style="list-style-type: none"> 1) accidental miscoding 2) coding performed by medical records clerks who have never seen the actual patient 3) contradictory or poorly described information in the medical record 4) lack of diagnostic specificity for congenital cardiac disease in the codes of the of International Classification of Diseases 5) inadequately trained medical coders <p>Citations for Evidence: 1. Jacobs JP, Mavroudis C, Jacobs ML, Maruszewski B, Tchervenkov CI, Lacour-Gayet FG, Clarke DR, Yeh T, Walters HL 3rd, Kurosawa H, Stellin G, Ebels T, Elliott MJ. What is Operative Mortality? Defining Death in a Surgical Registry Database: A Report from the STS Congenital Database Task Force and the Joint EACTS-STS Congenital Database Committee. The Annals of Thoracic Surgery, 81(5):1937-41, May 2006.</p> <p>2. Cronk CE, Malloy ME, Pelech AN, et al. Completeness of state administrative databases for surveillance of congenital heart disease. Birth Defects Res A Clin Mol Teratol 2003; 67: 597-603.</p> <p>3. Frohnert BK, Lussky RC, Alms MA, Mendelsohn NJ, Symonik DM, Falken MC. Validity of hospital discharge</p>

	<p>data for identifying infants with cardiac defects. J Perinatol 2005; 25: 737-742.</p> <p>4. Strickland MJ, Riehle-Colarusso TJ, Jacobs JP, Reller MD, Mahle WT, Botto LD, Tolbert PE, Jacobs ML, Lacour-Gayet FG, Tchervenkov CI, Mavroudis C, Correa A. The importance of nomenclature for congenital cardiac disease: implications for research and evaluation. In: 2008 Supplement to Cardiology in the Young: Databases and The Assessment of Complications associated with The Treatment of Patients with Congenital Cardiac Disease, Prepared by: The Multi-Societal Database Committee for Pediatric and Congenital Heart Disease, Jeffrey P. Jacobs, MD (editor). Cardiology in the Young, Volume 18, Issue S2 (Suppl. 2), pp 92-100, December 9, 2008</p> <p>Data/sample:</p> <p>Analytic Method:</p> <p>Testing Results:</p>
28 (2e)	<p>Risk Adjustment Testing Summarize the testing used to determine the need (or no need) for risk adjustment and the statistical performance of the risk adjustment method.</p> <p>Data/sample: None</p> <p>Analytic Method:</p> <p>Testing Results:</p> <p>► If outcome or resource use measure not risk adjusted, provide rationale:</p>
29 (2g)	<p>Testing comparability of results when more than 1 data method is specified (e.g., administrative claims or chart abstraction)</p> <p>Data/sample: Clinical data abstraction is the only method utilized</p> <p>Analytic Method:</p> <p>Results:</p>
30 (2f)	<p>Provide Measure Results from Testing or Current Use Results from current use</p> <p>Data/sample: The STS Congenital Heart Surgery Database</p> <p>Methods to identify statistically significant and practically/meaningfully differences in performance: Outliers can be identified with 95% confidence intervals based on the sample size, with complexity stratification for one and four-year time intervals</p> <p>Results: Currently being collected in the STS Congenital Heart Surgery Database. We know that 82 out of 122 pediatric heart surgery centers in the USA participate in the STS Congenital Heart Surgery Database.</p>
31 (2h)	<p>Identification of Disparities</p> <p>► If measure is stratified by factors related to disparities (i.e. race/ethnicity, primary language, gender, SES, health literacy), provide stratified results: NA</p> <p>► If disparities have been reported/identified, but measure is not specified to detect disparities, provide rationale:</p>
USABILITY	
32 (3)	<p>Current Use In use If in use, how widely used Nationally ► If "other," please describe: The STS Congenital Heart Surgery Database</p> <p><input type="checkbox"/> Used in a public reporting initiative, name of initiative: Sample report attached <input type="checkbox"/> OR Web page URL:</p>

<p>33 (3a)</p>	<p>Testing of Interpretability (<i>Testing that demonstrates the results are understood by the potential users for public reporting and quality improvement</i>)</p> <p>Data/sample: Post-operative mortality and morbidity data are currently being collected voluntarily by The Society of Thoracic Surgeons Congenital Cardiac Surgery Database. This measure is one of the nine structure measures that is being evaluated for NQF endorsement. All of the outcome metrics are used by clinicians as performance feedback and are tracked in the STS Database. No focused consumer testing has been done to date on any of these metrics. No public reporting has been done on any of these metrics to date. Pediatric and congenital heart surgery is very different from adult heart surgery. Separate metrics are necessary.</p> <p>Methods:</p> <p>Results:</p>
<p>34 (3b), (3c)</p>	<p>Relation to other NQF-endorsed™ measures</p> <p>► Is this measure similar or related to measure(s) already endorsed by NQF (on the same topic or the same target population)? <i>Measures can be found at www.qualityforum.org under Core Documents. Check all that apply</i></p> <p><input type="checkbox"/> Have not looked at other NQF measures <input checked="" type="checkbox"/> Other measure(s) on same topic <input type="checkbox"/> Other measure(s) for same target population <input type="checkbox"/> No similar or related measures</p> <p>Name and number of similar or related NQF-endorsed™ measure(s): <i>Similar or Related Measure - NQF # 0113 - Participation in a Systematic Database for Cardiac Surgery</i> Description: <i>Does the facility participate in a multicenter data collection and feedback program that provides benchmarking relative to peers and uses process and outcome measures?</i></p> <p>Rationale: NQF # 0113 states: "Participation in a Systematic Database for Cardiac Surgery"</p> <p><i>Our new proposed measure is different because it specifies participation in a National Database for Pediatric and Congenital Heart Surgery. Participation in a Systematic Database for Cardiac Surgery is not enough because for this group of patients, quality is consistent with participation in a cardiac surgery database specific for pediatric and congenital cardiac surgery.</i></p> <p>Are the measure specifications harmonized with existing NQF-endorsed™ measures? <i>Yes, fully harmonized</i> ► If not fully harmonized, provide rationale:</p> <p>Describe the distinctive, improved, or additive value this measure provides to existing NQF-endorsed measures:</p>
<p>FEASIBILITY</p>	
<p>35 (4a)</p>	<p>How are the required data elements generated? <i>Check all that apply</i></p> <p><input checked="" type="checkbox"/> <i>Data elements are generated concurrent with and as a byproduct of care processes during care delivery (e.g., blood pressure or other assessment recorded by personnel conducting the assessment)</i></p> <p><input type="checkbox"/> <i>Data elements are generated from a patient survey (e.g., CAHPS)</i></p> <p><input type="checkbox"/> <i>Data elements are generated through coding performed by someone other than the person who obtained the original information (e.g., DRG or ICD-9 coding on claims)</i></p> <p><input checked="" type="checkbox"/> <i>Other, Please describe: The Society of Thoracic Surgeons Congenital Heart Surgery Database participation agreement</i></p>
<p>36 (4b)</p>	<p>Electronic Sources <i>All data elements</i></p> <p>► <i>If all data elements are not in electronic sources, specify the near-term path to electronic collection by most providers:</i></p> <p>► <i>Specify the data elements for the electronic health record: Surgery</i></p>
<p>37</p>	<p>Do the specified exclusions require additional data sources beyond what is required for the other</p>

<p>(4c))</p>	<p>specifications? <i>No</i></p> <p>► <i>If yes, provide justification:</i></p>
<p>38 (4d))</p>	<p><i>Identify susceptibility to inaccuracies, errors, or unintended consequences of the measure:</i></p> <p><u><i>Inaccuracies and Errors:</i></u> <i>This measure may be susceptible to human error (i.e., recording the measure inaccurately or not recording the measure at all)</i></p> <p><u><i>Unintended Consequences:</i></u> <i>One should be cautious in drawing conclusions from the observation of these measures, especially in circumstances where there is a declining morbidity and mortality [1,2].</i></p> <p><i>1. Welke KF, Karamlou T, Ungerleider RM, Diggs BS. Mortality is not a valid indicator of quality differences between pediatric cardiac surgery programs. Ann Thoracic Surgery. (in press)</i></p> <p><i>2. O'Brien SM, Gauvreau K. Statistical issues in the analysis and interpretation of outcomes for congenital cardiac surgery. In: 2008 Cardiology in the Young Supplement: Databases and The Assessment of Complications associated with The Treatment of Patients with Congenital Cardiac Disease, Prepared by: The Multi-Societal Database Committee for Pediatric and Congenital Heart Disease, Jeffrey P. Jacobs, MD (editor). Cardiology in the Young. 2008;18(Suppl.2):145-151.</i></p> <p><i>Describe how could these potential problems be audited:</i></p> <p><u><i>Inaccuracies and Errors:</i></u> <i>Each participant is responsible for the quality and accuracy of the data they submit to the database. Each participant agrees to the following quality control measures in the participation agreement:</i></p> <p><i>i) "Participant hereby warrants that all data submitted for inclusion in the CHS Database will be accurate and complete, and acknowledges that such data may be subject to independent audit. Participant will use its best efforts to address any data or related deficiencies identified by the independent data warehouse service provider, and agrees to cooperate with and assist STS and its designees in connection with the performance of any independent audit.</i></p> <p><i>ii) Participant warrants that it will take all reasonable steps to avoid the submission of duplicative data for inclusion in the CHS Database, including but not limited to apprising the Director of the STS National Database and the independent data warehouse service provider about any other Participation Agreements in which an individual cardiothoracic surgeon named above or on Schedule A attached hereto (as amended from time to time) is also named."</i></p> <p><i>In addition, the Data warehouse and analysis center at Duke Clinical Research Institute, performs a series of internal quality controls on the submitted data and issues an annual data quality report</i></p> <p><u><i>Unintended Consequences:</i></u> <i>The Society of Thoracic Surgeons Database audit process is used. In addition, outliers can be identified with 95% confidence intervals based on the sample size with complexity stratification for one and four-year time intervals</i></p> <p><i>Did you audit for these potential problems during testing? Yes If yes, provide results:</i> <i>Clarke DR, Breen LS, Jacobs ML, Franklin RCG, Tobota Z, Maruszewski B, Jacobs JP. Verification of data in congenital cardiac surgery. In: 2008 Cardiology in the Young Supplement: Databases and The Assessment of Complications associated with The Treatment of Patients with Congenital Cardiac Disease, Prepared by: The Multi-Societal Database Committee for Pediatric and Congenital Heart Disease, Jeffrey P. Jacobs, MD (editor). Cardiology in the Young. 2008;18(Suppl. 2):177-187.</i></p>
<p>39</p>	<p><i>Testing feasibility Describe what have you learned/modified as a result of testing and/or operational use of the measure regarding data collection, availability of data/missing data, timing/frequency of data</i></p>

(4e) *collection, patient confidentiality, time/cost of data collection, other feasibility/ implementation issues:*
Lessons Learned:

- *The STS CHS database collects gender, race/ethnicity, age and geographic location information, so disparities and trends can be studied for populations at risk.*
- *Data elements required for the measure can be captured and the measure is actionable by the physician.*
- *There are no data availability issues.*
- *Cost to collect the data includes staff training and the use of specific software. However there are no additional costs over what a provider would pay to be a part of the STS CHS Database or other registry that collects this information.*
- *This measure can be used in a variety of care settings and at different levels of analysis (i.e. physician, hospital, etc.)*
- *Formal reliability testing was not done. Instead, the participant is bound by the participation agreement and his/her participation can be monitored by observing the data submitted on an annual basis.*
- *There are no confidentiality concerns. The data is de-identified, and the sites must be HIPAA compliant and obtain IRB approval for use of the database.*
- *The STS Congenital Quality Measures Sub-Committee meets at the STS Annual Meeting. The Subcommittee will review each STS congenital cardiac surgery measure on a yearly basis. Changes or updates to the measure will be at the recommendation of the committee.*
- *The STS has a yearly meeting (The Advances in Quality and Outcomes Conference) devoted to the Database for the clinicians and data coordinators.*
- *The audit process has demonstrated that data is very complete and accurate.1*

1. Clarke DR, Breen LS, Jacobs ML, Franklin RCG, Tobota Z, Maruszewski B, Jacobs JP. Verification of data in congenital cardiac surgery. In: 2008 Cardiology in the Young Supplement: Databases and The Assessment of Complications associated with The Treatment of Patients with Congenital Cardiac Disease, Prepared by: The Multi-Societal Database Committee for Pediatric and Congenital Heart Disease, Jeffrey P. Jacobs, MD (editor). Cardiology in the Young. 2008;18(Suppl. 2):177-187.

CONTACT INFORMATION

40 **Web Page URL for Measure Information** Describe where users (implementers) should go for more details on specifications of measures, or assistance in implementing the measure.
 Web page URL: www.sts.org

41 **Measure Steward Point of Contact**
 First Name: [Jane](#) MI: [M](#) Last Name: [Han](#) Credentials (MD, MPH, etc.): [MSW](#)
 Organization: [The Society of Thoracic Surgeons](#)
 Street Address: [633 N. Saint Clair St, Suite 2320](#) City: [Chicago](#) State: [IL](#) ZIP: [60611](#)
 Email: jhan@sts.org Telephone: [312-202-5856](#) ext:

42 **Measure Developer Point of Contact** **If different from Measure Steward**
 First Name: MI: Last Name: Credentials (MD, MPH, etc.):
 Organization:
 Street Address: City: State: ZIP:
 Email: Telephone: ext:

ADDITIONAL INFORMATION

43 **Workgroup/Expert Panel involved in measure development** [Workgroup/panel used](#)
 ► If workgroup used, describe the members' role in measure development: [The STS Task Force to Develop NQF Indicators for Pediatric and Congenital Cardiac Surgery members collectively formulated the numerator statement and defined its parameters in addition to identifying data elements and sources of data.](#)
 ► Provide a list of workgroup/panel members' names and organizations:

Name	Institution	Location
Erle H. Austin, III, MD	University of Louisville Kosair Children's Hospital	Louisville, KY

	Emile A. Bacha, MD	Children's Hospital Boston Department of Cardiovascular Surgery	Boston, MA
	Pedro J. del Nido, MD	Children's Hospital Boston Department of Cardiac Surgery	Boston, MA
	Charles D. Fraser, Jr., MD	Texas Children's Hospital Division of Congenital Heart Surgery	Houston, TX
	Frederick L. Grover, MD	University of Colorado Health Sciences Center Department of Surgery	Aurora, CO
	Jennifer C. Hirsch, MD	University of Michigan Health System Section of Cardiac Surgery	Ann Arbor, MI
	Jeffrey P. Jacobs, MD	The Congenital Heart Institute of Florida	Saint Petersburg, FL
	Marshall L. Jacobs, MD	Cleveland Clinic Center for Pediatric and Congenital Heart Diseases	Cleveland, OH
	David L. Morales, MD	Texas Children's Hospital Division of Congenital Heart Surgery	Houston, TX
	Kamal K. Pourmoghadam, MD	Geisinger Medical Center Department of Pediatric Cardiac Surgery	Danville, PA
	Jeffrey B. Rich, MD	Mid-Atlantic Cardiothoracic Surgeons, Ltd.	Norfolk, VA
	James S. Tweddell, MD	Children's Hospital of Wisconsin Department of Cardiothoracic Surgery	Milwaukee, WI
44	<p><i>Measure Developer/Steward Updates and Ongoing Maintenance</i> <i>Year the measure was first released: NA</i> <i>Month and Year of most recent revision: NA</i> <i>What is the frequency for review/update of this measure? Once a year at STS Annual Meeting</i> <i>When is the next scheduled review/update for this measure? January 2010</i></p>		
45	Copyright statement/disclaimers:		
46	Additional Information:		
47	I have checked that the submission is complete and any blank fields indicate that no information is provided. <input checked="" type="checkbox"/>		
48	Date of Submission (MM/DD/YY): 8/31/09		

Attachment

The Society of Thoracic Surgeons (STS) Congenital Heart Surgery Database

The Report of the 2005 STS Congenital Heart Surgery Practice and Manpower Survey, undertaken by the STS Workforce on Congenital Heart Surgery, documented that 122 centers in the United States of America perform pediatric and congenital heart surgery [1]. The Spring 2009 STS Congenital Heart Surgery Database Report [2, 3] contains data from 75 of these 122 centers from the United States of America.

The Society of Thoracic Surgeons Congenital Heart Surgery Database is the largest clinical database in North America dealing with congenital cardiac malformations [www.sts.org] [2, 3, 4]. It has grown annually since its inception, both in terms of the number of participating centers submitting data, and the number of operations analyzed (Figures 1, 2, and 3). The Spring 2009 STS Congenital Heart Surgery Database Report includes 87,271 operations performed in 76 centers in North America, 75 from the United States of America and 1 from Canada. One Japanese centre also submits data; however, these Japanese data are not included in the aggregate report produced by the STS.

The STS Congenital Heart Surgery Database has standardized methodology for meaningful multi-institutional analyses of the outcomes of patients undergoing surgery to treat pediatric and congenital heart disease, based on standardization of the following six elements:

1. use of a common language and nomenclature,
2. use of an established uniform core dataset for collection of information,
3. incorporation of a mechanism of evaluating case complexity,
4. availability of a mechanism to assure and verify the completeness and accuracy of the data collected,
5. collaboration between medical and surgical subspecialties, and
6. standardized protocols for life-long follow-up.

The STS Congenital Heart Surgery Database stratifies participating centers into low volume, medium volume, and high volume sites. The Spring 2009 STS Congenital Heart Surgery Database Report [2, 3] contains data from 75 sites in the United States of America with broad geographical representation and a wide variation on program size. In the Spring 2009 STS Congenital Heart Surgery Database Report, of the 75 sites, 13 were low volume, 25 were medium volume, and 37 were high volume.

FIGURE 1

The graph documents the annual growth in the STS Congenital Heart Surgery Database by number of participating centres submitting data. The aggregate report from Spring 2009 Harvest of the STS Congenital Heart Surgery Database includes data from 76 Congenital Heart Surgery Centres from the United States of America and Canada. One Japanese centre also submits data; however, these Japanese data are not included in the aggregate report produced by the STS.

Growth in the STS Congenital Heart Surgery Database





FIGURE 2

The graph documents the annual growth in the STS Congenital Heart Surgery Database (CHSD) by the number of operations per averaged 4 year data collection cycle. The aggregate report from the Spring 2009 Harvest of the STS CHSD included 87,271 operations performed in the 4 year period of January 1, 2005, through December 31, 2008, submitted from 76 centres from North America, 75 from the U.S. and 1 from Canada. One Japanese centre also submits data; however, these Japanese data are not included in the aggregate report produced by the STS.

Growth in the STS Congenital Heart Surgery Database




Operations per averaged 4 year data collection cycle.

FIGURE 3

The graph documents the annual growth in the STS Congenital Heart Surgery Database (CHSD) by the cumulative number of operations over time. The current number of cumulative total operations in the STS CHSD is 134,690. The aggregate report from the Spring 2009 Harvest of the STS CHSD included 87, 271 operations performed in the 4 year period of January 1, 2005, through December 31, 2008, submitted from 76 centres from North America, 75 from the U.S. and 1 from Canada. One Japanese centre also submits data; however, these Japanese data are not included in the aggregate report produced by the STS.

Growth in the STS Congenital Heart Surgery Database



Cumulative operations over time.

References

1. Jacobs ML, Mavroudis C, Jacobs JP, Tchervenkov CI, Pelletier GJ. Report of the 2005 STS Congenital Heart Surgery Practice and Manpower Survey: A Report from The STS Work Force on Congenital Heart Surgery. *The Annals of Thoracic Surgery*, 82(3):1152-8, 1159e1-5; discussion 1158-9, September 2006.
2. www.sts.org
3. Jacobs JP, Jacobs ML, Mavroudis C, Lacour-Gayet FG, Tchervenkov CI. Executive Summary: The Society of Thoracic Surgeons Congenital Heart Surgery Database - Tenth Harvest - (January 1, 2005 - December 31, 2008). The Society of Thoracic Surgeons (STS) and Duke Clinical Research Institute (DCRI), Duke University Medical Center, Durham, North Carolina, United States, Spring 2009 Harvest.
4. Jacobs JP, Jacobs ML, Mavroudis C, Backer CL, Lacour-Gayet FG, Tchervenkov CI, Franklin RCG, Béland MJ, Jenkins KJ, Walters III H, Bacha EA, Maruszewski B, Kurosawa H, Clarke DR, Gaynor JW, Spray TL, Stellin G, Ebels T, Krogmann ON, Aiello VD, Colan SD, Weinberg P, Giroud JM, Everett A, Wernovsky G, Martin J. Elliott MJ, Edwards FH. Nomenclature and databases for the surgical treatment of congenital cardiac disease - an updated primer and an analysis of opportunities for improvement. In: 2008 Supplement to *Cardiology in the Young: Databases and The Assessment of Complications associated with The Treatment of Patients with Congenital Cardiac Disease*, Prepared by: The Multi-Societal Database Committee for Pediatric and Congenital Heart Disease, Jeffrey P. Jacobs, MD (editor). *Cardiology in the Young*, Volume 18, Issue S2 (Suppl. 2), pp 38-62, December 9, 2008.

Procedure Code	Description	Definition
10	PFO, Primary closure	Suture closure of patent foramen ovale (PFO)
20	ASD repair, Primary closure	Suture closure of secundum (most frequently), coronary sinus, sinus venosus or common atrium ASD.
30	ASD repair, Patch	Patch closure (using any type of patch material) of secundum, coronary sinus, or sinus venosus ASD.
40	ASD repair, Device	Closure of any type ASD (including PFO) using a device
2110	ASD repair, Patch + PAPVC repair	
50	ASD, Common atrium (single atrium), Septation	Septation of common (single) atrium using any type patch material.
60	ASD creation/enlargement	Creation of an atrial septal defect or enlargement of an existing atrial septal defect using a variety of modalities including balloon septostomy, blade septostomy, or surgical septectomy. Creation may be accomplished with or without use of cardiopulmonary bypass.
70	ASD partial closure	Intentional partial closure of any type ASD (partial suture or fenestrated patch closure).
80	Atrial septal fenestration	Creation of a fenestration (window) in the septum between the atrial chambers. Usually performed using a hole punch, creating a specifically sized communication in patch material placed on the atrial septum.
85	Atrial fenestration closure	Closure of previously created atrial fenestration using any method including device, primary suture, or patch.
100	VSD repair, Primary closure	Suture closure of any type VSD.
110	VSD repair, Patch	Patch closure (using any type of patch material) of any type VSD.
120	VSD repair, Device	Closure of any type VSD using a device.
130	VSD, Multiple, Repair	Closure of more than one VSD using any method or combination of methods. Further information regarding each type of VSD closed and method of closure can be provided by additionally listing specifics for each VSD closed. In the case of multiple VSDs in which only one is closed the procedure should be coded as closure of a single VSD. The fundamental diagnosis, in this case, would be "VSD, Multiple" and a secondary diagnosis can be the morphological type of VSD that was closed at the time of surgery.
140	VSD creation/enlargement	Creation of a ventricular septal defect or enlargement of an existing ventricular septal defect.
150	Ventricular septal fenestration	Creation of a fenestration (window) in the septum between the ventricular chambers. Usually performed using a hole punch, creating a specifically sized communication in patch material placed on the ventricular septum.
170	AVC (AVSD) repair, Complete (CAVSD)	Repair of complete AV canal (AVSD) using one- or two patch or other technique, with or without mitral valve cleft repair.
180	AVC (AVSD) repair, Intermediate (Transitional)	Repair of intermediate AV canal (AVSD) using ASD and VSD patch, or ASD patch and VSD suture, or other technique, with or without mitral valve cleft repair.
190	AVC (AVSD) repair, Partial (Incomplete) (PAVSD)	Repair of partial AV canal defect (primum ASD), any technique, with or without repair of cleft mitral valve.
2300	Valvuloplasty, Common atrioventricular valve	

2250	Valvuloplasty converted to valve replacement in the same operation, Common atrioventricular valve	
2230	Valve replacement, Common atrioventricular valve	
210	AP window repair	Repair of AP window using one- or two-patch technique with cardiopulmonary bypass; or, without cardiopulmonary bypass, using transcatheter device or surgical closure.
220	Pulmonary artery origin from ascending aorta (hemitruncus) repair	Repair of pulmonary artery origin from the ascending aorta by direct reimplantation, autogenous flap, or conduit, with or without use of cardiopulmonary bypass.
230	Truncus arteriosus repair	Truncus arteriosus repair that most frequently includes patch VSD closure and placement of a conduit from RV to PA. In some cases, a conduit is not placed but an RV to PA connection is made by direct association. Very rarely, there is no VSD to be closed. Truncal valve repair or replacement should be coded separately (Valvuloplasty, Truncal valve; Valve replacement, Truncal valve), as would be the case as well with associated arch anomalies requiring repair (e.g., Interrupted aortic arch repair).
240	Valvuloplasty, Truncal valve	Truncal valve repair, any type.
2290	Valvuloplasty converted to valve replacement in the same operation, Truncal valve	
250	Valve replacement, Truncal valve	Replacement of the truncal valve with a prosthetic valve.
2220	Truncus + Interrupted aortic arch repair (IAA) repair	
260	PAPVC repair	PAPVC repair revolves around whether an intracardiac baffle is created to redirect pulmonary venous return to the left atrium or if the anomalous pulmonary vein is translocated and
270	PAPVC, Scimitar, Repair	In scimitar syndrome, PAPVC repair also revolves around whether an intracardiac baffle is created to redirect pulmonary venous return to the left atrium or if the anomalous pulmonary vein is translocated and connected to the left atrium directly. If there is an associated ASD and it is closed, that procedure should also be listed. Occasionally an
2120	PAPVC repair, Baffle redirection to left atrium with systemic vein translocation (Warden) (SVC sewn to right atrial appendage)	
280	TAPVC repair	Repair of TAPVC, any type. Issues surrounding TAPVC repair involve how the main pulmonary venous confluence anastomosis is fashioned, whether an associated ASD is closed or left open or enlarged (ASD closure and
2200	TAPVC repair + Shunt - systemic-to-pulmonary	

290	Cor triatriatum repair	Repair of cor triatriatum. Surgical decision making revolves around the approach to the membrane creating the cor triatriatum defect, how any associated ASD is closed, and how any associated anomalous pulmonary vein connection is addressed. Both ASD closure and anomalous pulmonary venous connection may be listed as separate procedures.
300	Pulmonary venous stenosis repair	Repair of pulmonary venous stenosis, whether congenital or acquired. Repair can be accomplished with a variety of approaches: sutureless, patch venoplasty, stent placement, etc.
310	Atrial baffle procedure (non-Mustard, non-Senning)	The atrial baffle procedure code is used primarily for repair of systemic venous anomalies, as in redirection of left superior vena cava drainage to the right atrium.
330	Anomalous systemic venous connection repair	With the exception of atrial baffle procedures (harvest code 310), anomalous systemic venous connection repair includes a range of surgical approaches, including, among others: ligation of anomalous vessels, reimplantation of anomalous vessels (with or without use of a conduit), or redirection of anomalous systemic venous flow through directly to the pulmonary circulation (bidirectional Glenn to redirect LSVC or RSVC to left or right pulmonary artery, respectively).
340	Systemic venous stenosis repair	Stenosis or obstruction of a systemic vein (most commonly SVC or IVC) may be relieved with patch or
350	TOF repair, No ventriculotomy	Tetralogy of Fallot repair (assumes VSD closure and relief of pulmonary stenosis at one or more levels), without use of an incision in the infundibulum of the right ventricle for exposure. In most cases this would be a transatrial and transpulmonary artery approach to repair the VSD and relieve the pulmonary stenosis. If the main pulmonary artery incision is extended proximally through the pulmonary annulus, this must be considered "transannular" and thus a ventricular incision, though the length of the incision onto the ventricle itself may be minimal.
360	TOF repair, Ventriculotomy, Nontransannular patch	Tetralogy of Fallot repair (assumes VSD closure and relief of pulmonary stenosis at one or more levels), with use of a ventriculotomy incision, but without placement of a transpulmonary annulus patch. If the main pulmonary artery incision is extended proximally through the pulmonary annulus, this must be considered "transannular" and thus a ventricular incision, though the length of the incision onto the ventricle itself may be minimal.
370	TOF repair, Ventriculotomy, Transannular patch	Tetralogy of Fallot repair (assumes VSD closure and relief of pulmonary stenosis at one or more levels), with use of a ventriculotomy incision and placement of a transpulmonary annulus patch. If the main pulmonary artery incision is extended proximally through the pulmonary annulus, this must be considered "transannular" and thus a ventricular incision, though the length of the incision onto the ventricle itself may be minimal.

380	TOF repair, RV-PA conduit	Tetralogy of Fallot repair (assumes VSD closure and relief of pulmonary stenosis at one or more levels), with placement of a right ventricle-to-pulmonary artery conduit. In this procedure the major components of pulmonary stenosis are relieved with placement of the RV-PA conduit.
390	TOF - AVC (AVSD) repair	Tetralogy of Fallot repair (assumes VSD closure and relief of pulmonary stenosis at one or more levels), with repair of associated AV canal defect. Repair of associated atrial septal defect or atrioventricular valve repair(s) should be listed as additional or secondary procedures under the primary TOF-AVC procedure.
400	TOF - Absent pulmonary valve repair	Repair of tetralogy of Fallot with absent pulmonary valve complex. In most cases this repair will involve pulmonary valve replacement (pulmonary or aortic homograft, porcine, other) and reduction pulmonary artery arterioplasty.
420	Pulmonary atresia - VSD (including TOF, PA) repair	For patients with pulmonary atresia with ventricular septal defect without MAPCAs, including those with tetralogy of Fallot with pulmonary atresia, repair may entail either a tetralogy-like repair with transannular patch placement, a VSD closure with placement of an RV-PA conduit, or an intraventricular tunnel VSD closure with transannular patch or RV-PA conduit placement. To assure an accurate count of repairs of pulmonary atresia-VSD without MAPCAs, even if a tetralogy-type repair or Rastelli-type repair is used, the pulmonary atresia-VSD code should be the code used, not Rastelli procedure or tetralogy of Fallot repair with transannular patch.
430	Pulmonary atresia - VSD - MAPCA (pseudotruncus) repair	In the presence of MAPCAs, this code implies implies pulmonary unifocalization (multi- or single-stage), repair of VSD (may be intraventricular tunnel or flat patch VSD closure), and placement of an RV-PA conduit.
440	Unifocalization MAPCA(s)	Anastomosis of aortopulmonary collateral arteries into the left, right, or main pulmonary artery or into a tube graft or other
450	Occlusion MAPCA(s)	Occlusion, or closing off, of MAPCAs. This may be done with a transcatheter occluding device, usually a
460	Valvuloplasty, Tricuspid	Reconstruction of the tricuspid valve may include but not be limited to a wide range of techniques including: leaflet patch extension, artificial chordae placement, papillary muscle translocation with or without detachment. Annuloplasty techniques that may be done solely or in combination with leaflet, chordae or muscle repair to achieve a competent valve include: eccentric annuloplasty, Kay annular plication, purse-
2280	Valvuloplasty converted to valve replacement in the same operation, Tricuspid	

465	Ebstein's repair	To assure an accurate count of repairs of Ebstein's anomaly of the tricuspid valve, this procedure code was included. Repair of Ebstein's anomaly may include, among other techniques, repositioning of the tricuspid valve, plication of the atrialized right ventricle, or right reduction atrioplasty. Often associated ASD's may be closed and arrhythmias addressed with surgical ablation procedures. These procedures should be entered as separate procedure codes.
470	Valve replacement, Tricuspid (TVR)	Replacement of the tricuspid valve with a prosthetic valve.
480	Valve closure, Tricuspid (exclusion, univentricular approach)	In a functional single ventricle heart, the tricuspid valve may be closed using a patch, thereby excluding the RV. Tricuspid valve closure may be used for infants with Ebstein's anomaly and
490	Valve excision, Tricuspid (without replacement)	Excision of the tricuspid valve without placement of a valve prosthesis.
500	Valve surgery, Other, Tricuspid	Other tricuspid valve surgery not specified in procedure codes.
510	RVOT procedure	Included in this procedural code would be all RVOT procedures not elsewhere specified in the nomenclature system. These might be, among others: resection of subvalvar pulmonary stenosis (not DCRV type; may be localized fibrous diaphragm or high infundibular stenosis), right ventricular patch augmentation, or reduction pulmonary artery arterioplasty.
520	1 1/2 ventricular repair	Partial biventricular repair; includes intracardiac repair with bidirectional cavopulmonary anastomosis to volume unload a small ventricle or poorly functioning ventricle
530	PA, reconstruction (plasty), Main (trunk)	Reconstruction of the main pulmonary artery trunk commonly using patch material. If balloon angioplasty is performed or a stent is placed in the main pulmonary artery intraoperatively, this code may be used in addition to the balloon dilation or stent placement code. If MPA reconstruction is performed with PA debanding, both codes should be listed.
540	PA, reconstruction (plasty), Branch, Central (within the hilar bifurcation)	Reconstruction of the right or left branch (or both right and left) pulmonary arteries (within the hilar bifurcation) commonly using patch material. If balloon angioplasty is performed or a stent is placed in the right or left (or both) pulmonary artery intraoperatively, this code may be used in addition to the balloon dilation or stent placement code. If, rarely, branch PA
550	PA, reconstruction (plasty), Branch, Peripheral (at or beyond the hilar bifurcation)	Reconstruction of the peripheral right or left branch (or both right and left) pulmonary arteries (at or beyond the hilar bifurcation) commonly using patch material. If balloon angioplasty is performed or a stent is placed in the right or left (or both) peripheral pulmonary artery intraoperatively, this code may be used in addition to the balloon dilation or stent placement code.

570	DCRV repair	Surgical repair of DCRV combines relief of the low infundibular stenosis (via muscle resection) and closure of a VSD when present. A ventriculotomy may be required and is repaired by patch enlargement of the infundibulum. VSD closure and patch enlargement of the infundibulum, if done, should be listed as separate procedure codes.
590	Valvuloplasty, Pulmonic	Valvuloplasty of the pulmonic valve may include a range of techniques including but not limited to: valvotomy with or without bypass, commissurotomy, and valvuloplasty.
2270	Valvuloplasty converted to valve replacement in the same operation, Pulmonic	
600	Valve replacement, Pulmonic (PVR)	Replacement of the pulmonic valve with a prosthetic valve. Care must be taken to differentiate between homograft pulmonic valve replacement and placement of a homograft RV-PA conduit.
630	Valve excision, Pulmonary (without replacement)	Excision of the pulmonary valve without placement of a valve prosthesis.
640	Valve closure, Semilunar	Closure of a semilunar valve (pulmonic or aortic) by any technique.
650	Valve surgery, Other, Pulmonic	Other pulmonic valve surgery not specified in procedure codes.
610	Conduit placement, RV to PA	Placement of a conduit, any type, from RV to PA.
620	Conduit placement, LV to PA	Placement of a conduit, any type, from LV to PA.
1774	Conduit placement, Ventricle to aorta	Placement of a conduit from the right or left ventricle to the aorta.
1772	Conduit placement, Other	Placement of a conduit from any chamber or vessel to any vessel, valved or valveless, not listed elsewhere.
580	Conduit reoperation	Conduit reoperation is the code to be used in the event of conduit failure, in whatever position (LV to aorta, LV to PA, RA to RV, RV to aorta, RV to PA, etc.), and from whatever cause (somatic growth, stenosis, insufficiency, infection, etc).
660	Valvuloplasty, Aortic	Valvuloplasty of the aortic valve for stenosis and/or insufficiency including, but not limited to the following techniques: valvotomy (open or closed), commissurotomy, aortic valve suspension, leaflet (left, right or noncoronary) partial resection, reduction, or leaflet shaving, extended valvuloplasty (freeing of leaflets, commissurotomy, and extension of leaflets using autologous or bovine pericardium), or annuloplasty (partial - interrupted or noncircumferential sutures, or complete - circumferential sutures).
2240	Valvuloplasty converted to valve replacement in the same operation, Aortic	
2310	Valvuloplasty converted to valve replacement in the same operation, Aortic – with Ross procedure	

2320	Valvuloplasty converted to valve replacement in the same operation, Aortic – with Ross-Konno procedure	
670	Valve replacement, Aortic (AVR)	Replacement of the aortic valve with a prosthetic valve (mechanical, bioprosthetic, or homograft). Use this code only if type of valve prosthesis is unknown or does not fit into the specific valve replacement codes available. Autograft valve replacement should be coded as a Ross procedure.
680	Valve replacement, Aortic (AVR), Mechanical	Replacement of the aortic valve with a mechanical prosthetic valve.
690	Valve replacement, Aortic (AVR), Bioprosthetic	Replacement of the aortic valve with a bioprosthetic prosthetic valve.
700	Valve replacement, Aortic (AVR), Homograft	Replacement of the aortic valve with a homograft prosthetic valve.
715	Aortic root replacement, Bioprosthetic	Replacement of the aortic root (that portion of the aorta attached to the heart; it gives rise to the coronary arteries) with a bioprosthesis (e.g., porcine) in a conduit, often composite.
720	Aortic root replacement, Mechanical	Replacement of the aortic root (that portion of the aorta attached to the heart; it gives rise to the coronary arteries) with a mechanical prosthesis in a composite conduit.
730	Aortic root replacement, Homograft	Replacement of the aortic root (that portion of the aorta attached to the heart; it gives rise to the coronary arteries) with a homograft.
735	Aortic root replacement, Valve sparing	Replacement of the aortic root (that portion of the aorta attached to the heart; it gives rise to the coronary arteries) without replacing the aortic valve (using a tube graft).
740	Ross procedure	Replacement of the aortic valve with a pulmonary autograft and replacement of the pulmonary valve with a homograft conduit.
750	Konno procedure	Relief of left ventricular outflow tract obstruction associated with aortic annular hypoplasia, aortic valvar stenosis and/or aortic valvar insufficiency via Konno aortoventriculoplasty. Components of the surgery include a longitudinal incision in the aortic septum, a vertical incision in the outflow tract of the right ventricle to join the septal incision, aortic valve replacement, and patch reconstruction of the outflow tracts of both ventricles.
760	Ross-Konno procedure	Relief of left ventricular outflow tract obstruction associated with aortic annular hypoplasia, aortic valvar stenosis and/or aortic valvar insufficiency via Konno aortoventriculoplasty using a pulmonary autograft root for the aortic root replacement.
770	Other annular enlargement procedure	Techniques included under this procedure code include those designed to effect aortic annular enlargement that are not included in other procedure codes. These include the Manouagian and Nicks aortic annular enlargement procedures.

780	Aortic stenosis, Subvalvar, Repair	Subvalvar aortic stenosis repair by a range of techniques including excision, excision and myotomy, excision and myomectomy, myotomy, myomectomy, initial placement of apical-aortic conduit (LV to aorta conduit replacement would be coded as conduit reoperation), Vouhé aortoventriculoplasty (aortic annular incision at commissure of left and right coronary cusps is carried down to the septum and RV infundibulum; septal muscle is resected, incisions are closed, and the aortic annulus is reconstituted), or other aortoventriculoplasty techniques.
2100	Aortic stenosis, Subvalvar, Repair, With myectomy for IHSS	
790	Aortic stenosis, Supravalvar, Repair	Repair of supravalvar aortic stenosis involving all techniques of patch aortoplasty and aortoplasty involving the use of all autologous tissue. In simple patch aortoplasty a diamond-shaped patch may be used, in the Doty technique an extended patch is placed (Y shaped patch, incision carried into two sinuses), and in the Brom repair the ascending aorta is transected, any fibrous ridge is resected, and the three sinuses are patched separately.
800	Valve surgery, Other, Aortic	Other aortic valve surgery not specified in other procedure codes.
810	Sinus of Valsalva, Aneurysm repair	Sinus of Valsalva aneurysm repair can be organized by site of aneurysm (left, right or noncoronary sinus), type of repair (suture, patch graft, or root repair by tube graft or valved conduit), and approach used (from chamber of origin (aorta) or from chamber of penetration (LV, RV, PA, left or right atrium, etc.). Aortic root replacement procedures in association with sinus of Valsalva aneurysm repairs are usually for associated uncorrectable aortic insufficiency or multiple sinus involvement and the aortic root replacement procedure should also be listed. Additional procedures also performed at the time of sinus of Valsalva aneurysm repair include but are not limited to VSD closure, repair or replacement of aortic valve, and coronary reconstruction; these procedures should also be coded separately from the sinus of Valsalva aneurysm repair.
820	LV to aorta tunnel repair	LV to aorta tunnel repair can be accomplished by suture, patch, or both, and may require reimplantation of the right coronary artery. Associated coronary artery procedures should be coded separately from the LV to aorta tunnel repair.
830	Valvuloplasty, Mitral	Repair of mitral valve including, but not limited to: valvotomy (closed or open heart), cleft repair, annuloplasty with or without ring, chordal reconstruction, commissurotomy, leaflet repair, or papillary muscle repair.
2260	Valvuloplasty converted to valve replacement in the same operation, Mitral	

840	Mitral stenosis, Supravalvar mitral ring repair	Supravalvar mitral ring repair.
850	Valve replacement, Mitral (MVR)	Replacement of mitral valve with prosthetic valve, any kind, in suprannular or annular position.
860	Valve surgery, Other, Mitral	Other mitral valve surgery not specified in procedure codes.
870	Norwood procedure	<p>The Norwood operation is synonymous with the term 'Norwood (Stage 1)' and is defined as an aortopulmonary connection and neoaortic arch construction resulting in univentricular physiology and pulmonary blood flow controlled with a calibrated systemic-to-pulmonary artery shunt, or a right ventricle to pulmonary artery conduit, or rarely, a cavopulmonary connection. When coding the procedure "Norwood procedure", the primary procedure of the operation should be "Norwood procedure". The second procedure (Procedure 2 after the Norwood procedure) must then document the source of pulmonary blood flow and be chosen from the following eight choices:</p> <ol style="list-style-type: none"> 1. Shunt, Systemic to pulmonary, Modified Blalock-Taussig Shunt (MBTS) 2. Shunt, Systemic to pulmonary, Central (from aorta or to main pulmonary artery) 3. Shunt, Systemic to pulmonary, Other 4. Conduit placement, RV to PA 5. Bidirectional cavopulmonary anastomosis (BDCPA) (bidirectional Glenn) 6. Glenn (unidirectional cavopulmonary anastomosis) (unidirectional Glenn) 7. Bilateral bidirectional cavopulmonary anastomosis (BBDCPA) (bilateral bidirectional Glenn) 8. HemiFontan
880	HLHS biventricular repair	Performed in patients who have small but adequately sized ventricles to support systemic circulation. These patients usually have small, but not stenotic, aortic and/or mitral valves. Primary biventricular repair has consisted of extensive aortic arch and ascending aorta enlargement with a patch, closure of interventricular and interatrial communications, and conservative approach for left ventricular outflow tract
2160	Hybrid Approach "Stage 1", Application of RPA & LPA bands	A "Hybrid Procedure" is defined as a procedure that combines surgical and transcatheter interventional approaches. The term "Hybrid approach" is used somewhat differently than the term "Hybrid Procedure". A "Hybrid approach" is defined as any of a group of procedures that fit into the general silo of procedures developed from the combined use of surgical and transcatheter interventional techniques. Therefore, not all procedures classified as "Hybrid approach" are truly "Hybrid Procedures".

2170	Hybrid Approach "Stage 1", Stent placement in arterial duct (PDA)	A "Hybrid Procedure" is defined as a procedure that combines surgical and transcatheter interventional approaches. The term "Hybrid approach" is used somewhat differently than the term "Hybrid Procedure". A "Hybrid approach" is defined as any of a group of procedures that fit into the general silo of procedures developed from the combined use of surgical and transcatheter interventional techniques. Therefore, not all procedures classified as "Hybrid approach" are truly "Hybrid Procedures".
2180	Hybrid Approach "Stage 1", Stent placement in arterial duct (PDA) + application of RPA & LPA bands	A "Hybrid Procedure" is defined as a procedure that combines surgical and transcatheter interventional approaches. The term "Hybrid approach" is used somewhat differently than the term "Hybrid Procedure". A "Hybrid approach" is defined as any of a group of procedures that fit into the general silo of procedures developed from the combined use of surgical and transcatheter interventional techniques. Therefore, not all procedures classified as "Hybrid approach" are truly "Hybrid Procedures".
2140	Hybrid approach "Stage 2", Aortopulmonary amalgamation + Superior Cavopulmonary anastomosis(es) + PA Debanding + Aortic arch repair (Norwood [Stage 1] + Superior Cavopulmonary anastomosis(es) + PA Debanding)	A "Hybrid Procedure" is defined as a procedure that combines surgical and transcatheter interventional approaches. The term "Hybrid approach" is used somewhat differently than the term "Hybrid Procedure". A "Hybrid approach" is defined as any of a group of procedures that fit into the general silo of procedures developed from the combined use of surgical and transcatheter interventional techniques. Therefore, not all procedures classified as "Hybrid approach" are truly "Hybrid Procedures". It should be acknowledged that a Hybrid approach "Stage 2" (Aortopulmonary amalgamation + Superior Cavopulmonary anastomosis(es) + PA Debanding, with or without Aortic arch repair) gets its name not because it has any actual hybrid elements, but because it is part of a planned staged approach that is typically commenced with a hybrid procedure.
2150	Hybrid approach "Stage 2", Aortopulmonary amalgamation + Superior Cavopulmonary anastomosis(es) + PA Debanding + Without aortic arch repair	A "Hybrid Procedure" is defined as a procedure that combines surgical and transcatheter interventional approaches. The term "Hybrid approach" is used somewhat differently than the term "Hybrid Procedure". A "Hybrid approach" is defined as any of a group of procedures that fit into the general silo of procedures developed from the combined use of surgical and transcatheter interventional techniques. Therefore, not all procedures classified as "Hybrid approach" are truly "Hybrid Procedures". It should be acknowledged that a Hybrid approach "Stage 2" (Aortopulmonary amalgamation + Superior Cavopulmonary anastomosis(es) + PA Debanding, with or without Aortic arch repair) gets its name not because it has any actual hybrid elements, but because it is part of a planned staged approach that is typically commenced with a hybrid procedure

890	Transplant, Heart	Heart transplantation, any technique, allograft or xenograft.
900	Transplant, Heart and lung	Heart and lung (single or double) transplantation.
910	Partial left ventriculectomy (LV volume reduction surgery) (Batista)	Wedge resection of LV muscle, with suturing of cut edges together, to reduce LV volume.
920	Pericardial drainage procedure	Pericardial drainage can include a range of therapies including, but not limited to: pericardiocentesis, pericardiostomy tube placement, pericardial window creation, and open pericardial drainage (pericardiotomy).
930	Pericardiectomy	Surgical removal of the pericardium.
940	Pericardial procedure, Other	Other pericardial procedures that include, but are not limited to: pericardial reconstruction for congenital absence of the pericardium, pericardial biopsy, pericardial mass or cyst excision.
950	Fontan, Atrio-pulmonary connection	Fontan-type procedure with atrio-pulmonary connection.
960	Fontan, Atrio-ventricular connection	Fontan-type procedure with atrio-ventricular connection, either direct or with RA-RV conduit, valved or nonvalved.
970	Fontan, TCPC, Lateral tunnel, Fenestrated	Total cavopulmonary connection using an intraatrial lateral tunnel construction, with fenestration.
980	Fontan, TCPC, Lateral tunnel, Nonfenestrated	Total cavopulmonary connection using an intraatrial lateral tunnel construction, with no fenestration.
1000	Fontan, TCPC, External conduit, Fenestrated	Total cavopulmonary connection using an external conduit to connect the infradiaphragmatic systemic venous return to the pulmonary artery, with fenestration.
1010	Fontan, TCPC, External conduit, Nonfenestrated	Total cavopulmonary connection using an external conduit to connect the infradiaphragmatic systemic venous return to the pulmonary artery, with no fenestration.
1025	Fontan revision or conversion (Re-do Fontan)	Revision of a previous Fontan procedure to a total cavopulmonary connection.
1030	Fontan, Other	Other Fontan procedure not specified in procedure codes. May include takedown of a Fontan procedure.
2340	Fontan + Atrioventricular valvuloplasty	
1035	Ventricular septation	Creation of a prosthetic ventricular septum. Surgical procedure used to septate univentricular hearts with two atrioventricular valves. Additional procedures, such as resection of subpulmonic stenosis, should be listed separately.
1050	Congenitally corrected TGA repair, Atrial switch and ASO (double switch)	Repair of congenitally corrected TGA by concomitant atrial switch (Mustard or Senning) and arterial switch operation. VSD closure is usually performed as well; this should be coded separately.
1060	Congenitally corrected TGA repair, Atrial switch and Rastelli	Repair of congenitally corrected TGA by concomitant atrial switch (Mustard or Senning) and VSD closure to the aortic valve with placement of an RV-to-PA conduit.
1070	Congenitally corrected TGA repair, VSD closure	Repair of congenitally corrected TGA by VSD closure only.

1080	Congenitally corrected TGA repair, VSD closure and LV to PA conduit	Repair of congenitally corrected TGA by VSD closure and placement of an LV-to-PA conduit.
1090	Congenitally corrected TGA repair, Other	Any procedures for correction of CCTGA not otherwise specified in other listed procedure codes.
1110	Arterial switch operation (ASO)	Arterial switch operation is used for repair of transposition of the great arteries (TGA). The pulmonary artery and aorta are transected and translocated so that the pulmonary artery arises from the right ventricle and the aorta from the left ventricle. Coronary artery transfer is also accomplished.
1120	Arterial switch operation (ASO) and VSD repair	Arterial switch operation is used for repair of transposition of the great arteries (TGA). The pulmonary artery and aorta are transected and translocated so that the pulmonary artery arises from the right ventricle and the aorta from the left ventricle. Coronary artery transfer is also accomplished. The VSD is closed, usually with a patch.
1123	Arterial switch procedure + Aortic arch repair	Concomitant arterial switch operation and repair of the aortic arch in patients with transposition of the great arteries with intact ventricular septum and associated coarctation of the aorta or interrupted aortic arch.
1125	Arterial switch procedure and VSD repair + Aortic arch repair	Concomitant arterial switch operation with VSD closure and repair of aortic arch in patients with transposition of the great arteries with VSD and associated coarctation of the aorta or interrupted aortic arch.
1130	Senning	Atrial baffle procedure for rerouting of venous flow in TGA effecting a "physiological repair". The caval flow is directed behind the baffle to the mitral valve, left ventricle and pulmonary artery while the pulmonary venous flow is directed in front of the baffle to the tricuspid valve, right ventricle, and aorta. The Senning procedure uses atrial wall to construct the baffle.
1140	Mustard	Atrial baffle procedure for rerouting of venous flow in TGA effecting a "physiological repair". The caval flow is directed behind the baffle to the mitral valve, left ventricle and pulmonary artery while pulmonary venous flow is directed in front of the baffle to the tricuspid valve, right ventricle, and aorta. The Mustard procedure uses patch material to construct the baffle.
1145	Atrial baffle procedure, Mustard or Senning revision	Revision of a previous atrial baffle procedure (either Mustard or Senning), for any reason (e.g., obstruction, baffle leak).
1150	Rastelli	Most often used for patients with TGA-VSD and significant LVOTO, the Rastelli operation consists of an

1160	REV	The Lecompte (REV) intraventricular repair is designed for patients with abnormalities of ventriculoarterial connection in whom a standard intraventricular tunnel repair cannot be performed. It is also suitable for patients in whom an arterial switch procedure with tunneling of the VSD to the pulmonary artery cannot be performed because of pulmonary (left ventricular outflow tract) stenosis. A right ventriculotomy incision is made. The infundibular (conal) septum, located between the two semilunar valves, is aggressively resected if its presence interferes with the construction of a tunnel from the VSD to the aorta. The VSD is then tunneled to the aorta. The decision to perform or not to perform the Lecompte maneuver should be made at the beginning of the operation. If the Lecompte maneuver is not performed the pulmonary artery is translocated to the right ventricular outflow tract on the side of the aorta that provides the shortest route. (When the decision to perform the Lecompte maneuver has been made, the great vessels are transected and this maneuver is performed at the
2190	Aortic root translocation over left ventricle (Including Nikaidoh procedure)	
2210	TGA, Other procedures (Kawashima, LV-PA conduit, other)	
1180	DORV, Intraventricular tunnel repair	Repair of DORV using a tunnel closure of the VSD to the aortic valve. This also includes the posterior straight tunnel repair of Kawashima
1200	DOLV repair	Because of the morphologic variability of DOLV, there are many approaches to repair, including: intraventricular tunnel repair directing the VSD to the pulmonary valve, the REV procedure, or the Rastelli procedure. In the case of DOLV use this code for tunnel closure to the pulmonary valve. If the REV or Rastelli procedures are performed then use those respective codes.
1210	Coarctation repair, End to end	Repair of coarctation of aorta by excision of the coarctation segment and end-to-end circumferential anastomosis of the aorta.
1220	Coarctation repair, End to end, Extended	Repair of coarctation of the aorta by excision of the coarctation segment and end-to-end anastomosis of the oblique ends of the aorta, creating an extended anastomosis.
1230	Coarctation repair, Subclavian flap	Repair of coarctation of the aorta by ligating, dividing, and opening the subclavian artery, incising the coarctation site, and folding down the subclavian artery onto the incision in the aorta, suturing the subclavian "flap" in place, creating a roof over the area of the previous coarctation.
1240	Coarctation repair, Patch aortoplasty	Repair of coarctation of the aorta by incising the coarctation site with placement of a patch sutured in place longitudinally along the aortotomy edge.

1250	Coarctation repair, Interposition graft	Repair of coarctation of the aorta by resection of the coarctation segment and placement of a prosthetic tubular interposition graft anastomosed circumferentially to the cut ends of the aorta.
1260	Coarctation repair, Other	Any repair of coarctation not specified in procedure codes. This may include, for example, a combination of two approaches for coarctation repair or extra-anatomic bypass graft, etc.
1275	Coarctation repair + VSD repair	Coarctation of aorta repair, any technique, and simultaneous VSD repair, any type VSD, any type repair.
1280	Aortic arch repair	Aortic arch repair, any technique.
1285	Aortic arch repair + VSD repair	Aortic arch repair, any technique, and simultaneous VSD repair, any type VSD, any type repair. This includes repair of IAA with VSD.
1290	Coronary artery fistula ligation	Coronary artery fistula repair using any technique. If additional technique information may be supplied by another procedure code, please list separately (e.g., bypass graft).
1291	Anomalous origin of coronary artery from pulmonary artery repair	Repair of anomalous origin of the coronary artery (any) from the pulmonary artery, by any technique (ligation, translocation with aortic implantation, Takeuchi operation, bypass graft). If additional technique information may be supplied by another procedure code, please list separately (for example, bypass graft).
1300	Coronary artery bypass	Coronary artery bypass graft procedure, any technique (with or without CPB, venous or arterial graft, one or more grafts, etc.), for any coronary artery pathology (coronary arterial fistula, aneurysm, coronary bridging, atresia of left main, acquired coronary artery disease, etc.).
1310	Coronary artery procedure, Other	Any coronary artery procedure not specifically listed.
1320	Interrupted aortic arch repair	Repair of interrupted aortic arch (any type) by any technique (direct anastomosis, prosthetic graft, etc). Does not include repair of IAA-VSD.
1330	PDA closure, Surgical	Closure of a PDA by any surgical technique (ligation, division, clip) using any approach (i.e., thoracotomy, thoracoscopic, etc).
1340	PDA closure, Device	Closure of a PDA by device using transcatheter techniques.
1360	Vascular ring repair	Repair of vascular ring (any type, except pulmonary artery sling) by any technique.
1365	Aortopexy	Surgical fixation of the aorta to another structure (usually the posterior aspect of the sternum) to relieve compression on another vessel or structure (e.g., trachea).
1370	Pulmonary artery sling repair	Pulmonary artery sling repair by any technique.
1380	Aortic aneurysm repair	Aortic aneurysm repair by any technique.
1390	Aortic dissection repair	Aortic dissection repair by any technique
1410	Transplant, lung(s)	Lung or lobe transplantation of any type.

1450	Pacemaker implantation, Permanent	Implantation of a permanent pacemaker of any type (e.g., single-chamber, dual-chamber, atrial antitachycardia), with any lead configuration or type (atrial, ventricular, atrial and ventricular, transvenous, epicardial, transmural), by any technique (sternotomy, thoracotomy etc).
1460	Pacemaker procedure	Any revision to a previously placed pacemaker system including revisions to leads, generators, pacemaker pockets. This may include explantation of pacemakers or leads as well.
2350	Explantation of pacing system	
1470	ICD (AICD) implantation	Implantation of an (automatic) implantable cardioverter defibrillator system.
1480	ICD (AICD) ([automatic] implantable cardioverter defibrillator) procedure	Any revision to a previously placed AICD including revisions to leads, pads, generators, pockets. This may include explantation procedures as well.
1490	Arrhythmia surgery - atrial, Surgical Ablation	Surgical ablation (any type) of any atrial arrhythmia.
1500	Arrhythmia surgery - ventricular, Surgical Ablation	Surgical ablation (any type) of any ventricular arrhythmia.
1590	Shunt, Systemic to pulmonary, Modified Blalock-Taussig Shunt (MBTS)	Placement of a tube graft from a branch of the aortic arch to the pulmonary artery with or without bypass, from any approach (thoracotomy, sternotomy).
1600	Shunt, Systemic to pulmonary, Central (from aorta or to main pulmonary artery)	A direct anastomosis or placement of a tube graft from the aorta to the pulmonary artery with or without bypass, from any approach (thoracotomy, sternotomy).
1610	Shunt, Systemic to pulmonary, Other	Placement of any other systemic-to-pulmonary artery shunt, with or without bypass, from any approach (thoracotomy, sternotomy) that is not otherwise coded. Includes classic Blalock-Taussig systemic-to-pulmonary artery shunt.
1630	Shunt, Ligation and takedown	Takedown of any shunt.
2095	Shunt, Reoperation	
1640	PA banding (PAB)	Placement of a pulmonary artery band, any type.
1650	PA debanding	Debanding of pulmonary artery. Please list separately any pulmonary artery reconstruction required.
1660	Damus-Kaye-Stansel procedure (DKS) (creation of AP anastomosis without arch reconstruction)	In the Damus-Kaye-Stansel procedure the proximal transected main pulmonary artery is connected by varying techniques to the aorta.
1670	Bidirectional cavopulmonary anastomosis (BDCPA) (bidirectional Glenn)	Superior vena cava to pulmonary artery anastomosis allowing flow to both pulmonary arteries with an end-to side superior vena-to-pulmonary artery anastomosis.
1680	Glenn (unidirectional cavopulmonary anastomosis) (unidirectional Glenn)	Superior vena cava to ipsilateral pulmonary artery anastomosis (i.e., LSVC to LPA, RSVC to RPA).

1690	Bilateral bidirectional cavopulmonary anastomosis (BBDCPA) (bilateral bidirectional Glenn)	Bilateral superior vena cava-to-pulmonary artery anastomoses (requires bilateral SVCs).
1700	HemiFontan	A HemiFontan is an operation that includes a bidirectional superior vena cava (SVC)-to-pulmonary artery anastomosis and the connection of this "SVCpulmonary artery amalgamation" to the atrium, with a "dam" between this "SVC-pulmonary artery amalgamation" and the atrium. This operation can be accomplished with a variety of operative strategies including the following two techniques and other techniques that combine elements of both of these approaches: (1) Augmenting both branch pulmonary arteries with a patch and suturing the augmented branch pulmonary arteries to an incision in the medial aspect of the superior vena cava. (With this approach, the pulmonary artery patch forms a roof over the SVC-topulmonary artery anastomosis and also forms a "dam" between the SVC-pulmonary artery amalgamation and the right atrium.) (2) Anastomosing both ends of the divided SVC to incisions in the top and bottom of the right pulmonary artery, and using a separate patch to close junction of the SVC and the right atrium.
2330	Superior cavopulmonary anastomosis(es) (Glenn or HemiFontan) + Atrioventricular valvuloplasty	
2130	Superior Cavopulmonary anastomosis(es) + PA reconstruction	
1720	Aneurysm, Ventricular, Right, Repair	Repair of right ventricular aneurysm, any technique.
1730	Aneurysm, Ventricular, Left, Repair	Repair of left ventricular aneurysm, any technique.
1740	Aneurysm, Pulmonary artery, Repair	Repair of pulmonary artery aneurysm, any technique.
1760	Cardiac tumor resection	Resection of cardiac tumor, any type.
1780	Pulmonary AV fistula repair/occlusion	Repair or occlusion of a pulmonary arteriovenous fistula.
1790	Ligation, Pulmonary artery	Ligation or division of the pulmonary artery. Most often performed as a secondary procedure.
1802	Pulmonary embolectomy, Acute pulmonary embolus	Acute pulmonary embolism (clot) removal, through catheter or surgery.
1804	Pulmonary embolectomy, Chronic pulmonary embolus	Chronic pulmonary embolism (clot) removal, through catheter or surgery.
1830	Ligation, Thoracic duct	Ligation of the thoracic duct; most commonly for persistent chylothorax.
1860	Mediastinal procedure	Any non-cardiovascular mediastinal procedure not otherwise listed.

