

**List of NQF-Endorsed Perinatal and Reproductive Health Consensus Standards  
November, 2011**

Measure Number	Title	Description	Steward
<b>REPRODUCTIVE HEALTH</b>			
<b>0502</b>	Pregnancy test for female abdominal pain patients.	Pregnancy test for female abdominal pain patients.	American College of Emergency Physicians
<b>0651</b>	Ultrasound determination of pregnancy location for pregnant patients with abdominal pain	Percentage of pregnant patients who present to the ED with a chief complaint of abdominal pain and or vaginal bleeding who receive a trans-abdominal or trans-vaginal ultrasound.	American College of Emergency Physicians
<b>PREGNANCY CARE</b>			
<b>0012*</b>	Prenatal Screening for Human Immunodeficiency Virus (HIV)	Percentage of patients who gave birth during a 12-month period who were screened for HIV infection during the first or second prenatal care visit.	AMA-PCPI
<b>0014*</b>	Prenatal Anti-D Immune Globulin	Percentage of D-negative, unsensitized patients who gave birth during a 12-month period who received anti-D immune globulin at 26-30 weeks gestation.	AMA-PCPI
<b>0015*</b>	Prenatal Blood Groups (ABO), D (Rh) Type	Percentage of patients who gave birth during a 12-month period who had a determination of blood group (ABO) and D (Rh) type by the second prenatal care visit.	AMA-PCPI
<b>0016*</b>	Prenatal Blood Group Antibody Testing	Percentage of patients who gave birth during a 12-month period who were screened for blood group antibodies during the first or second prenatal care visit.	AMA-PCPI
<b>0476</b>	Appropriate Use of Antenatal Steroids	Mothers receiving antenatal steroids during pregnancy at any time prior to delivery of a preterm infant	Providence St. Vincent Medical Center
<b>0582</b>	Diabetes and Pregnancy: Avoidance of Oral Hypoglycemic Agents	This measure identifies pregnant women with diabetes who are not taking an oral hypoglycemic agent.	Resolution Health, Inc.
<b>0606*</b>	Pregnant women that had HIV testing.	This measure identifies pregnant women who had an HIV test during their pregnancy.	Ingenix
<b>0607*</b>	Pregnant women that had syphilis screening.	This measure identifies pregnant women who had a syphilis test during their pregnancy.	Ingenix
<b>0608*</b>	Pregnant women that had HBsAg testing.	This measure identifies pregnant women who had a HBsAg (hepatitis B) test during their pregnancy.	Ingenix
<b>0652</b>	RH Immunoglobulin (rhogam) for RH negative pregnant women at risk of fetal blood exposure	Percent of RH negative pregnant women at risk of fetal blood exposure who receive Rhogam the ED.	American College of Emergency Physicians
<b>1391</b>	Frequency of Ongoing Prenatal Care (FPC)	The percentage of Medicaid deliveries between November 6 of the year prior to the measurement year and November 5 of the measurement year that received the following number of expected prenatal visits. <ul style="list-style-type: none"> <li>•&gt;21 percent of expected visits</li> <li>•21 percent–40 percent of expected visits</li> <li>•41 percent–60 percent of expected visits</li> <li>•61 percent–80 percent of expected visits</li> <li>•=81 percent of expected visits</li> </ul>	NCQA
<b>1517</b>	Prenatal and Postpartum Care	The percentage of deliveries of live births between November 6 of the year prior to the measurement year and November 5 of the measurement year. For these women, the	NCQA

\*Measures have not been resubmitted for endorsement; endorsement will be retired at the completion of the Perinatal and Reproductive Healthcare Project.

**List of NQF-Endorsed Perinatal and Reproductive Health Consensus Standards  
November, 2011**

		<p>measure assesses the following facets of prenatal and postpartum care.</p> <ul style="list-style-type: none"> <li>• Rate 1: Timeliness of Prenatal Care. The percentage of deliveries that received a prenatal care visit as a member of the organization in the first trimester or within 42 days of enrollment in the organization.</li> <li>• Rate 2: Postpartum Care. The percentage of deliveries that had a postpartum visit on or between 21 and 56 days after delivery.</li> </ul>	
<b>CHILDBIRTH</b>			
<b>0333*</b>	Severity-Standardized ALOS - Deliveries	Standardized ALOS for deliveries	Leapfrog Group
<b>0278</b>	Low birth weight (PQI 9)	This measure is used to assess the number of low birth weight infants per 100 births. See Notes.	AHRQ
<b>0469</b>	Elective delivery prior to 39 completed weeks gestation	Percentage of babies electively delivered prior to 39 completed weeks gestation	Hospital Corporation of America
<b>0470</b>	Incidence of Episiotomy	Percentage of vaginal deliveries during which an episiotomy is performed	Christiana Care Health System
<b>0471</b>	Cesarean Rate for low-risk first birth women (aka NTSV CS rate)	Percentage of low-risk first birth women (aka NTSV CS rate: nulliparous, term, singleton, vertex) with a Cesarean rate that has the most variation among practitioners, hospitals, regions and states. Unlike other cesarean measures, it focuses attention on the proportion of cesarean births that is affected by elective medical practices such as induction and early labor admission. Furthermore, the success (or lack thereof) of management of the first labor directly impacts the remainder of the woman's reproductive life (especially given the current high rate of repeat cesarean births).	California Maternal Quality Care Collaborative
<b>0472</b>	Prophylactic Antibiotic Received Within One Hour Prior to Surgical Incision or at the Time of Delivery – Cesarean section.	Percentage of patients undergoing cesarean section who receive prophylactic antibiotics within one hour prior to surgical incision or at the time of delivery.	Massachusetts General Hospital/Partners Health Care System
<b>0473</b>	Appropriate DVT prophylaxis in women undergoing cesarean delivery	Measure adherence to current ACOG, ACCP recommendations for use of DVT prophylaxis in women undergoing cesarean delivery	Hospital Corporation of America
<b>0474</b>	Birth Trauma Rate: Injury to Neonates (PSI #17)	Percentage of neonates with specific birth trauma per 1000 births. Exclude infants with injury to skeleton and osteogenesis imperfecta, subdural or cerebral hemorrhage in preterm infant.	AHRQ, National Perinatal Information Center
<b>0477</b>	Under 1500g infant Not Delivered at Appropriate Level of Care	The number per 1,000 livebirths of <1500g infants delivered at hospitals not appropriate for that size infant.	California Maternal Quality Care Collaborative
<b>NEWBORN CARE</b>			
<b>0303</b>	Late sepsis or meningitis in neonates (risk-adjusted)	Percentage of infants born at the hospital, whose birth weight is between 401 and 1500 grams OR whose gestational age is between 22 weeks 0 days and 29 weeks 6 days with late sepsis or meningitis with one or more of the following criteria: Bacterial Pathogen,	Vermont Oxford Network

\*Measures have not been resubmitted for endorsement; endorsement will be retired at the completion of the Perinatal and Reproductive Healthcare Project.

**List of NQF-Endorsed Perinatal and Reproductive Health Consensus Standards  
November, 2011**

		Coagulase Negative Staphylococcus, Fungal Infection	
<b>0304</b>	Late sepsis or meningitis in Very Low Birth Weight (VLBW) neonates (risk-adjusted)	Percentage of infants born at the hospital, whose birth weight is between 401 and 1500 grams OR whose gestational age is between 22 weeks 0 days and 29 weeks 6 days, who have late sepsis or meningitis, with one or more of the following criteria: Bacterial Pathogen, Coagulase Negative Staphylococcus, Fungal Infection	Vermont Oxford Network
<b>0475</b>	Measurement of Hepatitis B Vaccine Administration to All Newborns Prior to Hospital or Birthing Facility Discharge	Percentage of newborns administered hepatitis B vaccine prior to discharge from the birthing facility or hospital, subtract the number of newborns who died prior to discharge, and divide this number by the number of live newborns discharged from the birthing facility or hospital during a given time period (perhaps annually) to identify the hepatitis B vaccine coverage rate for newborns at a single birthing facility or hospital.	CDC
<b>0478</b>	Nosocomial Blood Stream Infections in Neonates (NQI #3)	Percentage of qualifying neonates with selected bacterial blood stream infections	AHRQ
<b>0479</b>	Birth dose of hepatitis B vaccine and hepatitis immune globulin for newborns of mothers with chronic hepatitis B	Percentage of newborns to hepatitis B surface antigen (HBsAg)-positive mothers who receive a birth dose of hepatitis B virus (HBV) vaccine and hepatitis B immune globulin (HBIG)	Asian Liver Center at Stanford University
<b>0480</b>	Exclusive Breastfeeding during Birth Hospitalization	Exclusive Breastfeeding (BF) for the first 6 mos of neonatal life has long been the expressed goal of WHO, DHHS, APA, and ACOG. Holding perinatal and intrapartum providers accountable is an important way to incent greater efforts during the critical prenatal and immediate postpartum periods where BF attitudes are solidified.	California Maternal Quality Care Collaborative
<b>0481</b>	First temperature measured within one hour of admission to the NICU.	Percent of NICU admissions with a birth weight of 501-1500g with a first temperature taken within 1 hour of NICU admission.	Vermont Oxford Network
<b>0482</b>	First NICU Temperature < 36 degrees C	Percent of all NICU admissions with a birth weight of 501-1500g whose first temperature was measured within one hour of admission to the NICU and was below 36 degrees Centigrade.	Vermont Oxford Network
<b>0483</b>	Proportion of infants 22 to 29 weeks gestation screened for retinopathy of prematurity.	Proportion of infants 22 to 29 weeks screened for retinopathy of prematurity using the guidelines from the American Academy of Pediatrics	Vermont Oxford Network
<b>0484</b>	Proportion of infants 22 to 29 weeks gestation treated with surfactant who are treated within 2 hours of birth.	Number of infants 22 to 29 weeks gestation treated with surfactant within 2 hours of birth	Vermont Oxford Network
<b>0485*</b>	Neonatal Immunization	Percent of neonates with a length of stay greater than 60 days receiving DPT, Hepatitis B, Polio, Hib, and PCV immunizations in adherence with current guidelines.	Child Health Corporation of America
<b>0145*</b>	Neonate immunization administration	Percentage of patient refined diagnostic-related groups (APR-DRG) who received neonate immunization administration	Child Health Corporation of America

\*Measures have not been resubmitted for endorsement; endorsement will be retired at the completion of the Perinatal and Reproductive Healthcare Project.

**List of NQF-Endorsed Perinatal and Reproductive Health Consensus Standards  
November, 2011**

<b>0716</b>	Healthy term newborn	Percent of term singleton livebirths (excluding those with diagnoses originating in the fetal period) who DO NOT have significant complications during birth or the nursery care.	California Maternal Quality Care Collaborative
<b>1354</b>	Hearing screening prior to hospital discharge (EHDI-1a)	This measure assesses the proportion of births that have been screened for hearing loss before hospital discharge.	CDC
<b>1351</b>	Proportion of infants covered by Newborn Bloodspot Screening (NBS)	What percentage of infants had bloodspot newborn screening performed as mandated by state of birth?	HRSA-MCHB
<b>714</b>	Standardized mortality ratio for neonates undergoing non-cardiac surgery	Ratio of observed to expected rate of in-hospital mortality following non-cardiac surgery among infants less than or equal to 30 days of age, risk-adjusted.	Children's Hospital Boston - Program for Patient Safety & Quality

\*Measures have not been resubmitted for endorsement; endorsement will be retired at the completion of the Perinatal and Reproductive Healthcare Project.