

## NQF-Endorsed® Preventive Services Endorsement Maintenance Standards

Measure	Steward	Description
#0023: Body Mass Index (BMI) in adults > 18 years of age	City of New York Department of Health and Mental Hygiene	Percentage of adults with BMI documentation in the past 24 month.
#0024: Body Mass Index (BMI) 2 through 18 years of age	National Committee for Quality Assurance	Percentage children, 2 through 18 years of age, whose weight is classified based on BMI percentile for age and gender
#0421: Adult Weight Screening and Follow-Up	Centers for Medicare & Medicaid Services	<p>Percentage of patients aged 18 years and older with a calculated BMI documented in the medical record AND if the most recent BMI is outside the parameters, a follow up plan is documented.</p> <p>Parameters: age 65 and older BMI &gt; or = 30 or &lt; 22; age 18-64 BMI &gt; or = 25 or &lt; 18.5</p>
#0026: Measure pair - a. Tobacco use prevention for infants, children and adolescents, b. Tobacco use cessation for infants, children and adolescents	Institute for Clinical Systems Improvement	<p>Percentage of patients' charts showing either that there is no tobacco use/exposure or (if a user) that the current use was documented at the most recent clinic visit</p> <p>Percentage of patients with documented tobacco use or exposure at the latest visit who also have documentation that their cessation interest was assessed or that they received advice to quit</p>
#0027: Smoking Cessation, Medical assistance: a. Advising Smokers to Quit, b. Discussing Smoking Cessation Medications, c. Discussing Smoking Cessation Strategies	National Committee for Quality Assurance	<p>Percentage of patients who received advice to quit smoking</p> <p>Percentage of patients whose practitioner recommended or discussed smoking cessation medications</p>
#0028: Measure pair: a. Tobacco Use Assessment, b. Tobacco Cessation Intervention	American Medical Association - Physician Consortium for Performance Improvement	<p>Percentage of patients who were queried about tobacco use one or more times during the two-year measurement period</p> <p>Percentage of patients identified as tobacco users who received cessation intervention during the two-year measurement period</p>

## NQF-Endorsed® Preventive Services Endorsement Maintenance Standards

#0629: Male Smokers or Family History of Abdominal Aortic Aneurysm (AAA) - Consider Screening for AAA	ActiveHealth Management	Percentage of men age 65-75 years with history of tobacco use or men age 60 yrs and older with a family history of abdominal aortic aneurysm who were screened for AAA
#0029: Counseling on physical activity in older adults - a. Discussing Physical Activity, b. Advising Physical Activity	National Committee for Quality Assurance	Percentage patients 65 years of age and older who reported: discussing their level of exercise or physical activity with a doctor or other health provider in the last 12 months
#0031: Breast Cancer Screening	National Committee for Quality Assurance	Percentage of eligible women 50-69 who receive a mammogram in a two year period
#0032: Cervical Cancer Screening	National Committee for Quality Assurance	Percentage of women 18-64 years of age, who received one or more Pap tests during the measurement year or the two years prior to the measurement year
#0034: Colorectal Cancer Screening	National Committee for Quality Assurance	Percentage of adults 50-80 years of age who had appropriate screening for colorectal cancer (CRC) including fecal occult blood test during the measurement year or, flexible sigmoidoscopy during the measurement year or the four years prior to the measurement year or, double contrast barium enema during the measurement year or the four years prior to the measurement year or, colonoscopy during the measurement year or the nine years prior to the measurement year
#0579: Annual Cervical Cancer Screening for High-Risk Patients	Resolution Health, Inc.	This measure identifies women age 12 to 65 diagnosed with cervical dysplasia (CIN 2), cervical carcinoma-in-situ, or HIV/AIDS prior to the measurement year, and who still have a cervix, who had a cervical CA screen during the measurement year.

## NQF-Endorsed® Preventive Services Endorsement Maintenance Standards

#0033: Chlamydia screening in women	National Committee for Quality Assurance	Percentage of eligible women who were identified as sexually active who had at least one test for chlamydia during the measurement year
#0037: Osteoporosis testing in older women	National Committee for Quality Assurance	Percentage of female patients aged 65 and older who reported receiving a bone density test (BMD) to check for osteoporosis
#0614: Steroid Use - Osteoporosis Screening	ActiveHealth Management	Percentage of patients, 18 and older, who have been on chronic steroids for at least 180 days in the past 9 months and who had a bone density evaluation or osteoporosis treatment
#0038: Childhood Immunization Status	National Committee for Quality Assurance	Percentage of children 2 years of age who had four DtaP/DT, three IPV, one MMR, three H influenza type B, three hepatitis B, one chicken pox vaccine (VZV) and four pneumococcal conjugate vaccines by their second birthday. The measure calculates a rate for each vaccine and two separate combination rates
#0145: Neonate immunization administration	Child Health Corporation of America	Percentage of patient refined diagnostic-related groups (APR-DRG) who received neonate immunization administration
#0039: Flu shots for Adults Ages 50-64	National Committee for Quality Assurance	Percentage of patients age 50-64 who report having received an influenza vaccination during the past influenza vaccination season
#0040: Flu Shot for Older Adults	National Committee for Quality Assurance	Percentage of patients age 65 and over who received an influenza vaccination from September through December of the year
#0041: Influenza Vaccination	American Medical Association - Physician Consortium for Performance Improvement	Percentage of patients who received an influenza vaccination
#0149: Influenza vaccination	Centers for Medicare & Medicaid Services	Percentage of patients discharged during October, November, December, January, or February with pneumonia, age 50 and older, who were screened for influenza vaccine status and were vaccinated prior to discharge, if indicated

## NQF-Endorsed® Preventive Services Endorsement Maintenance Standards

#0226: Influenza Vaccination in the ESRD Population- Facilities	Kidney Care Partners	Percentage of all ESRD patients aged 6 months and older receiving hemodialysis and/or peritoneal dialysis during the time from October 1 (or when the influenza vaccine became available) and March 31 who receive an influenza vaccination, were assessed and offered an influenza vaccination but decline, or were assessed and determined to have a medical contraindication to the influenza vaccination.
#0227: Influenza Immunization	American Medical Association - Physician Consortium for Performance Improvement	Percentage of patients aged 18 years and older with a diagnosis of ESRD and receiving dialysis who received the influenza immunization during the flu season (September through February)
#0431: Influenza Vaccination Coverage among Healthcare Personnel	Centers for Disease Control and Prevention	Percentage of healthcare personnel (HCP) who receive the influenza vaccination.
#0522: Influenza Immunization Received for Current Flu Season	Centers for Medicare & Medicaid Services	Percent of patients who received influenza immunization for the current flu season from this home health agency
#0043: Pneumonia vaccination status for older adults	National Committee for Quality Assurance	Percentage of patients 65 years of age and older who ever received a pneumococcal vaccination
#0044: Pneumonia Vaccination	National Committee for Quality Assurance	Percentage of patients who ever received a pneumococcal vaccination
#0150: Pneumococcal vaccination	Centers for Medicare & Medicaid Services	Percentage of patients with pneumonia, age 65 and older, who were screened for pneumococcal vaccine status and were vaccinated prior to discharge, if indicated
#0525: Pneumococcal Polysaccharide Vaccine (PPV) Ever Received	Centers for Medicare & Medicaid Services	Percent of patients who have ever received Pneumococcal Polysaccharide Vaccine (PPV)
#0617: High Risk for Pneumococcal Disease - Pneumococcal Vaccination	ActiveHealth Management	Percentage of patients age 5-64 with a high risk condition or age 65 years and older who received the pneumococcal vaccine

## NQF-Endorsed® Preventive Services Endorsement Maintenance Standards

#0508: Inappropriate use of “probably benign” assessment category in mammography screening	American Medical Association - Physician Consortium for Performance Improvement	Percentage of final reports for screening mammograms that are classified as “probably benign”
#0509: Reminder system for mammograms	American Medical Association - Physician Consortium for Performance Improvement	Percentage of patients aged 40 years and older undergoing a screening mammogram whose information is entered into a reminder system* with a target due date for the next mammogram
#0573: HIV Screening: Members at high risk of HIV	Health Benchmarks, Inc, IMS Health	To ensure that members at increased risk of HIV infection be screened for HIV