

# NATIONAL QUALITY FORUM

## Population Health: Prevention EM Submitted Measures

Measure ID/ Title	Measure Description	Measure Steward	Topic Area
<b>#0421</b> <b>Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up</b>	Percentage of patients aged 18 years and older with a calculated BMI in the past six months or during the current visit documented in the medical record AND if the most recent BMI is outside of normal parameters, a follow-up plan is documented Normal Parameters: Age 65 and older BMI > than or = to 23 and <30, Age 18 – 64 BMI > than or = to 18.5 and <25.	Centers for Medicare & Medicaid Services	Weight/ Physical Activity
<b>#0431</b> <b>Influenza Vaccination Coverage Among Healthcare Personnel</b>	Percentage of healthcare personnel (HCP) who receive the influenza vaccination.	Centers for Disease Control and Prevention	Immunization
<b>#0522</b> <b>Influenza Immunization Received for Current Flu Season</b>	Percentage of home health episodes of care during which patients received influenza immunization for the current flu season.	Centers for Medicare & Medicaid Services	Immunization
<b>#0525</b> <b>Pneumococcal Polysaccharide Vaccine (PPV) Ever Received</b>	Percentage of home health episodes of care during which patients were determined to have ever received Pneumococcal Polysaccharide Vaccine (PPV).	Centers for Medicare & Medicaid Services	Immunization
<b>#0023</b> <b>Body Mass Index (BMI) in adults &gt; 18 years of age</b>	Percentage of adults with BMI documentation in the past 24 months.	City of New York Dept. of Health and Mental Hygiene	Weight/ Physical Activity
<b>#0226</b> <b>Influenza Immunization in the ESRD Population (Facility Level)</b>	Percentage of end stage renal disease (ESRD) patients aged 6 months and older receiving hemodialysis or peritoneal dialysis during the time from October 1 (or when the influenza vaccine became available) to March 31 who either received, were offered and declined, or were determined to have a medical contraindication to the influenza vaccine.	Kidney Care Quality Alliance	Immunization
<b>#0614</b> <b>Steroid Use - Osteoporosis Screening</b>	The percentage of patients, 18 and older, who have been on chronic steroids for at least 180 days in the past 9 months and who had a bone density evaluation or osteoporosis treatment.	ActiveHealth Management	Screening
<b>#0617</b> <b>High Risk for Pneumococcal Disease - Pneumococcal Vaccination</b>	The percentage of patients age 5-64 with a high risk condition or age 65 years and older who received a pneumococcal vaccine.	ActiveHealth Management	Immunization
<b>#0629</b> <b>Male Smokers or Family History of Abdominal Aortic Aneurysm (AAA) - Consider Screening for AAA</b>	The percentage of men age 65-75 years with history of tobacco use or men age 60 yrs and older with a family history of abdominal aortic aneurysm who were screened for AAA.	ActiveHealth Management	Screening

Measure ID/ Title	Measure Description	Measure Steward	Topic Area
<b>#1653 Pneumococcal Immunization (PPV 23)</b>	Inpatients age 65 years and older and 6-64 years of age who have a high risk condition who are screened for 23-valent Pneumococcal Polysaccharide Vaccine (PPV23)status and vaccinated prior to discharge if indicated.	Centers for Medicare & Medicaid Services	Immunization
<b>#1659 Influenza Immunization</b>	Inpatients age 6 months and older discharged during October, November, December, January, February or March who are screened for influenza vaccine status and vaccinated prior to discharge if indicated.	Centers for Medicare & Medicaid Services	Immunization
<b>#0041 Influenza Immunization</b>	Percentage of patients aged 6 months and older seen for a visit between October 1 and the end of February of the one-year measurement period who have documented administration of influenza immunization OR patient reported previous receipt of an influenza immunization.	American Medical Association- Physician Consortium for Performance Improvement	Immunization
<b>#0579 Annual Cervical Cancer Screening for High-Risk Patients</b>	This measure identifies women age 12 to 65 diagnosed with cervical dysplasia (CIN 2), cervical carcinoma-in-situ, or HIV/AIDS prior to the measurement year, and who still have a cervix, who had a cervical CA screen during the measurement year.	Resolution Health, Inc.	Screening
<b>#1651 TOB-1 Tobacco Use Screening</b>	Hospitalized patients age 18 years and older who are screened during the hospital stay for tobacco use (cigarettes, smokeless tobacco, pipe and cigars) within the past 30 days. This measure is intended to be used as part of a set of 4 linked measures addressing Tobacco Use (TOB-2 Tobacco Use Treatment Provided or Offered (during the hospital stay); TOB-3 Tobacco Use Treatment Provided or offered at Discharge; TOB-4 Tobacco Use: Assessing Status After Discharge.)	The Joint Commission	Tobacco Use
<b>#1654 TOB - 2 Tobacco Use Treatment Provided or Offered and the subset measure TOB-2a Tobacco Use Treatment</b>	The measure is reported as an overall rate which includes all hospitalized patients 18 years of age and older to whom tobacco use treatment was provided during the hospital stay, or offered and refused, and a second rate, a subset of the first, which includes only those patients who received tobacco use treatment during the hospital stay. Refer to section 2a1.10 Stratification Details/Variables for the rationale for the addition of the subset measure. These measures are intended to be used as part of a set of 4 linked measures addressing Tobacco Use (TOB-1 Tobacco Use Screening; TOB-3 Tobacco Use Treatment Provided or Offered at Discharge; TOB-4 Tobacco Use: Assessing Status After Discharge.)	The Joint Commission	Tobacco Use

Measure ID/ Title	Measure Description	Measure Steward	Topic Area
<b>#1656</b> <b>TOB-3 Tobacco Use Treatment Provided or Offered at Discharge and the subset measure TOB-3a Tobacco Use Treatment at Discharge</b>	<p>The measure is reported as an overall rate which includes all hospitalized patients 18 years of age and older to whom tobacco use treatment was provided, or offered and refused, at the time of hospital discharge, and a second rate, a subset of the first, which includes only those patients who received tobacco use treatment at discharge. Treatment at discharge includes a referral to outpatient counseling and a prescription for one of the FDA-approved tobacco cessation medications. Refer to section 2a1.10 Stratification Details/Variables for the rationale for the addition of the subset measure. These measures are intended to be used as part of a set of 4 linked measures addressing Tobacco Use (TOB-1 Tobacco Use Screening; TOB-2 Tobacco Use Treatment Provided or Offered During the Hospital Stay; TOB-4 Tobacco Use: Assessing Status After Discharge).</p>	The Joint Commission	Tobacco Use
<b>#1657</b> <b>TOB-4 Tobacco Use: Assessing Status after Discharge</b>	<p>Hospitalized patients 18 years of age and older who are identified through the screening process as having used tobacco products (cigarettes, smokeless tobacco, pipe, and cigars) within the past 30 days who are contacted within 30 days after hospital discharge and follow-up information regarding tobacco use status is collected. This measure is intended to be used as part of a set of 4 linked measures addressing Tobacco Use (TOB-1 Tobacco Use Screening; TOB-2 Tobacco Use Treatment Provided or Offered (during hospital stay); TOB-3 Tobacco Use Treatment Provided or Offered at Discharge).</p>	The Joint Commission	Tobacco Use
<b>#1661</b> <b>SUB-1 Alcohol Use Screening</b>	<p>Hospitalized patients 18 years of age and older who are screened during the hospital stay using a validated screening questionnaire for unhealthy alcohol use. This measure is intended to be used as part of a set of 4 linked measures addressing Substance Use (SUB-1 Alcohol Use Screening ; SUB-2 Alcohol Use Brief Intervention Provided or Offered; SUB-3 Alcohol and Other Drug Use Disorder Treatment Provided or Offered at Discharge; SUB-4 Alcohol and Drug Use: Assessing Status after Discharge).</p>	The Joint Commission	Alcohol Use
<b>#1663</b> <b>SUB-2 Alcohol Use Brief Intervention Provided or Offered and SUB-2a Alcohol Use Brief Intervention</b>	<p>The measure is reported as an overall rate which includes all hospitalized patients 18 years of age and older to whom a brief intervention was provided, or offered and refused, and a second rate, a subset of the first, which includes only those patients who received a brief intervention. The Provided or Offered rate (SUB-2), describes patients who screened positive for unhealthy alcohol use who received or refused a brief intervention during the hospital stay. The Alcohol Use Brief Intervention (SUB-2a) rate describes only those who received the brief intervention during the hospital stay. Those who refused are not included.</p> <p>These measures are intended to be used as part of a set of 4 linked measures addressing Substance Use (SUB-1 Alcohol Use Screening ; SUB-2 Alcohol Use Brief Intervention Provided or Offered; SUB-3 Alcohol and Other Drug Use Disorder Treatment Provided or Offered at Discharge; SUB-4 Alcohol and Drug Use: Assessing Status after Discharge).</p>	The Joint Commission	Alcohol Use

Measure ID/ Title	Measure Description	Measure Steward	Topic Area
<b>#1664</b> <b>SUB-3 Alcohol &amp; Other Drug Use Disorder Treatment Provided or Offered at Discharge and SUB-3a Alcohol &amp; Other Drug Use Disorder Treatment at Discharge</b>	<p>The measure is reported as an overall rate which includes all hospitalized patients 18 years of age and older to whom alcohol or drug use disorder treatment was provided, or offered and refused, at the time of hospital discharge, and a second rate, a subset of the first, which includes only those patients who received alcohol or drug use disorder treatment at discharge. The Provided or Offered rate (SUB-3) describes patients who are identified with alcohol or drug use disorder who receive or refuse at discharge a prescription for FDA-approved medications for alcohol or drug use disorder, OR who receive or refuse a referral for addictions treatment. The Alcohol and Other Drug Disorder Treatment at Discharge (SUB-3a) rate describes only those who receive a prescription for FDA-approved medications for alcohol or drug use disorder OR a referral for addictions treatment. Those who refused are not included.</p> <p>These measures are intended to be used as part of a set of 4 linked measures addressing Substance Use (SUB-1 Alcohol Use Screening ; SUB-2 Alcohol Use Brief Intervention Provided or Offered; SUB-3 Alcohol and Other Drug Use Disorder Treatment Provided or Offered at Discharge; SUB-4 Alcohol and Drug Use: Assessing Status after Discharge).</p>	The Joint Commission	Alcohol Use
<b>#1665</b> <b>SUB-4 Alcohol &amp; Drug Use: Assessing Status After Discharge</b>	<p>Hospitalized patients age 18 years and older who screened positive for unhealthy alcohol use or who received a diagnosis of alcohol or drug disorder during their inpatient stay, who are contacted within 30 days after hospital discharge and follow-up information regarding their alcohol or drug use status post discharge is collected.</p> <p>This measure is intended to be used as part of a set of 4 linked measures addressing Substance Use (SUB-1 Alcohol Use Screening ; SUB-2 Alcohol Use Brief Intervention Provided or Offered; SUB-3 Alcohol and Other Drug Use Disorder Treatment Provided or Offered at Discharge; SUB-4 Alcohol and Drug Use: Assessing Status after Discharge).</p>	The Joint Commission	Alcohol Use
<b>#0024</b> <b>Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents</b>	<p>Percentage of children 3-17 years of age who had an outpatient visit with a primary care physician or an OB/GYN and who had evidence of body mass index (BMI) percentile documentation, counseling for nutrition and counseling for physical activity during the measurement year.</p>	National Committee for Quality Assurance	Weight/ Physical Activity

Measure ID/ Title	Measure Description	Measure Steward	Topic Area
<b>#0027</b> <b>Medical Assistance With Smoking Cessation</b>	<p>Assesses different facets of providing medical assistance with smoking and tobacco use cessation:</p> <p>-Advising Smokers and Tobacco Users to Quit: A rolling average represents the percentage of members 18 years of age and older who were current smokers or tobacco users and who received advice to quite during the measurement year.</p> <p>-Discussing Cessation Medications: A rolling average represents the percentage of members 18 years of age and older who were current smokers or tobacco users and who discussed or were recommended cessation medications during the measurement year.</p> <p>-Discussing Cessation Strategies: A rolling average represents the percentage of members 18 years of age and older who were current smokers or tobacco users and who discussed or were provided smoking cessation methods or strategies during the measurement year.</p>	National Committee for Quality Assurance	Tobacco Use
<b>#0029</b> <b>Counseling on physical activity in older adults - a. Discussing Physical Activity, b. Advising Physical Activity</b>	<p>Percentage patients 65 years of age and older who reported: discussing their level of exercise or physical activity with a doctor or other health provider in the last 12 months.</p> <p>Percentage patients 65 years of age and older who reported receiving advice to start, increase, or maintain their level of exercise or physical activity from a doctor or other health provider in the last 12 months.</p>	National Committee for Quality Assurance	Weight/ Physical Activity
<b>#0032</b> <b>Cervical Cancer Screening</b>	<p>Percentage of women 21–64 years of age received one or more Pap tests to screen for cervical cancer.</p>	National Committee for Quality Assurance	Screening
<b>#0033</b> <b>Chlamydia screening in women</b>	<p>Assesses the percentage of women 16–24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.</p>	National Committee for Quality Assurance	Screening
<b>#0034</b> <b>Colorectal Cancer Screening</b>	<p>The percentage of members 50–75 years of age who had appropriate screening for colorectal cancer.</p>	National Committee for Quality Assurance	Screening
<b>#0037</b> <b>Osteoporosis testing in older women</b>	<p>Percentage of female patients aged 65 and older who reported receiving a bone density test (BMD) to check for osteoporosis.</p>	National Committee for Quality Assurance	Screening
<b>#0038</b> <b>Childhood Immunization Status</b>	<p>Percentage of children 2 years of age who had four diphtheria, tetanus and acellular pertussis (DtaP); three polio (IPV); one measles, mumps and rubella (MMR); three H influenza type B (HiB); three hepatitis B (HepB); one chicken pox (VZV); four pneumococcal conjugate (PCV); two hepatitis A (HepA); two or three rotavirus (RV); and two influenza (flu) vaccines by their second birthday. The measure calculates a rate for each vaccine and nine separate combination rates.</p>	National Committee for Quality Assurance	Immunization

Measure ID/ Title	Measure Description	Measure Steward	Topic Area
<b>#0039</b> <b>Flu Shots for Adults Ages 50 and Over</b>	<p>This measure represents the percentage of adults aged 50 and over who received an influenza vaccine within the measurement period within the respective age-stratified CAHPS surveys. This measure is only reported by age group stratification. The terms FSA and FSO, defined below, will be used to identify any differences between the two age stratifications.</p> <p>FSA - A rolling average represents the percentage of members 50–64 years of age who received an influenza vaccination between September 1 of the measurement year and the date on which the CAHPS 4.0H adult survey was completed.</p> <p>FSO - The percentage of Medicare members 65 years of age and older who received an influenza vaccination between September 1 of the measurement year and the date on which the Medicare CAHPS survey was completed.</p>	National Committee for Quality Assurance	Immunization
<b>#0043</b> <b>Pneumonia vaccination status for older adults</b>	Percentage of patients 65 years of age and older who ever received a pneumococcal vaccination.	National Committee for Quality Assurance	Immunization
<b>#0046</b> <b>Osteoporosis: Screening or Therapy for Women Aged 65 Years and Older</b>	Percentage of female patients aged 65 years and older who have a central DXA measurement ordered or performed at least once since age 60 or pharmacologic therapy prescribed within 12 months.	National Committee for Quality Assurance	Screening
<b>#1690</b> <b>Adult BMI Assessment</b>	The percentage of adults 18–74 years of age who had body mass index (BMI) documented.	National Committee for Quality Assurance	Weight/ Physical Activity
<b>#0031</b> <b>Breast Cancer Screening</b>	Percentage of eligible women 40-69 who receive a mammogram in a two year period.	National Committee for Quality Assurance	Screening