

NATIONAL QUALITY FORUM

Measure Evaluation 4.1 December 2009

This form contains the measure information submitted by stewards. Blank fields indicate no information was provided. Attachments also may have been submitted and are provided to reviewers. The subcriteria and most of the footnotes from the [evaluation criteria](#) are provided in Word comments within the form and will appear if your cursor is over the highlighted area. Hyperlinks to the evaluation criteria and ratings are provided in each section.

TAP/Workgroup (if utilized): Complete all **yellow highlighted** areas of the form. Evaluate the extent to which each subcriterion is met. Based on your evaluation, summarize the strengths and weaknesses in each section.

Note: *If there is no TAP or workgroup, the SC also evaluates the subcriteria (yellow highlighted areas).*

Steering Committee: Complete all **pink** highlighted areas of the form. Review the workgroup/TAP assessment of the subcriteria, noting any areas of disagreement; then evaluate the extent to which each major criterion is met; and finally, indicate your recommendation for the endorsement. Provide the rationale for your ratings.

Evaluation ratings of the extent to which the criteria are met

C = Completely (unquestionably demonstrated to meet the criterion)

P = Partially (demonstrated to partially meet the criterion)

M = Minimally (addressed BUT demonstrated to only minimally meet the criterion)

N = Not at all (NOT addressed; OR incorrectly addressed; OR demonstrated to NOT meet the criterion)

NA = Not applicable (only an option for a few subcriteria as indicated)

(for NQF staff use) NQF Review #: 0125	NQF Project: Surgery Endorsement Maintenance 2010
MEASURE DESCRIPTIVE INFORMATION	
De.1 Measure Title: Timing of Antibiotic Prophylaxis for Cardiac Surgery Patients	
De.2 Brief description of measure: Percent of patients aged 18 years and older undergoing cardiac surgery who received prophylactic antibiotics within one hour of surgical incision or start of procedure if no incision was required (two hours if receiving vancomycin or fluoroquinolone)	
1.1-2 Type of Measure: Process	
De.3 If included in a composite or paired with another measure, please identify composite or paired measure	
De.4 National Priority Partners Priority Area: Safety	
De.5 IOM Quality Domain: Safety	
De.6 Consumer Care Need: Getting better	

CONDITIONS FOR CONSIDERATION BY NQF	
Four conditions must be met before proposed measures may be considered and evaluated for suitability as voluntary consensus standards:	NQF Staff
<p>A. The measure is in the public domain or an intellectual property (measure steward agreement) is signed. <i>Public domain only applies to governmental organizations. All non-government organizations must sign a measure steward agreement even if measures are made publicly and freely available.</i></p> <p>A.1 Do you attest that the measure steward holds intellectual property rights to the measure and the right to use aspects of the measure owned by another entity (e.g., risk model, code set)? Yes</p> <p>A.2 Indicate if Proprietary Measure (as defined in measure steward agreement):</p> <p>A.3 Measure Steward Agreement: Agreement will be signed and submitted prior to or at the time of measure submission</p> <p>A.4 Measure Steward Agreement attached: STS Measure Steward Agreement. Fully Executed-634267323027557342.pdf</p>	<p>A</p> <p>Y <input type="checkbox"/></p> <p>N <input type="checkbox"/></p>

<p>B. The measure owner/steward verifies there is an identified responsible entity and process to maintain and update the measure on a schedule that is commensurate with the rate of clinical innovation, but at least every 3 years. Yes, information provided in contact section</p>	<p>B Y <input type="checkbox"/> N <input type="checkbox"/></p>
<p>C. The intended use of the measure includes <u>both</u> public reporting <u>and</u> quality improvement. ► Purpose: Public Reporting, Quality Improvement (Internal to the specific organization), Quality Improvement with Benchmarking (external benchmarking to multiple organizations)</p>	<p>C Y <input type="checkbox"/> N <input type="checkbox"/></p>
<p>D. The requested measure submission information is complete. Generally, measures should be fully developed and tested so that all the evaluation criteria have been addressed and information needed to evaluate the measure is provided. Measures that have not been tested are only potentially eligible for a time-limited endorsement and in that case, measure owners must verify that testing will be completed within 12 months of endorsement. D.1 Testing: Yes, fully developed and tested D.2 Have NQF-endorsed measures been reviewed to identify if there are similar or related measures? Yes</p>	<p>D Y <input type="checkbox"/> N <input type="checkbox"/></p>
<p>(for NQF staff use) Have all conditions for consideration been met? Staff Notes to Steward (if submission returned):</p>	<p>Met Y <input type="checkbox"/> N <input type="checkbox"/></p>
<p>Staff Notes to Reviewers (issues or questions regarding any criteria):</p>	
<p>Staff Reviewer Name(s):</p>	

<p>TAP/Workgroup Reviewer Name:</p>	
<p>Steering Committee Reviewer Name:</p>	
<p>1. IMPORTANCE TO MEASURE AND REPORT</p>	
<p>Extent to which the specific measure focus is important to making significant gains in health care quality (safety, timeliness, effectiveness, efficiency, equity, patient-centeredness) and improving health outcomes for a specific high impact aspect of healthcare where there is variation in or overall poor performance. Measures must be judged to be important to measure and report in order to be evaluated against the remaining criteria. (evaluation criteria) 1a. High Impact</p>	<p>Eval Rating</p>
<p>(for NQF staff use) Specific NPP goal:</p>	
<p>1a.1 Demonstrated High Impact Aspect of Healthcare: Affects large numbers, Frequently performed procedure, Leading cause of morbidity/mortality, High resource use, Severity of illness, Patient/societal consequences of poor quality 1a.2 1a.3 Summary of Evidence of High Impact: Postoperative mediastinitis is an infection of the mediastinal space after cardiac surgery. The incidence of deep sternal infections (mediastinitis) associated with cardiac surgery ranges between 0.25% and 4% [1]. The incidence of postoperative mediastinitis can be decrease by assuring that “patients aged 18 years and older undergoing cardiac surgery receive prophylactic antibiotics within one hour of surgical incision or start of procedure if no incision was required (two hours if receiving vancomycin or fluoroquinolone)”. Reference 1 below states: “Postoperative mediastinitis carries a very high hospital mortality [3-5] and is also associated with reduced long-term survival [3]. This complication invariably involves an additional operation, a prolonged hospitalization, a significant toll in clinical resources, and dramatically increased costs. Anyone who has provided care for a patient with mediastinitis also knows well the emotional cost not only for the patient but also for the family, the nursing staff, and the surgeons. Truly one of the most devastating infections in all of surgery, this dreaded complication influences the perioperative management strategy of virtually all cardiothoracic surgeons.”</p>	<p>1a C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>

1a.4 Citations for Evidence of High Impact: 1. Edwards FH, Engelman RM, Houck P, Shahian DM, Bridges CR; Society of Thoracic Surgeons. The Society of Thoracic Surgeons Practice Guideline Series: Antibiotic Prophylaxis in Cardiac Surgery, Part I: Duration. Ann Thorac Surg. 2006 Jan;81(1):397-404. No abstract available. PMID: 16368422

2. Engelman R, Shahian D, Shemin R, Guy TS, Bratzler D, Edwards F, Jacobs M, Fernando H, Bridges C; Workforce on Evidence-Based Medicine, Society of Thoracic Surgeons. The Society of Thoracic Surgeons practice guideline series: Antibiotic prophylaxis in cardiac surgery, part II: Antibiotic choice. Ann Thorac Surg. 2007 Apr;83(4):1569-76. Review. No abstract available. PMID: 17383396

3. Braxton JH, Marrin CAS, McGrath PD, et al. 10-year follow-up of patients with and without mediastinitis. Sem Thorac Cardiovasc Surg 2004;16:70-6.

4. Demmy TL, Park SB, Liebler GA, et al. Recent experience with major sternal wound complications. Ann Thorac Surg 1990;49:458-62.

5. Tang GHL, Maganti M, Weisel RD, Borger MA. Prevention and management of deep sternal wound infection. Sem Thorac Cardiovasc Surg 2004;16:62-9.

6. American Society of Health-System Pharmacists. ASHP Therapeutic Guidelines on Antimicrobial Prophylaxis in Surgery; March 23, 2004. Available at www.ashp.org. Last accessed April 20, 2004.

7. Centers for Disease Control and Prevention (CDC) National Nosocomial Infections Surveillance (NNIS) System. National nosocomial infections surveillance (NNIS) system report, data summary from January 1992 to June 2003, issued August 2003. Am J Infect Control. 2003;31:481-498.

8. Classen DC, Evans RS, Pestotnik SL, Horn SD, Menlove RL, Burke JP. The timing of prophylactic administration of antibiotics and the risk of surgical-wound infection. N Engl J Med. 1992;326(5):281-286.

1b. Opportunity for Improvement

1b.1 Benefits (improvements in quality) envisioned by use of this measure: The incidence of deep sternal infections (mediastinitis) associated with cardiac surgery ranges between 0.25% and 4% [1]. The incidence of postoperative mediastinitis can be decrease by assuring that “patients aged 18 years and older undergoing cardiac surgery who received prophylactic antibiotics within one hour of surgical incision or start of procedure if no incision was required (two hours if receiving vancomycin or fluoroquinolone)”.

1b.2 Summary of data demonstrating performance gap (variation or overall poor performance) across providers:

Please see attachment and below

Measurement Timing of Antibiotic Administration for Cardiac Surgery Patients

N	786
Mean	98.0%
1st	83.2%
5th	93.2%
10th	95.2%
25th	97.7%
Median	99.2%
75th	99.9%
90th	100.0%
95th	100.0%
99th	100.0%

Outlier 347 (44.1%)

High 259

Low 88

1b.3 Citations for data on performance gap:

Dates: January 1, 2009-December 31, 2009

Analysis includes 786 STS Adult Cardiac Surgery Database Participants who had at least 100 eligible cases for the measure and reported data to STS for all 12 months.

1b.4 Summary of Data on disparities by population group:

1b
 C
 P
 M
 N

please see attachment

1b.5 Citations for data on Disparities:

Analysis includes STS Adult Cardiac Surgery Database Participants that had more than 50 eligible cases in 2008 and 2009, and reported data for at least 15 months.

- 375888 Patients from 887 Participants were included in the Gender = Male sub-group.
- 175058 Patients from 819 Participants were included in the Gender = Female sub-group.
- 29844 Patients from 231 Participants were included in the Race = Black sub-group.
- 477888 Patients from 881 Participants were included in the Race = White sub-group.
- 25994 Patients from 192 Participants were included in the Race = Other sub-group.
- 19142 Patients from 151 Participants were included in the Ethnicity = Hispanic sub-group.
- 526816 Patients from 887 Participants were included in the Ethnicity = Non-Hispanic sub-group.

1c. Outcome or Evidence to Support Measure Focus

1c.1 Relationship to Outcomes (*For non-outcome measures, briefly describe the relationship to desired outcome. For outcomes, describe why it is relevant to the target population*): “Postoperative mediastinitis carries a very high hospital mortality and is also associated with reduced long-term survival [3]. This complication invariably involves an additional operation, a prolonged hospitalization, a significant toll in clinical resources, and dramatically increased costs. Anyone who has provided care for a patient with mediastinitis also knows well the emotional cost not only for the patient but also for the family, the nursing staff, and the surgeons. Truly one of the most devastating infections in all of surgery, this dreaded complication influences the perioperative management strategy of virtually all cardiothoracic surgeons.”

Reference:

Edwards FH, Engelman RM, Houck P, Shahian DM, Bridges CR; Society of Thoracic Surgeons. The Society of Thoracic Surgeons Practice Guideline Series: Antibiotic Prophylaxis in Cardiac Surgery, Part I: Duration. *Ann Thorac Surg.* 2006 Jan;81(1):397-404. No abstract available. PMID: 16368422

The incidence of deep sternal infections (mediastinitis) associated with cardiac surgery ranges between 0.25% and 4% [1]. The incidence of postoperative mediastinitis can be decrease by assuring that “patients aged 18 years and older undergoing cardiac surgery receive prophylactic antibiotics within one hour of surgical incision or start of procedure if no incision was required (two hours if receiving vancomycin or fluoroquinolone)”.

1c.2-3. Type of Evidence: Observational study, Expert opinion, Systematic synthesis of research, Other Clinical results from approximately 90% of cardiac surgery centers in the US

1c.4 Summary of Evidence (*as described in the criteria; for outcomes, summarize any evidence that healthcare services/care processes influence the outcome*):

“Postoperative mediastinitis carries a very high hospital mortality and is also associated with reduced long-term survival [3]. This complication invariably involves an additional operation, a prolonged hospitalization, a significant toll in clinical resources, and dramatically increased costs. Anyone who has provided care for a patient with mediastinitis also knows well the emotional cost not only for the patient but also for the family, the nursing staff, and the surgeons. Truly one of the most devastating infections in all of surgery, this dreaded complication influences the perioperative management strategy of virtually all cardiothoracic surgeons.”

Reference:

Edwards FH, Engelman RM, Houck P, Shahian DM, Bridges CR; Society of Thoracic Surgeons. The Society of Thoracic Surgeons Practice Guideline Series: Antibiotic Prophylaxis in Cardiac Surgery, Part I: Duration. *Ann Thorac Surg.* 2006 Jan;81(1):397-404. No abstract available. PMID: 16368422

The incidence of deep sternal infections (mediastinitis) associated with cardiac surgery ranges between 0.25% and 4% [1]. The incidence of postoperative mediastinitis can be decreased by assuring that “patients aged 18 years and older undergoing cardiac surgery receive prophylactic antibiotics within one hour of

1c
C
P
M
N

surgical incision or start of procedure if no incision was required (two hours if receiving vancomycin or fluoroquinolone)”.

“In patients for whom cefazolin is the appropriate prophylactic antibiotic for cardiac surgery, administration within 60 minutes of the skin incision is indicated (Class I, Level of Evidence A).”

Reference:

Engelman R, Shahian D, Shemin R, Guy TS, Bratzler D, Edwards F, Jacobs M, Fernando H, Bridges C; Workforce on Evidence-Based Medicine, Society of Thoracic Surgeons. The Society of Thoracic Surgeons practice guideline series: Antibiotic prophylaxis in cardiac surgery, part II: Antibiotic choice. *Ann Thorac Surg*. 2007 Apr;83(4):1569-76. Review. No abstract available. PMID: 17383396

“In patients for whom vancomycin is an appropriate prophylactic antibiotic for cardiac surgery, a dose of 1 to 1.5 g or a weight-adjusted dose of 15 mg/kg administered intravenously slowly over 1 hour, with completion within 1 hour of the skin incision, is recommended (Class I, Level of Evidence A).”

Reference:

Engelman R, Shahian D, Shemin R, Guy TS, Bratzler D, Edwards F, Jacobs M, Fernando H, Bridges C; Workforce on Evidence-Based Medicine, Society of Thoracic Surgeons. The Society of Thoracic Surgeons practice guideline series: Antibiotic prophylaxis in cardiac surgery, part II: Antibiotic choice. *Ann Thorac Surg*. 2007 Apr;83(4):1569-76. Review. No abstract available. PMID: 17383396

1c.5 Rating of strength/quality of evidence (also provide narrative description of the rating and by whom):

Class I, Level of Evidence A - “In patients for whom cefazolin is the appropriate prophylactic antibiotic for cardiac surgery, administration within 60 minutes of the skin incision is indicated.” Class I, Level of Evidence A - “In patients for whom vancomycin is an appropriate prophylactic antibiotic for cardiac surgery, a dose of 1 to 1.5 g or a weight-adjusted dose of 15 mg/kg administered intravenously slowly over 1 hour, with completion within 1 hour of the skin incision, is recommended.” -- Reference: Engelman R, Shahian D, Shemin R, Guy TS, Bratzler D, Edwards F, Jacobs M, Fernando H, Bridges C; Workforce on Evidence-Based Medicine, Society of Thoracic Surgeons. The Society of Thoracic Surgeons practice guideline series: Antibiotic prophylaxis in cardiac surgery, part II: Antibiotic choice. *Ann Thorac Surg*. 2007 Apr;83(4):1569-76. Review. No abstract available. PMID: 17383396

1c.6 Method for rating evidence: n/a

1c.7 Summary of Controversy/Contradictory Evidence: n/a

- 1c.8 Citations for Evidence (other than guidelines):**
1. Edwards FH, Engelman RM, Houck P, Shahian DM, Bridges CR; Society of Thoracic Surgeons. The Society of Thoracic Surgeons Practice Guideline Series: Antibiotic Prophylaxis in Cardiac Surgery, Part I: Duration. *Ann Thorac Surg*. 2006 Jan;81(1):397-404. No abstract available. PMID: 16368422
 2. Engelman R, Shahian D, Shemin R, Guy TS, Bratzler D, Edwards F, Jacobs M, Fernando H, Bridges C; Workforce on Evidence-Based Medicine, Society of Thoracic Surgeons. The Society of Thoracic Surgeons practice guideline series: Antibiotic prophylaxis in cardiac surgery, part II: Antibiotic choice. *Ann Thorac Surg*. 2007 Apr;83(4):1569-76. Review. No abstract available. PMID: 17383396
 3. Braxton JH, Marrin CAS, McGrath PD, et al. 10-year follow-up of patients with and without mediastinitis. *Sem Thorac Cardiovasc Surg* 2004;16:70-6.
 4. Demmy TL, Park SB, Liebler GA, et al. Recent experience with major sternal wound complications. *Ann Thorac Surg* 1990;49:458-62.
 5. Tang GHL, Maganti M, Weisel RD, Borger MA. Prevention and management of deep sternal wound infection. *Sem Thorac Cardiovasc Surg* 2004;16:62-9.
 6. American Society of Health-System Pharmacists. ASHP Therapeutic Guidelines on Antimicrobial Prophylaxis in Surgery; March 23, 2004. Available at www.ashp.org. Last accessed April 20, 2004.
 7. Centers for Disease Control and Prevention (CDC) National Nosocomial Infections Surveillance (NNIS) System. National nosocomial infections surveillance (NNIS) system report, data summary from January 1992 to June 2003, issued August 2003. *Am J Infect Control*. 2003;31:481-498.

<p>8. Classen DC, Evans RS, Pestotnik SL, Horn SD, Menlove RL, Burke JP. The timing of prophylactic administration of antibiotics and the risk of surgical-wound infection. N Engl J Med. 1992;326(5):281-286.</p> <p>1c.9 Quote the Specific guideline recommendation (including guideline number and/or page number): n/a</p> <p>1c.10 Clinical Practice Guideline Citation: n/a</p> <p>1c.11 National Guideline Clearinghouse or other URL: n/a</p> <p>1c.12 Rating of strength of recommendation (also provide narrative description of the rating and by whom): n/a</p> <p>1c.13 Method for rating strength of recommendation (If different from USPSTF system, also describe rating and how it relates to USPSTF): n/a</p> <p>1c.14 Rationale for using this guideline over others: n/a</p>	
<p>TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for <i>Importance to Measure and Report</i>?</p>	<p>1</p>
<p>Steering Committee: Was the threshold criterion, <i>Importance to Measure and Report</i>, met? Rationale:</p>	<p>1 Y <input type="checkbox"/> N <input type="checkbox"/></p>
<p style="text-align: center;">2. SCIENTIFIC ACCEPTABILITY OF MEASURE PROPERTIES</p>	
<p>Extent to which the measure, <u>as specified</u>, produces consistent (reliable) and credible (valid) results about the quality of care when implemented. (evaluation criteria)</p>	<p>Eval Rating</p>
<p style="text-align: center;">2a. MEASURE SPECIFICATIONS</p>	
<p>S.1 Do you have a web page where current detailed measure specifications can be obtained? S.2 If yes, provide web page URL:</p>	
<p>2a. Precisely Specified</p>	
<p>2a.1 Numerator Statement (Brief, text description of the numerator - what is being measured about the target population, e.g. target condition, event, or outcome): Number of patients undergoing cardiac surgery patients who received prophylactic antibiotics within one hour of surgical incision or start of procedure if no incision was required (two hours if vancomycin or fluoroquinolone)</p>	
<p>2a.2 Numerator Time Window (The time period in which cases are eligible for inclusion in the numerator): Within one hour of surgical incision or start of procedure if no incision was required (two hours if vancomycin or fluoroquinolone)</p>	
<p>Rationale: Due to the longer infusion time required for vancomycin or a fluoroquinolone, it is acceptable to start these antibiotics within two hours prior to incision time.</p>	
<p>2a.3 Numerator Details (All information required to collect/calculate the numerator, including all codes, logic, and definitions): Number of cardiac surgery procedures in which timing of appropriate antibiotic administration [AbxTiming (STS Adult Cardiac Surgery Database Version 2.73)] is marked “yes”</p>	<p>2a-specs C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>
<p>2a.4 Denominator Statement (Brief, text description of the denominator - target population being measured): Number of patients undergoing cardiac surgery</p>	

2a.5 Target population gender: Female, Male
2a.6 Target population age range: 18 and older

2a.7 Denominator Time Window (*The time period in which cases are eligible for inclusion in the denominator*):
 12 months

2a.8 Denominator Details (*All information required to collect/calculate the denominator - the target population being measured - including all codes, logic, and definitions*):
 Number of cardiac surgery procedures;

A cardiac procedure is determined as a procedure for which at least one of the following is not marked “no” or “missing” (note: full terms for STS field names are provided in brackets []):
 OpCAB[Coronary Artery Bypass], OpValve[Valve Surgery], VADProc [VAD Implanted or Removed], VSAV [Aortic Valve Procedure], VSMV [Mitral Valve Procedure], OpTricus [Tricuspid Valve Procedure Performed], OpPulm[Pulmonic Valve Procedure Performed], OpOCard [Other Cardiac Procedure other than CABG or Valve], OCarLVA [Left Ventricular Aneurysm Repair], OCarVSD [Ventricular Septal Defect Repair], OCarSVR [Surgical Ventricular Restoration], OCarCong [Congenital Defect Repair], OCarTrma [surgical procedure for an injury due to Cardiac Trauma], OCarCrTx [Cardiac Transplant], OCarACD [Arrhythmia Correction Surgery], OCAoProcType[Aortic Procedure Type], EndoProc [Endovascular Procedure (TEVAR)], OCTumor [resection of an intracardiac tumor], OCPulThromDis [Pulmonary Thromboembolism], OCarOthr [Other Cardiac Procedure other than those listed previously], ECMO [Extracorporeal Membrane Oxygenation], OCarLasr [-Transmyocardial Laser Revascularization], OCarASD [Atrial Septal Defect Repair], OCarAFibSur [Atrial Fibrillation Surgical Procedure]

2a.9 Denominator Exclusions (*Brief text description of exclusions from the target population*): Cases are removed from the denominator if the patient had a documented contraindication or rationale for not administering antibiotic in medical record.

Other exclusions include:

- Patients who had a principal diagnosis suggestive of preoperative infectious diseases
- Patients whose ICD-9-CM principal procedure was performed entirely by Laparoscope
- Patients enrolled in clinical trials
- Patients with documented infection prior to surgical procedure of interest
- Patients who were receiving antibiotics more than 24 hours prior to surgery
- Patients who were receiving antibiotics within 24 hours prior to arrival

This list will be provided in the STS Adult Cardiac Surgery Database Data Manager’s Training Manual as acceptable exclusions.

2a.10 Denominator Exclusion Details (*All information required to collect exclusions to the denominator, including all codes, logic, and definitions*):
 Timing of appropriate antibiotic administration (AbxTiming) is marked “Exclusion”

2a.11 Stratification Details/Variables (*All information required to stratify the measure including the stratification variables, all codes, logic, and definitions*):
 N/A

2a.12-13 Risk Adjustment Type: No risk adjustment necessary

2a.14 Risk Adjustment Methodology/Variables (*List risk adjustment variables and describe conceptual models, statistical models, or other aspects of model or method*):
 N/A

2a.15-17 Detailed risk model available Web page URL or attachment:

2a.18-19 Type of Score: Rate/proportion

2a.20 Interpretation of Score: Better quality = Higher score

2a.21 Calculation Algorithm (*Describe the calculation of the measure as a flowchart or series of steps*):

<p>N/A</p>	
<p>2a.22 Describe the method for discriminating performance (e.g., significance testing): Two-sided 95% binomial confidence intervals; a confidence interval is calculated for each database participant. If the overall STS database result falls within the participant’s 95% binomial confidence interval, the participant’s performance is considered not significantly different from the overall database result. If the overall STS database result falls to the right of the participant’s 95% binomial confidence interval, then the participant’s performance is considered significantly lower than the overall database results. If the overall STS database result falls to the left of the participant’s 95% binomial confidence interval, then the participant’s performance is considered significantly higher than the overall database results.</p>	
<p>2a.23 Sampling (Survey) Methodology <i>If measure is based on a sample (or survey), provide instructions for obtaining the sample, conducting the survey and guidance on minimum sample size (response rate):</i> N/A</p>	
<p>2a.24 Data Source <i>(Check the source(s) for which the measure is specified and tested)</i> Electronic Clinical Data : Registry</p>	
<p>2a.25 Data source/data collection instrument <i>(Identify the specific data source/data collection instrument, e.g. name of database, clinical registry, collection instrument, etc.):</i> STS Adult Cardiac Surgery Database - Version 2.73</p>	
<p>2a.26-28 Data source/data collection instrument reference web page URL or attachment: URL Data Collection Form http://www.sts.org/sites/default/files/documents/STSAultCVDataCollectionForm2_73_Annotated.pdf</p>	
<p>2a.29-31 Data dictionary/code table web page URL or attachment: URL http://www.sts.org/sites/default/files/documents/STSAultCVDataSpecificationsV2_73.pdf</p>	
<p>2a.32-35 Level of Measurement/Analysis <i>(Check the level(s) for which the measure is specified and tested)</i> Clinician : Group/Practice, Facility, Population : County or City, Population : National, Population : Regional, Population : State</p>	
<p>2a.36-37 Care Settings <i>(Check the setting(s) for which the measure is specified and tested)</i> Hospital/Acute Care Facility</p>	
<p>2a.38-41 Clinical Services <i>(Healthcare services being measured, check all that apply)</i> Clinicians: Physicians (MD/DO)</p>	
TESTING/ANALYSIS	
<p>2b. Reliability testing</p>	
<p>2b.1 Data/sample <i>(description of data/sample and size):</i> STS Adult Cardiac Surgery Database - Compared results between two proximate time periods: January 2008-December 2008 and January 2009-December 2009.</p>	
<p>2b.2 Analytic Method <i>(type of reliability & rationale, method for testing):</i> Compared results between two proximate time periods: January 2008-December 2008 and January 2009-December 2009. Excluded from analysis are participants that did not submit results for both time periods. As database participants can change their underlying care processes at any time, we would not expect perfect correlation between two sets of results from even proximate time periods.</p>	<p>2b C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>
<p>2b.3 Testing Results <i>(reliability statistics, assessment of adequacy in the context of norms for the test conducted):</i> Please see attachment</p>	
<p>2c. Validity testing</p>	<p>2c C <input type="checkbox"/></p>

<p>2c.1 Data/sample (<i>description of data/sample and size</i>): STS Adult Cardiac Surgery Database</p> <p>Audits conducted in 2010, all cases performed in 2009; N = 40 randomly selected sites participating in the STS Adult Cardiac Surgery Database</p> <p>2c.2 Analytic Method (<i>type of validity & rationale, method for testing</i>): Participating sites are randomly selected for participation in STS Adult Cardiac Surgery Database Audit, which is designed to evaluate the accuracy, consistency, and comprehensiveness of data collection and ultimately validate the integrity of the data contained in the database. The Iowa Foundation for Medical Care (IFMC), the quality improvement organization for Iowa and Illinois, has conducted audits on behalf of STS since 2006.</p> <p>Each year, the IFMC conducts audits at randomly selected sites throughout the country and tracks the individual agreement rates by variable and by year. More specifically, for each site, agreement rates are calculated for 73 individual elements. In addition, aggregate agreement rates for each element, variable category (e.g., pre-operative risk factors, previous interventions, etc), and overall for all categories are calculated for all sites. While this is not region specific, it is data point specific and comparison agreement rates confirm the improvement over time as well as the consistency.</p> <p>2c.3 Testing Results (<i>statistical results, assessment of adequacy in the context of norms for the test conducted</i>):</p>	<p>P <input type="checkbox"/></p> <p>M <input type="checkbox"/></p> <p>N <input type="checkbox"/></p>
<p>2d. Exclusions Justified</p> <p>2d.1 Summary of Evidence supporting exclusion(s):</p> <p>2d.2 Citations for Evidence:</p> <p>2d.3 Data/sample (<i>description of data/sample and size</i>): Immediately prior to this NQF measure endorsement maintenance period, stewardship of this measure was transferred to STS. Exclusions could not be captured using the previous version of the STS Database (STS Adult Cardiac Surgery Database Version 2.61).</p> <p>Released in December 2010, STS Adult Cardiac Surgery Database Version 2.73, which is designed to address changes in technology and practice, allow for easier identification of devices, and permit improved capture of preoperative risk factors, operative information and postoperative evaluation, has the capability of capturing exclusions data for this measure. Therefore, during the next NQF endorsement maintenance period, scheduled to take place in the year 2013, STS will be able to provide data on exclusions. STS Adult Cardiac Surgery Database Version 2.73 will be implemented for all cases with a surgery date of 7/1/2011 or later.</p> <p>2d.4 Analytic Method (<i>type analysis & rationale</i>):</p> <p>2d.5 Testing Results (<i>e.g., frequency, variability, sensitivity analyses</i>):</p>	<p>2d</p> <p>C <input type="checkbox"/></p> <p>P <input type="checkbox"/></p> <p>M <input type="checkbox"/></p> <p>N <input type="checkbox"/></p> <p>NA <input type="checkbox"/></p>
<p>2e. Risk Adjustment for Outcomes/ Resource Use Measures</p> <p>2e.1 Data/sample (<i>description of data/sample and size</i>): n/a</p> <p>2e.2 Analytic Method (<i>type of risk adjustment, analysis, & rationale</i>):</p> <p>2e.3 Testing Results (<i>risk model performance metrics</i>):</p>	<p>2e</p> <p>C <input type="checkbox"/></p> <p>P <input type="checkbox"/></p> <p>M <input type="checkbox"/></p> <p>N <input type="checkbox"/></p> <p>NA <input type="checkbox"/></p>

<p>2e.4 If outcome or resource use measure is not risk adjusted, provide rationale:</p>	
<p>2f. Identification of Meaningful Differences in Performance</p> <p>2f.1 Data/sample from Testing or Current Use (<i>description of data/sample and size</i>): 786 STS Adult Cardiac Surgery Database Participants who had at least 100 eligible cases for the measure and reported data to STS for all 12 months; January 1, 2009-December 31, 2009</p> <p>2f.2 Methods to identify statistically significant and practically/meaningfully differences in performance (<i>type of analysis & rationale</i>): Two-sided 95% binomial confidence intervals; a confidence interval is calculated for each database participant. If the overall STS database result falls within the participant’s 95% binomial confidence interval, the participant’s performance is considered not significantly different from the overall database result. If the overall STS database result falls to the right of the participant’s 95% binomial confidence interval, then the participant’s performance is considered significantly lower than the overall database results. If the overall STS database result falls to the left of the participant’s 95% binomial confidence interval, then the participant’s performance is considered significantly higher than the overall database results.</p> <p>2f.3 Provide Measure Scores from Testing or Current Use (<i>description of scores, e.g., distribution by quartile, mean, median, SD, etc.; identification of statistically significant and meaningfully differences in performance</i>): Please see attachment</p>	<p>2f C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>
<p>2g. Comparability of Multiple Data Sources/Methods</p> <p>2g.1 Data/sample (<i>description of data/sample and size</i>): n/a</p> <p>2g.2 Analytic Method (<i>type of analysis & rationale</i>):</p> <p>2g.3 Testing Results (<i>e.g., correlation statistics, comparison of rankings</i>):</p>	<p>2g C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/></p>
<p>2h. Disparities in Care</p> <p>2h.1 If measure is stratified, provide stratified results (<i>scores by stratified categories/cohorts</i>): n/a</p> <p>2h.2 If disparities have been reported/identified, but measure is not specified to detect disparities, provide follow-up plans:</p>	<p>2h C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/></p>
<p>TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for <i>Scientific Acceptability of Measure Properties</i>?</p>	<p>2</p>
<p>Steering Committee: Overall, to what extent was the criterion, <i>Scientific Acceptability of Measure Properties</i>, met? Rationale:</p>	<p>2 C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>
<p>3. USABILITY</p>	
<p>Extent to which intended audiences (e.g., consumers, purchasers, providers, policy makers) can understand the results of the measure and are likely to find them useful for decision making. (evaluation criteria)</p>	<p>Eval Rating</p>
<p>3a. Meaningful, Understandable, and Useful Information</p> <p>3a.1 Current Use: In use</p> <p>3a.2 Use in a public reporting initiative (disclosure of performance results to the public at large) (<i>If used in a public reporting initiative, provide name of initiative(s), locations, Web page URL(s). If not publicly</i></p>	<p>3a C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>

<p><u>reported, state the plans to achieve public reporting within 3 years):</u> Currently being considered for NQF endorsement, the STS CABG Composite Score is a multidimensional performance measure comprised of four domains consisting of 11 individual NQF-endorsed cardiac surgery metrics: (1) Operative Care--use of the internal mammary artery; (2) Perioperative Medical Care (use of preoperative beta blockade; discharge beta blockade, antiplatelet agents, and lipid-lowering agents--an "all-or-none" measure); (3) Risk-adjusted Operative Mortality; and (4) Risk-Adjusted Postoperative Morbidity (occurrence of postoperative stroke, renal failure, prolonged ventilation, re-exploration, or deep sternal wound infection--an "any-or-none" measure). Composite star ratings are presented on the STS website, www.sts.org/publicreporting and in the health section of the Consumers Union website, www.ConsumerReportsHealth.org. There are approximately 330 STS Adult Cardiac Surgery Database Participants who voluntarily participate in the Consumer's Union public reporting initiative. In addition, approximately 352 STS Adult Cardiac Surgery Database Participants voluntarily take part in STS Public Reporting Online.</p> <p>STS plans to publicly report more measures in the future. There is no definite date yet assigned to this measure; however, STS staff and surgeon leadership have engaged in initial internal STS discussions regarding this matter.</p> <p>3a.3 If used in other programs/initiatives (If used in quality improvement or other programs/initiatives, name of initiative(s), locations, Web page URL(s). <u>If not used for QI, state the plans to achieve use for QI within 3 years):</u> CMS Physician Quality Reporting Initiative (PQRI), www.cms.hhs.gov/pqri</p> <p>Testing of Interpretability (Testing that demonstrates the results are understood by the potential users for public reporting and quality improvement)</p> <p>3a.4 Data/sample (description of data/sample and size): See 3a.6 below</p> <p>3a.5 Methods (e.g., focus group, survey, QI project):</p> <p>3a.6 Results (qualitative and/or quantitative results and conclusions): Please see attached</p>	
<p>3b/3c. Relation to other NQF-endorsed measures</p> <p>3b.1 NQF # and Title of similar or related measures: ...</p>	
<p>(for NQF staff use) Notes on similar/related <u>endorsed</u> or submitted measures:</p>	
<p>3b. Harmonization If this measure is related to measure(s) already <u>endorsed by NQF</u> (e.g., same topic, but different target population/setting/data source <u>or</u> different topic but same target population): 3b.2 Are the measure specifications harmonized? If not, why? N/A; however, data definitions and key elements have been established by a multi-societal writing committee called the "ACCF/AHA Writing Committee to Develop Acute Coronary Syndromes and Coronary Artery Disease Clinical Data Standards" with representatives from each of the following organizations:</p> <p>Agency for Healthcare Research and Quality American College of Cardiology American College of Chest Physicians American College of Emergency Physicians American College of Physicians American College of Preventative Medicine American Heart Association American Medical Association Centers for Disease Control and Prevention Emergency Nurses Association Food and Drug Administration Joint Commission on Accreditation of Healthcare Organizations</p>	<p>3b C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/></p>

<p>National Association of Emergency Medical Technicians National Association of EMS Physicians National Heart, Lung, and Blood Institute Preventive Cardiovascular Nurses Association Society for Academic Emergency Medicine Society of Chest Pain Centers and Providers Society of General Internal Medicine Society of Thoracic Surgeons</p>	
<p>3c. Distinctive or Additive Value 3c.1 Describe the distinctive, improved, or additive value this measure provides to existing NQF-endorsed measures: n/a</p> <p>5.1 If this measure is similar to measure(s) already endorsed by NQF (i.e., on the same topic and the same target population), Describe why it is a more valid or efficient way to measure quality: n/a</p>	<p>3c C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/></p>
<p>TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for <i>Usability</i>?</p>	<p>3</p>
<p>Steering Committee: Overall, to what extent was the criterion, <i>Usability</i>, met? Rationale:</p>	<p>3 C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>
4. FEASIBILITY	
<p>Extent to which the required data are readily available, retrievable without undue burden, and can be implemented for performance measurement. (evaluation criteria)</p>	<p>Eval Rating</p>
<p>4a. Data Generated as a Byproduct of Care Processes</p> <p>4a.1-2 How are the data elements that are needed to compute measure scores generated? Data generated as byproduct of care processes during care delivery (Data are generated and used by healthcare personnel during the provision of care, e.g., blood pressure, lab value, medical condition), Coding/abstraction performed by someone other than person obtaining original information (E.g., DRG, ICD-9 codes on claims, chart abstraction for quality measure or registry)</p>	<p>4a C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>
<p>4b. Electronic Sources</p> <p>4b.1 Are all the data elements available electronically? (<i>elements that are needed to compute measure scores are in defined, computer-readable fields, e.g., electronic health record, electronic claims</i>) Yes</p> <p>4b.2 If not, specify the near-term path to achieve electronic capture by most providers.</p>	<p>4b C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>
<p>4c. Exclusions</p> <p>4c.1 Do the specified exclusions require additional data sources beyond what is required for the numerator and denominator specifications? No</p> <p>4c.2 If yes, provide justification.</p>	<p>4c C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/></p>
<p>4d. Susceptibility to Inaccuracies, Errors, or Unintended Consequences</p> <p>4d.1 Identify susceptibility to inaccuracies, errors, or unintended consequences of the measure and describe how these potential problems could be audited. If audited, provide results. This measure may be susceptible to human error (i.e., recording the measure inaccurately or not at all).</p>	<p>4d C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>

<p>When data collection on this measure is done through participation in the STS Adult Cardiac Surgery Database, an auditing strategy is in place.</p> <p>Both STS and the Duke Clinical Research Institute have a list of database participants making participation in the STS Adult Cardiac Surgery Database easy to track.</p> <p>Each participant is responsible for the quality and accuracy of the data they submit to the database. The participant agrees to the following quality control measures in the participation agreement:</p> <p>i) Participant hereby warrants that all data submitted for inclusion in the STS National Database will be accurate and complete, and acknowledges that such data may be subject to independent audit. Participant will use its best efforts to address any data or related deficiencies identified by the independent data warehouse service provider and agrees to cooperate with and assist STS and its designees in connection with the performance of any independent audit.</p> <p>ii) Participant warrants that it will take all reasonable steps to avoid the submission of duplicative data for inclusion in the STS National Database, including but not limited to apprising the Director of the STS National Database and the independent data warehouse service provider about any other Participation Agreements in which an individual cardiothoracic surgeon named above or on Schedule A attached hereto (as amended from time to time) is also named.</p> <p>STS audited for these potential problems during testing.</p>	
<p>4e. Data Collection Strategy/Implementation</p> <p>4e.1 Describe what you have learned/modified as a result of testing and/or operational use of the measure regarding data collection, availability of data/missing data, timing/frequency of data collection, patient confidentiality, time/cost of data collection, other feasibility/ implementation issues:</p> <p>4e.2 Costs to implement the measure (costs of data collection, fees associated with proprietary measures): Data Collection: There are no direct costs to collect the data for this measure. Costs to develop the measure included volunteer cardiothoracic surgeon time, STS staff time, and DCRI statistician and project management time.</p> <p>Other fees: STS Adult Cardiac Surgery Database participants (single cardiothoracic surgeons or a group of surgeons) pay annual participant fees of \$2,950 or \$3,700, depending on whether participants are STS members (or whether the majority of surgeons in a group are STS members). As a benefit of STS membership, STS members are charged the lesser of the two fees.</p> <p>4e.3 Evidence for costs:</p> <p>4e.4 Business case documentation:</p>	<p>4e C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>
<p>TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for <i>Feasibility</i>?</p>	<p>4</p>
<p>Steering Committee: Overall, to what extent was the criterion, <i>Feasibility</i>, met? Rationale:</p>	<p>4 C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/></p>
<p>RECOMMENDATION</p>	
<p>(for NQF staff use) Check if measure is untested and only eligible for time-limited endorsement.</p>	<p>Time-limited</p>

	<input type="checkbox"/>
Steering Committee: Do you recommend for endorsement? Comments:	Y <input type="checkbox"/> N <input type="checkbox"/> A <input type="checkbox"/>
CONTACT INFORMATION	
Co.1 Measure Steward (Intellectual Property Owner) Co.1 Organization Society of Thoracic Surgeons, 633 North Saint Clair Street, Suite 2320, Chicago, Illinois, 60611	
Co.2 Point of Contact Jane, Han, MSW, jhan@sts.org, 312-202-5856-	
Measure Developer If different from Measure Steward Co.3 Organization Society of Thoracic Surgeons, 633 North Saint Clair Street, Suite 2320, Chicago, Illinois, 60611	
Co.4 Point of Contact Jane, Han, MSW, jhan@sts.org, 312-202-5856-	
Co.5 Submitter If different from Measure Steward POC Jane, Han, MSW, jhan@sts.org, 312-202-5856-, Society of Thoracic Surgeons	
Co.6 Additional organizations that sponsored/participated in measure development	
ADDITIONAL INFORMATION	
Workgroup/Expert Panel involved in measure development Ad.1 Provide a list of sponsoring organizations and workgroup/panel members' names and organizations. Describe the members' role in measure development. Members of the STS Task Force on Quality Initiatives provide clinical expertise as needed. The STS Workforce on National Databases meets at the STS Annual Meeting and reviews the measures on a yearly basis. Changes or updates to the measure will be at the recommendation of the Workforce.	
Ad.2 If adapted, provide name of original measure: Ad.3-5 If adapted, provide original specifications URL or attachment	
Measure Developer/Steward Updates and Ongoing Maintenance Ad.6 Year the measure was first released: 2004 Ad.7 Month and Year of most recent revision: 12, 2010 Ad.8 What is your frequency for review/update of this measure? annually Ad.9 When is the next scheduled review/update for this measure? 2011	
Ad.10 Copyright statement/disclaimers:	
Ad.11 -13 Additional Information web page URL or attachment: Attachment 0125 Sections 1b.2, 1b.4, 2b.3, 2f.3, 3a.6.pdf	
Date of Submission (MM/DD/YY): 06/13/2011	

1b.2. Summary of Measure Results Demonstrating Performance Gap (*Descriptive statistics for performance results for this measure - distribution of scores for measured entities by quartile/decile, mean, median, SD, min, max, etc.*)

<i>Measurement</i>	<i>Timing of Antibiotic Administration for Cardiac Surgery Patients</i>
N	786
Mean	98.0%
1 st	83.2%
5 th	93.2%
10 th	95.2%
25 th	97.7%
Median	99.2%
75 th	99.9%
90 th	100.0%
95 th	100.0%
99 th	100.0%
Outlier	347 (44.1%)
High	259
Low	88

1b.4. Summary of Measure Results on Disparities by Population Group (*Descriptive statistics for performance results for this measure by population group*)

Timing of Antibiotic Administration for Cardiac Surgery Patients

<i>Measurement</i>	<i>Population Group</i>	
	<i>Men</i>	<i>Women</i>
N	887	819
Mean	97.6%	97.4%
1 st	70.0%	70.9%
5 th	91.7%	90.6%
10 th	94.7%	93.8%
25 th	97.5%	97.1%
Median	98.9%	98.9%
75 th	99.7%	100.0%
90 th	100.0%	100.0%
95 th	100.0%	100.0%
99 th	100.0%	100.0%
Outlier	462 (52.1%)	294 (35.9%)
High	366	211
Low	96	83

Timing of Antibiotic Administration for Cardiac Surgery Patients

<i>Measurement</i>	<i>Population Group</i>		
	<i>Black</i>	<i>White</i>	<i>Other</i>
N	231	881	192
Mean	97.2%	97.6%	96.3%
1 st	71.1%	69.2%	34.3%
5 th	87.0%	91.8%	86.9%
10 th	93.7%	94.2%	90.2%
25 th	97.3%	97.4%	96.7%
Median	98.7%	98.9%	98.5%
75 th	100.0%	99.7%	100.0%
90 th	100.0%	100.0%	100.0%
95 th	100.0%	100.0%	100.0%
99 th	100.0%	100.0%	100.0%

Timing of Antibiotic Administration for Cardiac Surgery Patients

Population Group

	<i>Black</i>	<i>White</i>	<i>Other</i>
<i>Measurement</i>			
Outlier	52 (22.5%)	532 (60.4%)	57 (29.7%)
High	29	424	35
Low	23	108	22

Timing of Antibiotic Administration for Cardiac Surgery Patients


Population Group

	<i>Hispanic</i>	<i>Non-Hispanic</i>
<i>Measurement</i>		
N	151	887
Mean	96.6%	97.6%
1 st	48.6%	69.4%
5 th	89.6%	91.5%
10 th	92.8%	94.2%
25 th	96.3%	97.4%
Median	98.6%	98.9%
75 th	100.0%	99.7%
90 th	100.0%	100.0%
95 th	100.0%	100.0%
99 th	100.0%	100.0%
Outlier	34 (22.5%)	546 (61.6%)
High	25	436
Low	9	110

2b.3. Testing Results (*Reliability statistics, assessment of adequacy in the context of norms for the test conducted*)

Testing results: $\rho = 0.60$

Timing of Antibiotic Administration for Cardiac Surgery Patients ($\rho=0.6$)



2f.3. Measure Scores from Testing or Current Use (*Description of scores, e.g., distribution by quartile, mean, median, SD, etc.; identification of statistically significant and meaningfully differences in performance*)

Results below are from January 1, 2009-December 31, 2009. The sample contains 786 STS Adult Cardiac Surgery Database Participants who had at least 100 eligible cases for the measure and reported data to STS for all 12 months.

<i>Measurement</i>	<i>Timing of Antibiotic Administration for Cardiac Surgery Patients</i>
N	786
Mean	98.0%
1 st	83.2%
5 th	93.2%
10 th	95.2%
25 th	97.7%
Median	99.2%
75 th	99.9%
90 th	100.0%
95 th	100.0%
99 th	100.0%
Outlier†	347 (44.1%)
High	259
Low	88

†Represents the number of participants that are outliers according to two-sided 95% binomial confidence interval.

3a.6. Results *(Qualitative or quantitative results and conclusions)*

Although formal testing of interpretability has not been performed, this measure has been used and reported for STS Adult Cardiac Surgery database participants since 2007. Current report presentation and interpretation manuals are presented below. These materials are updated as needed based upon feedback from database participants.

1) Report Overview and Interpretation Manual:

The NQF Measures Report

a. Organization

This report section is separated into three areas corresponding to: 1) NQF volume measures, 2) NQF process measures, and 3) NQF outcomes measures, in that order. The header at the top of each page references the report section for that page. Each NQF measure is presented on a single row in the section. Tabular data are on the left-hand side of each page and a standard graphic representation is shown on the right-hand side.

b. Statistical Calculation and Details – NQF Measures

Time period: This report section contains information on the individual STS participant and overall STS performance for the most recent 12 months for volume, process and CABG outcomes measures and the most recent 60 months for Valve and Valve + CABG outcomes. The 5 years (60 months) of performance for outcomes involving Valve procedures is necessary due to smaller sample sizes.

Volume Measures: The NQF report provides average annual case volumes data for three surgery categories: i) Isolated CABG, ii) Valve without CABG, and iii) combined CABG + Valve. Definitions of the three surgery categories are provided in Table 2 of this NQF Report Overview. For each type of surgery, the participant's annualized volume is calculated as:

$$\text{Participant Annualized Volume} = 12 \times (\# \text{ of surgeries}) / (\# \text{ of months})$$

where (# of surgeries) denotes the number of surgeries of the specified type performed by the participant during the specified time period, and (# of months) is the number of months during the specified time period for which the participant submitted at least one cardiac surgery of any type. The intent of calculating “annualized” volumes is to adjust for participants who participated in the database for fewer months than the time period specified. For participants who participated in the database and submitted cases every month during 2006, the annualized volume for 2006 is simply the total number of cases.

The STS Average Annualized Volume is the average value of all of the participant annualized volumes across the entire population of STS participants. The Participant Percentile indicates the percent of STS participants whose annualized volumes are less than, or equal to, your own. Higher percentiles indicate higher volumes in relation to other STS participant sites. The Distribution of Participant Values shows the range and percentiles of the distribution of participant annualized volumes across all database participants. For example, 90% of participants have annualized volumes less than or equal to the value marked “90th percentile.” Confidence intervals are not provided for volume measures, as volume is known with certainty and is not estimated.

Process Measures: The NQF process measures provide data on the frequency of usage of five therapies among subsets of Isolated CABG patients. The therapies are: i) preoperative beta blockade therapy, ii) use of IMA, iii) discharge anti-platelet medication, iv) discharge beta blockade therapy, and v) discharge anti-lipid medication. The patient population for each measure differs, in accordance with the NQF specifications (see Table 2 of this NQF Report Overview for details). The number of Eligible

Procedures is the number of cases performed by the participant during the specified time period who meet the eligibility requirements to be included in the calculations when summarizing the participant's data. ***Beginning with the 2008 Harvest 3 report (covering the procedure time period through 6/30/2008), STS implementation of NQF medication process measures using data version 2.61 excludes records for which the medication was contraindicated/not indicated from the eligible population.*** The main summary statistic, Participant Usage, is the percent of eligible Isolated CABG cases during the specified time period for which the patient received the specified therapy. The Overall STS Usage is the percent of all eligible patients in the entire STS population during the specified time period who received the specified therapy. ***In calculating these percentages, missing data are treated as a "No", emphasizing the importance of having complete data in these fields.***

The Participant Percentile indicates the percent of STS participants who applied the therapy in their respective populations less frequently than or as frequently as did your institution. The Distribution of Participant Values shows the range and percentiles of the distribution of participant usage across all participants in the database. For example, 90% of participants use the therapy less frequently than the amount indicated by the "90th percentile". A bar identified as "Participant" indicates the point estimate and limits of a 95% Confidence Interval (CI) for the participant's usage of therapy. The underlying parameter being estimated is the long-run usage rate that would be observed in a large sample of patients. The 95% CI indicates the range of usage rates that are consistent with the data in light of sampling variability.

Outcomes Measures: The NQF outcomes data provide risk-adjusted analyses of mortality and morbidity for Isolated CABG surgery as well as risk-adjusted operative mortality for Isolated AVR, Isolated MVR, AVR+CABG, and MVR+CABG. The main summary statistic provided is the Participant's Estimated Odds Ratio (OR) based on a hierarchical logistic regression analysis. The OR measures the impact that a participant's performance level has on a patient's probability of experiencing an adverse outcome. The interpretation is similar to that of an O/E ratio (see the Risk-Adjusted Results: Overview portion of the General Report Overview for details on STS risk adjustment). An OR greater than 1.0 implies that the participant increases a patient's risk of experiencing the outcome, relative to an "average" STS participant. An OR less than 1.0 implies that the participant decreases a patient's risk of experiencing the outcome, relative to an "average" STS participant. Each measure is calculated among patients undergoing surgery of the type specified during the time period specified who additionally meet certain eligibility requirements. The column labeled Eligible Procedures indicates the number of patients who met the inclusion criteria to be included in the analysis for the indicated measure. The Participant Percentile is the percent of STS participants who have an estimated OR that is greater than or equal to your estimated OR. Note that this is different than performance percentiles for process measures, where the percentile indicates the percentage of STS participants with performance that is *less than* the specified number. This simply reflects the fact that high process compliance is desirable, whereas a high OR is undesirable.

The Observed Participant Rate is the percent of eligible patients who experienced the specified outcome. Unlike the participant estimated OR, the observed participant rate is not risk-adjusted. The estimated OR is the main summary statistic for summarizing the NQF measure in this report.

The Distribution of Participant Values shows the range and percentiles of the distribution of estimated Odds Ratios across all STS participants. For example, 90% of STS participants have an OR greater than the value indicated by the "90th percentile." The line that extends to the left and right of the Participant Value indicates the lower and upper limits of a 95% Confidence Interval (CI) surrounding the participant's estimated OR.

c. Technical Notes

Calculation of Percentiles for the Distribution of Participant Values: The graph provided for each measure contains information about the distribution of the value of the measure across all STS

participants, namely the minimum, maximum, 10th percentile, 50th percentile, and 90th percentile. The “Xth” percentile, denoted P_x , is loosely defined as the number having the property that X% of the participant values are less than P_x , and (100 – X)% of the participant values are greater than P_x . **For process measures, participants with greater than 5% missing data were excluded when calculating percentiles of the STS distribution and do not have a calculated participant percentile.** For participants having less than 5% missing data on a process measure, the missing values on the process measure were converted to “No” before calculating percentiles. For outcomes measures, all participants submitting at least one eligible case were included when calculating percentiles of the STS distribution. Missing data on outcomes variables were treated as “No.”


NQF/STS Results Comparison: Participants may see some differences between summaries of their data provided in the NQF section of the report and summaries of their data reported elsewhere in the STS report. These differences are due to subtle variations in variable definitions, patient inclusion and exclusion criteria, and rules for handling missing data in the NQF section versus the rest of the report. Definitions used in the NQF report were designed to match current NQF specifications as closely as possible. It is expected that these differences will eventually disappear as the NQF measures are refined. Some important differences are:

Case Volumes – The NQF report section presents “annualized” volumes. These are case volumes that have been adjusted for the number of months that a participant was an active contributor to the database. Elsewhere in the STS report, total case volumes are presented without adjustment for the length of participation.

Eligible Cases - The NQF report also presents the number of “eligible cases” for each measure. Separate inclusion criteria are applied to each measure, and these inclusion criteria do not always match the definitions used elsewhere in the STS report. Please refer to the footnotes in each section for specific details.

Interpretation Manual

In addition to the statistics provided for each of the STS Composite Quality Domains and NQF measures, a figure representing the distribution of values for the entire STS population is provided.



The figure allows participants to quickly judge their performance relative to the overall STS. The scale of the figure is set up such that the right side of the distribution represents the most favorable performance and the left side represents the least favorable performance (Note that in some cases smaller numbers will be on the left; in other instances, smaller numbers will be on the right. For example, for the Pre-operative Beta Blockade Therapy measure, the far left side of the distribution will contain the *lowest* percentage Beta Blockade Therapy for an STS participant – this corresponds to least

favorable performance. Alternatively, for the Operative Mortality Measure, the far left side of the distribution will contain the *highest* Estimated Odds Ratio – this also corresponds to least favorable performance). If a participant’s value for a given measure is to the left of the STS overall value, the participant is performing worse on that measure than the overall STS. Conversely, if the participant’s value for a given measure is located to the right of the overall STS value, the participant is performing better than the overall STS.

NOTE! Care should be given to reading these figures. In some instances, the various percentiles presented cluster very close together in the data. In such cases, the label for the percentile is not necessarily located immediately at the point on the distribution where the percentile occurs. An example of this is apparent in the figure above: The 50th percentile corresponds to a value of 93.7 and looks to align fairly closely with the STS overall value as represented by the large black dot. However, the expandable figure marking actually points to a place somewhere to the right of the STS overall value for the 50th percentile marking. So the STS overall value would be some amount less than 93.7.

Also, please note that in some cases, small sample sizes preclude valid comparisons between the participant and the STS overall. Such instances are clearly noted in the report output.

a. NQF Measures Interpretation Example

Sample CABG Operative Mortality results – tabular and figure representation.


NQF Measure	Eligible Procedures	Participant Estimated OR	Participant Percentile	Participant Observed Rate
2005 CABG Operative Mortality	74	1.14	26.3	5.4%

Eligible Procedures: 74 patients met the inclusion criteria for the indicated measure.

Participant Estimated OR (Odds Ratio): The main summary statistic measuring the impact that a participant’s performance has on a patient’s probability of experiencing an adverse outcome has a value of 1.14 indicating worse than expected performance.

Participant Percentile: 26.3% of STS participants had an estimated OR greater than or equal to your estimated OR. In other words, 26.3% had the same or worse performance.

Participant Observed Rate: 5.4% of the 74 eligible patients experienced the specified outcome.



The highest OR among all STS participants = 2.29
 The lowest OR among all STS participants = 0.45
 The STS average OR is 1.00

The 95% confidence interval for the participant's OR spans from <0.45 to ~1.90

2) Sample page from section of the report that contains NQF measure results:



**NQF Measures
Process Measures
Participant 99999
STS Period Ending 12/31/2008**



NQF Measure	Eligible Procedures	Participant Usage (95% CI)	Participant Percentile	Overall STS Usage	Distribution of Participant Values ● = Overall STS Usage
Jan 2008 - Dec 2008 Preoperative Beta Blockade Therapy ¹	541	89.3% (86.4 , 91.8)	69.9	82.1%	
Jan 2008 - Dec 2008 Use of IMA ²	536	96.5% (94.5 , 97.9)	63.3	94.2%	
Jan 2008 - Dec 2008 Discharge Anti-Platelet Medication ³	536	98.7% (97.3 , 99.5)	68.7	96.1%	
Jan 2008 - Dec 2008 Discharge Beta Blockade Therapy ⁴	538	96.1% (94.1 , 97.6)	53.4	93.7%	
Jan 2008 - Dec 2008 Discharge Anti-Lipid Treatment ⁴	535	91.8% (89.1 , 94.0)	40.7	91.4%	

¹Excludes v2.61 contraindicated / not indicated records.

²Excludes patients with prior CABG surgery

³Anti-platelet use includes Aspirin and ADP Inhibitors, and excludes in-hospital mortalities. Excludes v2.61 contraindicated / not indicated records.

⁴Excludes in-hospital mortalities. Excludes v2.61 contraindicated / not indicated records.