

NATIONAL QUALITY FORUM

Measure Evaluation 4.1 December 2009

This form contains the measure information submitted by stewards. Blank fields indicate no information was provided. Attachments also may have been submitted and are provided to reviewers. The subcriteria and most of the footnotes from the [evaluation criteria](#) are provided in Word comments within the form and will appear if your cursor is over the highlighted area. Hyperlinks to the evaluation criteria and ratings are provided in each section.

TAP/Workgroup (if utilized): Complete all **yellow highlighted** areas of the form. Evaluate the extent to which each subcriterion is met. Based on your evaluation, summarize the strengths and weaknesses in each section.

Note: If there is no TAP or workgroup, the SC also evaluates the subcriteria (**yellow highlighted areas**).

Steering Committee: Complete all **pink** highlighted areas of the form. Review the workgroup/TAP assessment of the subcriteria, noting any areas of disagreement; then evaluate the extent to which each major criterion is met; and finally, indicate your recommendation for the endorsement. Provide the rationale for your ratings.

Evaluation ratings of the extent to which the criteria are met

C = Completely (unquestionably demonstrated to meet the criterion)

P = Partially (demonstrated to partially meet the criterion)

M = Minimally (addressed BUT demonstrated to only minimally meet the criterion)

N = Not at all (NOT addressed; OR incorrectly addressed; OR demonstrated to NOT meet the criterion)

NA = Not applicable (only an option for a few subcriteria as indicated)

(for NQF staff use) NQF Review #: 1532	NQF Project: Surgery Endorsement Maintenance 2010
MEASURE DESCRIPTIVE INFORMATION	
De.1 Measure Title: Plasma Transfusion Indication	
De.2 Brief description of measure: Percentage of transfused plasma units (bags) with pre-transfusion PT/INR result and clinical indication documented - applicable to inpatients of all ages	
1.1-2 Type of Measure: Process	
De.3 If included in a composite or paired with another measure, please identify composite or paired measure PBM-03 is a part of the Patient Blood Management (PBM) measure set: PBM-01 (Transfusion Consent), PBM-02 (RBC Transfusion Indication), PBM-04 (Platelet Transfusion Indication), PBM-05 (Blood Administration Documentation), PBM-06 (Preoperative Anemia Screening), PBM-07 (Preoperative Blood Type Testing and Antibody Screening).	
De.4 National Priority Partners Priority Area: Care coordination, Safety, Overuse	
De.5 IOM Quality Domain: Effectiveness, Patient-centered, Safety	
De.6 Consumer Care Need: Getting better, Living with illness	

CONDITIONS FOR CONSIDERATION BY NQF	
Four conditions must be met before proposed measures may be considered and evaluated for suitability as voluntary consensus standards:	NQF Staff
A. The measure is in the public domain or an intellectual property (measure steward agreement) is signed. <i>Public domain only applies to governmental organizations. All non-government organizations must sign a measure steward agreement even if measures are made publicly and freely available.</i> A.1 Do you attest that the measure steward holds intellectual property rights to the measure and the right to use aspects of the measure owned by another entity (e.g., risk model, code set)? Yes A.2 Indicate if Proprietary Measure (as defined in measure steward agreement): A.3 Measure Steward Agreement: Agreement will be signed and submitted prior to or at the time of measure submission A.4 Measure Steward Agreement attached:	A Y <input type="checkbox"/> N <input type="checkbox"/>

B. The measure owner/steward verifies there is an identified responsible entity and process to maintain and update the measure on a schedule that is commensurate with the rate of clinical innovation, but at least every 3 years. Yes, information provided in contact section	B Y <input type="checkbox"/> N <input type="checkbox"/>
C. The intended use of the measure includes <u>both</u> public reporting <u>and</u> quality improvement. ► Purpose: Public reporting, Internal quality improvement Accountability	C Y <input type="checkbox"/> N <input type="checkbox"/>
D. The requested measure submission information is complete. Generally, measures should be fully developed and tested so that all the evaluation criteria have been addressed and information needed to evaluate the measure is provided. Measures that have not been tested are only potentially eligible for a time-limited endorsement and in that case, measure owners must verify that testing will be completed within 12 months of endorsement. D.1 Testing: Yes, fully developed and tested D.2 Have NQF-endorsed measures been reviewed to identify if there are similar or related measures? Yes	D Y <input type="checkbox"/> N <input type="checkbox"/>
(for NQF staff use) Have all conditions for consideration been met? Staff Notes to Steward (if submission returned):	Met Y <input type="checkbox"/> N <input type="checkbox"/>
Staff Notes to Reviewers (issues or questions regarding any criteria):	
Staff Reviewer Name(s):	

TAP/Workgroup Reviewer Name:	
Steering Committee Reviewer Name:	
1. IMPORTANCE TO MEASURE AND REPORT	
Extent to which the specific measure focus is important to making significant gains in health care quality (safety, timeliness, effectiveness, efficiency, equity, patient-centeredness) and improving health outcomes for a specific high impact aspect of healthcare where there is variation in or overall poor performance. Measures must be judged to be important to measure and report in order to be evaluated against the remaining criteria. (evaluation criteria) 1a. High Impact	Eval Rating
(for NQF staff use) Specific NPP goal :	
1a.1 Demonstrated High Impact Aspect of Healthcare: Frequently performed procedure, Leading cause of morbidity/mortality, Severity of illness, Patient/societal consequences of poor quality 1a.2 1a.3 Summary of Evidence of High Impact: The use of plasma has increased in the US and is disproportionally high compared to other countries with similar levels of health care. Indications for transfusing plasma are very limited, and as a result, published studies often show unjustifiable use of plasma. According to the National Heart Lung and Blood Institute, plasma should be administered only to increase the level of clotting factors in patients with a demonstrated deficiency. If the prothrombin time (PT) and partial thromboplastin time (PTT) are < 1.5 times normal, a plasma transfusion is rarely needed. 1a.4 Citations for Evidence of High Impact: Wilson K, Mac Dougall L, Fergusson D, et al. The effectiveness of interventions to reduce physician's levels of inappropriate transfusion: What can be learned from a systematic review of the literature? Transfusion 2002;42:1224-1229. Stansworth SJ, Brunskill SJ, Hyde CJ, et al. Is fresh frozen clinically effective? A systematic review of randomized controlled trials. Br J Haematology 2004;126:139-52. Roback JD, Caldwell S, Carson, et al. Evidence-based practice guidelines for plasma transfusions. Transfusion 2010;50:1227-39. DeAnda A Jr, Baker KM, Roseff SD, et al. Developing a blood conservation program in cardiac surgery. AM J Med Qual. 2006;21(4):230-237.	1a C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>

Helm RE, Rosengart TK, Gomez M. et al. Comprehensive multimodality blood conservation: 100 consecutive CABG operations without transfusion. *Ann Thorac Surg.* 1998;65(1):125-136.

Rosengart TK, Helm RE, DeBois WJ, Garcia N, Krieger KH, et al. Open heart operations without transfusion using a multimodality blood conservation strategy in 50 Jehovah's Witness patients: implications for a "bloodless" surgical technique. *J Am Coll Surg.* 1997;184(6):618-629.

Guerrero EB, Zhao Y, Obrien SM, Ferguson TB, Peterson ED, et al. Variation in use of blood transfusion in coronary artery bypass graft surgery. *JAMA* 2010;304(14) 1568-1575.

1b. Opportunity for Improvement

1b.1 Benefits (improvements in quality) envisioned by use of this measure: Studies show that plasma use varies across a large number of hospitals with no significant difference in mortality rates. Experts believe that the absence of differences in mortality strongly suggests inappropriate transfusions. Plasma is frequently transfused to patients with mild-to moderate elevations in PT despite numerous studies that have not shown a correlation between the risk of bleeding and mild-to moderate test results. Measuring and monitoring patients that receive platelets will provide data that can be used to determine if patients are receiving the best care based on the guidelines.

1b.2 Summary of data demonstrating performance gap (variation or overall poor performance) across providers:

In a study by Wahab et al, transfusion of plasma for mild abnormalities of coagulation values resulted in a partial normalization in a minority of patients, and failed to correct the PT in 99% of the patients. In a 2004 study by Hui, the need to correct prolonged international normalized ratios (INRs) for patients on warfarin emerged as the primary indication for plasma allowed by massive transfusions. Since blood transfusions may cause more harm than benefit, hospitals need to begin to monitor and evaluate plasma transfusions using a patient-centered approach that carefully evaluates patients for the need for each unit.

In the recent transfusion Requirements After Cardiac Surgery (TRAC) randomized controlled trial, 408 hospitals plasma use for patients undergoing isolated primary coronary artery bypass graft (CABG) surgery ranged from 0% to 97.5%.

Numerous studies have concluded that the routine administration of small quantities of plasma to perioperative patients with minor coagulopathies is probably of little hemostatic benefit and exposes the patient to numerous adverse reactions including volume overload and transfusion-related acute lung injury. Two meta-analyses were conducted where patients undergoing a variety of minor procedures were analyzed as to whether the perioperative PT/INR predicts the risk of major bleeding during those procedures. It was concluded that the preprocedure international normalized ration (INR) does not likely predict the bleeding risk.

Another meta-analysis determined that plasma administered to perioperative patients does not have a beneficial effect in reducing transfusion requirements or surgical blood loss.

One hospital collected data on fresh frozen plasma (FFP) and red blood cell (RBC) transfusions and measures of hospital activity and mortality over a 12-year period. Plasma orders were discouraged if the INR was less than 2.0, in the absence of bleeding. The use of vitamin K was encouraged if the patient was receiving warfarin. The program resulted in about an 80% reduction in transfused FFP. No unexpected bleeding was reported and inpatient mortality decreased during the 12 years while overall hospital activity remained largely unchanged.

1b.3 Citations for data on performance gap:

Tavares, M, DiQuattro P, Nolette N, Conti G, Sweeney J. Reduction in plasma after enforcement of transfusion guidelines. Article first published on line 4 Oct 2010.

Stansworth SJ, Brunskill SJ, Hyde CJ, et al. Is fresh frozen clinically effective? A systematic review of randomized controlled trials. *Br J Haematology* 2004;126:139-52.

Segal JB, Dzik WH. Paucity of studies to support t that abnormal coagulation test results predict bleeding in the setting of invasive procedures: An evidence-based review. *Transfusion* 2005;45:1413-25.

Guerrero EB, Zhao Y, Obrien SM, Ferguson TB, Peterson ED, et al. Variation in use of blood transfusion in coronary artery bypass graft surgery. *JAMA* 2010;304(14) 1568-1575.

Hui C, Williams I, Davis K. Clinical audit of the use of fresh-frozen plasma and platelets in a tertiary teaching hospital and the impact of a new transfusion request form. *Int Med J.* 2005;35:283-288.

Wallis JP, Dzik S. Is fresh frozen plasma over transfused in the United States? *Transfusion.* 2004;44:1674-75.

Ardel-Wahab OI, Healy B, Dzik WH. Effect of fresh-frozen plasma transfusion on prothrombin time and bleeding in patients with mild coagulation abnormalities. *Transfusion.* 2006;46:1479-1285.

1b

C ☐P ☐M ☐N ☐

Segal J, Dzik WH; Transfusion Medicine/Hemostasis Clinical Trials Network. Paucity of studies to support that abnormal coagulation test results predict bleeding in the setting of invasive procedures: an evidenced-based review. *Transfusion*. 2005;45:1413-25.

Tavares M, Diquattro P, Nolette N et al. Reduction in plasma transfusion after enforcement of transfusion guidelines. Retrieved from the world wide web at <http://www.ncbi.nlm.nih.gov/pubmed/20946197>

1b.4 Summary of Data on disparities by population group:

None

1b.5 Citations for data on Disparities:

NA

1c. Outcome or Evidence to Support Measure Focus

1c.1 Relationship to Outcomes (*For non-outcome measures, briefly describe the relationship to desired outcome. For outcomes, describe why it is relevant to the target population*): Plasma transfusions are often ordered for diverse indications and in many cases, the adverse effects of plasma may outweigh any potential benefit. Plasma transfusions can increase the risk of acute lung injury and has been shown to increase mortality. Once data is collected at the patient level, adverse events can be tracked and reported to the National Hemovigilance Database that will use the data to identify ways to improve patient outcomes.

1c.2-3. Type of Evidence: Observational study, Evidence-based guideline, Randomized controlled trial, Expert opinion, Systematic synthesis of research, Meta-analysis

1c.4 Summary of Evidence (*as described in the criteria; for outcomes, summarize any evidence that healthcare services/care processes influence the outcome*):

Studies have demonstrated that use of a blood conservation program can significantly decrease transfusion rates over time. Plasma transfusions are commonly prescribed for a variety of indications, but the scientific evidence supporting many plasma transfusion practices is limited and in many cases, the adverse effects of plasma may outweigh any potential benefit. Even when transfusion criteria are met, the clinical efficacy of prophylactic plasma is questionable. A systematic review of 57 randomized controlled trials involving the use of plasma for a variety of indications found insufficient evidence to support or refute any value in treating with plasma.

The AABB convened a panel who reviewed the data for plasma and only two recommendations could be made. A systematic review and meta-analysis of the plasma transfusion literature was performed. Then, six questions were developed and a methodology called GRADE (Grading for Recommendations, Assessment, Development and Evaluation) was used. The panel suggested that plasma be transfused to patients requiring massive transfusion (quality of evidence = moderate) and that plasma be transfused in patients with warfarin anticoagulation-related intracranial hemorrhage (quality of evidence = low).

A retrospective study was done in the intensive care unit (ICU) that showed that critically ill patients frequently receive inappropriate FFP transfusions. However after review, many transfusions may be appropriate for the ICU setting even though they were inconsistent with the expert recommendations. Until more randomized clinical trials can be done, they suggested that education, audits or request forms and feedback about adverse events and costs of FFP transfusions be compiled.

1c.5 Rating of strength/quality of evidence (*also provide narrative description of the rating and by whom*):

NA

1c.6 Method for rating evidence: NA

1c.7 Summary of Controversy/Contradictory Evidence: None

1c.8 Citations for Evidence (*other than guidelines*): Wilson K, Mac Dougall L, Fergusson D, et al. The effectiveness of interventions to reduce physician's levels of inappropriate transfusion: What can be learned from a systematic review of the literature? *Transfusion* 2002;42:1224-1229.

Stansworth SJ, Brunskill SJ, Hyde CJ, et al. Is fresh frozen clinically effective? A systematic review of randomized controlled trials. *Br J Haematol* 2004;126:139-52.

1c

C ☐

P ☐

M ☐

N ☐

Roback JD, Caldwell S, Carson, et al. Evidence-based practice guidelines for plasma transfusions. *Transfusion* 2010;50:1227-39.

DeAnda A Jr, Baker KM, Roseff SD, et al. Developing a blood conservation program in cardiac surgery. *AM J Med Qual.* 2006;21(4):230-237.

Helm RE, Rosengart TK, Gomez M. et al. Comprehensive multimodality blood conservation: 100 consecutive CABG operations without transfusion. *Ann Thorac Surg.* 1998;65(1):125-136.

Rosengart TK, Helm RE, DeBois WJ, Garcia N, Krieger KH, et al. Open heart operations without transfusion using a multimodality blood conservation strategy in 50 Jehovah's Witness patients: implications for a "bloodless" surgical technique. *J Am Coll Surg.* 1997;184(6):618-629.

Guerrero EB, Zhao Y, O'Brien SM, Ferguson TB, Peterson ED, et al. Variation in use of blood transfusion in coronary artery bypass graft surgery. *JAMA* 2010;304(14) 1568-1575.

Hajjar LA, Vincent JL, Galas FRBG, Nakamura RE, Silva CMP, et al. Transfusion requirements after cardiac surgery: the TRACS randomized controlled trial. *JAMA* 2010; 304(14)1559-1567.

Shander AS, Goodnough LT. Blood transfusion as a quality indicator in cardiac surgery. *JAMA* 2010;(14)1610-1611.

1c.9 Quote the Specific guideline recommendation (including guideline number and/or page number):

It is reasonable to transfuse non-red cell hemostatic blood products based on clinical evidence of bleeding and preferably guided by point-of-care tests that assess hemostatic function in a timely and accurate manner. (#4-2, p. S36).

1c.10 Clinical Practice Guideline Citation: Perioperative Blood Transfusion and Blood Conservation in Cardiac Surgery: The Society of Thoracic Surgeons and The Society of Cardiovascular Anesthesiologists Clinical Practice Guideline. *Ann. Thorac. Surg.*, May 2007; 83: S27 - S86.

1c.11 National Guideline Clearinghouse or other URL:

<http://www.sts.org/sections/aboutthesociety/practiceguidelines>

1c.12 Rating of strength of recommendation (also provide narrative description of the rating and by whom):

Level of evidence is C, Class IIa

1c.13 Method for rating strength of recommendation (If different from [USPSTF system](#), also describe rating and how it relates to USPSTF):

The classification system is the same as that used by the Joint Task Force for Guidelines of the American College of Cardiology (ACC) and the American Heart Association (AHA) available

at: http://circ.ahajournals.org/manual/manual_IIstep6.shtml

Classification of Recommendations

Class I: Conditions for which there is evidence and/or general agreement that a given procedure or treatment is useful and effective.

Class II: Conditions for which there is conflicting evidence and/or a divergence of opinion about the usefulness/efficacy of a procedure or treatment.

IIa. Weight of evidence/opinion is in favor of usefulness/efficacy

IIb. Usefulness/efficacy is less well established by evidence/opinion.

Class III: Conditions for which there is evidence and/or general agreement that the procedure/treatment is not useful/effective, and in some cases may be harmful.

Level of Evidence

Level of Evidence A: Data derived from multiple randomized clinical trials

Level of Evidence B: Data derived from a single randomized trial, or non-randomized studies

Level of Evidence C: Consensus opinion of experts

1c.14 Rationale for using this guideline over others:

This measure set includes elective cardiac surgery patients. This guideline is cited because it supports plasma usage based on clinical evidence and prefers that point-of-care testing is used to assess hemostatic function prior to transfusion.

TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for *Importance to Measure and Report*?

1

Steering Committee: Was the threshold criterion, <i>Importance to Measure and Report</i>, met? Rationale:	1 Y <input type="checkbox"/> N <input type="checkbox"/>	
2. SCIENTIFIC ACCEPTABILITY OF MEASURE PROPERTIES		
Extent to which the measure, as specified, produces consistent (reliable) and credible (valid) results about the quality of care when implemented. (evaluation criteria)	Eval Rating	
2a. MEASURE SPECIFICATIONS		
S.1 Do you have a web page where current detailed measure specifications can be obtained? S.2 If yes, provide web page URL: 2a. Precisely Specified		
2a.1 Numerator Statement (<i>Brief, text description of the numerator - what is being measured about the target population, e.g. target condition, event, or outcome</i>): Number of plasma doses(bags) with pre-transfusion PT/INR result and clinical indication documented 2a.2 Numerator Time Window (<i>The time period in which cases are eligible for inclusion in the numerator</i>): Episode of care 2a.3 Numerator Details (<i>All information required to collect/calculate the numerator, including all codes, logic, and definitions</i>): The units in the numerator are a subset of the denominator units. The following data elements are collected for the numerator: Clinical Indication for Plasma, Plasma ID, and Pre-transfusion PT/INR Result. Detailed descriptions are provided in attachment for Section 2a.30. 2a.4 Denominator Statement (<i>Brief, text description of the denominator - target population being measured</i>): Number of transfused plasma units evaluated 2a.5 Target population gender: Female, Male 2a.6 Target population age range: All age patients who received plasma 2a.7 Denominator Time Window (<i>The time period in which cases are eligible for inclusion in the denominator</i>): Episode of care 2a.8 Denominator Details (<i>All information required to collect/calculate the denominator - the target population being measured - including all codes, logic, and definitions</i>): Transfused units are identified using ICD-9-CM Procedure Codes or Blood Bank Records. The following data elements are collected for the denominator: Admission Date, Blood Administration Location, Discharge Date, and ICD-9-CM Principal or Other Procedure Codes or Blood Bank Records. Detailed descriptions are provided in attachment for Section 2a.30. 2a.9 Denominator Exclusions (<i>Brief text description of exclusions from the target population</i>): Trauma patients 2a.10 Denominator Exclusion Details (<i>All information required to collect exclusions to the denominator, including all codes, logic, and definitions</i>): Patients are excluded using ICD-9-CM Principal or Other Diagnosis Trauma Codes in Appendix A, Table 9.7. 2a.11 Stratification Details/Variables (<i>All information required to stratify the measure including the stratification variables, all codes, logic, and definitions</i>): Units may be stratified according to the blood administration location at the start of the transfusion. The definition is the location where the blood transfusion started. Allowable values for settings are: Intraoperative or Non-intraoperative Settings. 2a.12-13 Risk Adjustment Type: No risk adjustment necessary		
		2a-specs C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>

2a.14 Risk Adjustment Methodology/Variables (List risk adjustment variables and describe conceptual models, statistical models, or other aspects of model or method):

2a.15-17 Detailed risk model available Web page URL or attachment:

2a.18-19 Type of Score: Rate/proportion

2a.20 Interpretation of Score: Better quality = Higher score

2a.21 Calculation Algorithm (Describe the calculation of the measure as a flowchart or series of steps): Algorithms are provided in attachment for Section 2a.30.

2a.22 Describe the method for discriminating performance (e.g., significance testing):

During the six-month pilot, the distribution of the hospital rates was reviewed over time.

2a.23 Sampling (Survey) Methodology If measure is based on a sample (or survey), provide instructions for obtaining the sample, conducting the survey and guidance on minimum sample size (response rate):

For pilot testing, hospitals were requested to submit 10 cases of patients that were discharged from the designated six months with plasma transfusions. Post pilot, the sample size will be based on the number of plasma units transfused per discharge month or quarter.

Hospitals that choose to sample have the option of sampling quarterly or monthly. A hospital may choose to use a larger sample size than required. Hospitals with an initial population size less than the minimum number of units/bags transfused per quarter/month for the measure, cannot apply sampling to the measure.

2a.24 Data Source (Check the source(s) for which the measure is specified and tested)

Paper medical record/flow-sheet, Electronic administrative data/claims, Lab data

2a.25 Data source/data collection instrument (Identify the specific data source/data collection instrument, e.g. name of database, clinical registry, collection instrument, etc.):

The Joint Commission developed a web-based data collection tool that was used by hospitals and for reliability testing during the pilot test. When the measures are made part of The Joint Commission's ORYX data collection and reporting program, the data would be collected using contracted Performance Measurement Systems (vendors) that develop data collection tools based on the measure specifications. The tools are verified and tested by Joint Commission staff to confirm the accuracy of the data collection tool with the specifications.

2a.26-28 Data source/data collection instrument reference web page URL or attachment: Attachment The_Patient Blood_Management_Tool [1]-634279215776770950.pdf

2a.29-31 Data dictionary/code table web page URL or attachment: Attachment PBMSpecifications-634279442348907962.pdf

2a.32-35 Level of Measurement/Analysis (Check the level(s) for which the measure is specified and tested)

Facility/Agency, Can be measured at all levels

2a.36-37 Care Settings (Check the setting(s) for which the measure is specified and tested)

Hospital

2a.38-41 Clinical Services (Healthcare services being measured, check all that apply)

Clinicians: PA/NP/Advanced Practice Nurse, Clinicians: Physicians (MD/DO)

TESTING/ANALYSIS

2b. Reliability testing

2b.1 Data/sample (description of data/sample and size): A sample of 194 medical records were reabstracted at 12 randomly selected pilot hospitals July through September 2010.

2b.2 Analytic Method (type of reliability & rationale, method for testing):

2b
C ☐
P ☐
M ☐
N ☐

Hospitals for reliability testing were randomly selected based on multiple characteristics, including region (west, south, north central, northeast), hospital type (teaching/non-teaching, rural/urban), and bed size (0-99, 100-199, 200-299, 300+). The objectives of the reliability site visits included: evaluation of the reliability of the individual measures and associated data elements, assessment of data collection effort including abstraction time and estimated cost, assessment of measure specifications including definitions, abstraction guidelines, etc. and assessment of sampling strategies. To prepare for the reliability site visits, the data collection tool that was used by the pilot hospitals was enhanced and tested. During the reliability site visit, Joint Commission staff re-abstracted a sub-set of records that had been previously submitted by the hospital into the enhanced data collection tool without knowing the measure specific data values that the hospital had submitted. When reabstraction was completed for each record, the results from the hospital and Joint Commission staff were compared and differences adjudicated in the program. Focus group interviews were conducted at each hospital and findings were discussed with each hospital to understand what aspects could be improved. A comparison of calculated indicator rates using data originally abstracted by hospitals and the data that were reabstracted by The Joint Commission staff was adjudicated on each measure and the individual data elements. Statistical analysis utilized Kappa scores and p values.

2b.3 Testing Results (*reliability statistics, assessment of adequacy in the context of norms for the test conducted*):

The number of originally abstracted denominator cases was 49 events with a computed original measure rate of 78%. The number of re-abstracted denominator cases was 53 with a re-abstracted measure rate of 70%. The absolute difference was 7.7% with a Kappa score of 0.460. The percent of hospital identified population verified as 98%. The match rate for 55 cases for the individual data elements was: Clinical Indication for Plasma 45%, Plasma Event ID 100%, Plasma Event Total Doses 94%, and Pre-transfusion Laboratory Testing 67%. Measure specifications have been revised to strengthen and provide additional clarity to the data element definitions and abstraction guidelines.

2c. Validity testing

2c.1 Data/sample (*description of data/sample and size*): Face validity was tested by a total of 63 hospitals of various sizes and geographic locations across the country that represented over 300 individuals during August and May 2009. Measure specifications were sent to the test hospitals for review. In addition, on-site focus interviews were conducted at five hospitals. Criterion validity was evaluated during the reliability site visits mentioned above as well as through an online survey that the participating hospitals completed.

2c.2 Analytic Method (*type of validity & rationale, method for testing*):

The measure information form and the data dictionary were evaluated for face validity. The following parts of the measure information form were evaluated: numerator statement, numerator inclusions, numerator exclusions, denominator statement, denominator inclusions, denominator exclusions and an overall understanding of the measure information form. Each area was scored utilizing a five-point Likert scale. For each data element, the hospitals were asked to comment on the clarity and understanding of the abstraction guidelines and data definitions. In addition, the data dictionary was reviewed for overall understanding, usefulness and clarity utilizing a five-point Likert scale. Qualitative analysis was performed on measure feedback received during the focus group interviews and from the online surveys.

2c.3 Testing Results (*statistical results, assessment of adequacy in the context of norms for the test conducted*):

A total of 58 hospitals completed the face validity evaluation and rated the overall understanding of the numerator and denominator statements an average 4.3% that ranked the measure 4th out of the 10 measures. Modifications to improve the understanding and clarity of the measure specifications were made prior to pilot testing based on feedback received from the hospitals during the face validity evaluation. Analysis of the online survey revealed 98% (57/58) of the pilot hospitals recommended moving the measure forward to the pilot test with suggested modifications. Note: For alpha testing, a sample of all three blood products were the proposed population for one measure.

2c
C ☐
P ☐
M ☐
N ☐

2d. Exclusions Justified

2d.1 Summary of Evidence supporting exclusion(s):

2d
C ☐
P ☐
M ☐

<p>2d.2 Citations for Evidence:</p> <p>2d.3 Data/sample (<i>description of data/sample and size</i>):</p> <p>2d.4 Analytic Method (<i>type analysis & rationale</i>):</p> <p>2d.5 Testing Results (<i>e.g., frequency, variability, sensitivity analyses</i>):</p>	<p>N <input type="checkbox"/></p> <p>NA <input type="checkbox"/></p>
<p>2e. Risk Adjustment for Outcomes/ Resource Use Measures</p> <p>2e.1 Data/sample (<i>description of data/sample and size</i>):</p> <p>2e.2 Analytic Method (<i>type of risk adjustment, analysis, & rationale</i>):</p> <p>2e.3 Testing Results (<i>risk model performance metrics</i>):</p> <p>2e.4 If outcome or resource use measure is not risk adjusted, provide rationale:</p>	<p>2e</p> <p>C <input type="checkbox"/></p> <p>P <input type="checkbox"/></p> <p>M <input type="checkbox"/></p> <p>N <input type="checkbox"/></p> <p>NA <input type="checkbox"/></p>
<p>2f. Identification of Meaningful Differences in Performance</p> <p>2f.1 Data/sample from Testing or Current Use (<i>description of data/sample and size</i>): A sample of patients was selected from the eligible measure population. For each patient, a maximum of the first three 'events' (based on transfusion order) that could include up to three units or doses of blood from each of the three types of blood products were used for measurement purposes.</p> <p>2f.2 Methods to identify statistically significant and practically/meaningfully differences in performance (<i>type of analysis & rationale</i>): Z-scores were used to determine hospital measure rates that were significantly different from the overall average.</p> <p>2f.3 Provide Measure Scores from Testing or Current Use (<i>description of scores, e.g., distribution by quartile, mean, median, SD, etc.; identification of statistically significant and meaningfully differences in performance</i>): Mean Rate for All Hospitals = 76.2% Overall Rate for All Hospitals = 73.7% Standard Deviation = 26.9% Median Rate for All Hospitals = 87.8% Min. = 0.0% Max. = 100% Lower Quartile = 61.5% Upper Quartile = 97.5% Z < -2* = 2 Z < 2** = 0</p>	<p>2f</p> <p>C <input type="checkbox"/></p> <p>P <input type="checkbox"/></p> <p>M <input type="checkbox"/></p> <p>N <input type="checkbox"/></p>
<p>2g. Comparability of Multiple Data Sources/Methods</p> <p>2g.1 Data/sample (<i>description of data/sample and size</i>):</p> <p>2g.2 Analytic Method (<i>type of analysis & rationale</i>):</p> <p>2g.3 Testing Results (<i>e.g., correlation statistics, comparison of rankings</i>):</p>	<p>2g</p> <p>C <input type="checkbox"/></p> <p>P <input type="checkbox"/></p> <p>M <input type="checkbox"/></p> <p>N <input type="checkbox"/></p> <p>NA <input type="checkbox"/></p>

2h. Disparities in Care 2h.1 If measure is stratified, provide stratified results (<i>scores by stratified categories/cohorts</i>): 2h.2 If disparities have been reported/identified, but measure is not specified to detect disparities, provide follow-up plans:	2h C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/>
TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for <i>Scientific Acceptability of Measure Properties</i>?	2
Steering Committee: Overall, to what extent was the criterion, <i>Scientific Acceptability of Measure Properties</i>, met? Rationale:	2 C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
3. USABILITY	
Extent to which intended audiences (e.g., consumers, purchasers, providers, policy makers) can understand the results of the measure and are likely to find them useful for decision making. (evaluation criteria)	Eval Rating
3a. Meaningful, Understandable, and Useful Information 3a.1 Current Use: Not in use but testing completed 3a.2 Use in a public reporting initiative (disclosure of performance results to the public at large) (<i>If used in a public reporting initiative, provide name of initiative(s), locations, Web page URL(s). <u>If not publicly reported</u>, state the plans to achieve public reporting within 3 years</i>): We intend to incorporate these Patient Blood Management measures into our ORYX initiative with associated public reporting on Quality Check when there is a national call for measures. 3a.3 If used in other programs/initiatives (<i>If used in quality improvement or other programs/initiatives, name of initiative(s), locations, Web page URL(s). <u>If not used for QI</u>, state the plans to achieve use for QI within 3 years</i>): The specifications will be posted on the Joint Commission website for public use in 2011. Testing of Interpretability (<i>Testing that demonstrates the results are understood by the potential users for public reporting and quality improvement</i>) 3a.4 Data/sample (<i>description of data/sample and size</i>): 3a.5 Methods (<i>e.g., focus group, survey, QI project</i>): 3a.6 Results (<i>qualitative and/or quantitative results and conclusions</i>):	3a C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
3b/3c. Relation to other NQF-endorsed measures	
3b.1 NQF # and Title of similar or related measures:	
(for NQF staff use) Notes on similar/related endorsed or submitted measures:	
3b. Harmonization If this measure is related to measure(s) already endorsed by NQF (e.g., same topic, but different target population/setting/data source <u>or</u> different topic but same target population): 3b.2 Are the measure specifications harmonized? If not, why?	3b C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/>
3c. Distinctive or Additive Value 3c.1 Describe the distinctive, improved, or additive value this measure provides to existing NQF-endorsed measures:	3c C <input type="checkbox"/> P <input type="checkbox"/>

5.1 If this measure is similar to measure(s) already endorsed by NQF (i.e., on the same topic and the same target population), Describe why it is a more valid or efficient way to measure quality:	M <input type="checkbox"/> N <input type="checkbox"/> NA <input type="checkbox"/>
TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for <i>Usability</i> ?	3
Steering Committee: Overall, to what extent was the criterion, <i>Usability</i> , met? Rationale:	3 C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
4. FEASIBILITY	
Extent to which the required data are readily available, retrievable without undue burden, and can be implemented for performance measurement. (evaluation criteria)	Eval Rating
4a. Data Generated as a Byproduct of Care Processes	
4a.1-2 How are the data elements that are needed to compute measure scores generated? Data generated as byproduct of care processes during care delivery (Data are generated and used by healthcare personnel during the provision of care, e.g., blood pressure, lab value, medical condition), Coding/abstraction performed by someone other than person obtaining original information (E.g., DRG, ICD-9 codes on claims, chart abstraction for quality measure or registry)	4a C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
4b. Electronic Sources	
4b.1 Are all the data elements available electronically? (<i>elements that are needed to compute measure scores are in defined, computer-readable fields, e.g., electronic health record, electronic claims</i>) No	4b C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
4b.2 If not, specify the near-term path to achieve electronic capture by most providers. The project will begin Phase III in January 2011 to retool the specifications for retrieval from an electronic health record.	
4c. Exclusions	
4c.1 Do the specified exclusions require additional data sources beyond what is required for the numerator and denominator specifications? No	4c C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
4c.2 If yes, provide justification.	NA <input type="checkbox"/>
4d. Susceptibility to Inaccuracies, Errors, or Unintended Consequences	
4d.1 Identify susceptibility to inaccuracies, errors, or unintended consequences of the measure and describe how these potential problems could be audited. If audited, provide results.	4d C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>
4e. Data Collection Strategy/Implementation	
4e.1 Describe what you have learned/modified as a result of testing and/or operational use of the measure regarding data collection, availability of data/missing data, timing/frequency of data collection, patient confidentiality, time/cost of data collection, other feasibility/ implementation issues: Abstraction time for PBM-03 varied based on whether the patient received plasma and the number of plasma units transfused to each patient. Less plasma was transfused during testing, so the extra layer of abstraction by 'event' was not as critical to reliability. However, for consistency, all blood products will be abstracted by unit and the initial four plasma units that are transfused will be evaluated. There were	4e C <input type="checkbox"/> P <input type="checkbox"/> M <input type="checkbox"/> N <input type="checkbox"/>

similar issues related to the difficulty abstractors had in determining how to match the hospital indication with the pilot indication as mentioned for PBM-02 for the data element Clinical Indication for Plasma, but post pilot, hospitals will not have to categorize the indication to a pre-defined list of reasons. Intraoperatively, documentation of a blood transfusion pre-transfusion lab results and clinical indication was lacking in most paper-based records. So, in order to assist hospitals to focus their efforts on areas with low rates of compliance, this measure will be stratified so that hospitals can track results based on administration location. The “closest” PT/INR value or TEG will be abstracted without a “within 24 hour timeframe” requirement for consistency with the other transfusion measures.

Pilot hospitals were requested to estimate the time to abstract one unit of plasma for the six-month pilot. Twenty hospitals estimated an average time of 30 minutes to abstract a unit of blood with an average cost of \$21-25 per hour. However, these costs do not include the time or cost involved in identifying the patient population, staff training or data collection tool instruction. It should also be noted that the learning curve varied widely due to the staff experience and expertise that were utilized for a ‘time-limited’ project. Due to the amount of time needed to manually abstract the volume of blood transfusions, we believe that these measures are most suitable for abstraction from an electronic medical record (EHR). Retrieval from an EHR could capture 100% of all units that were transfused and would decrease or eliminate the associated abstraction burden. This method would also improve the identification of patients who received blood since procedure codes to document blood use are not standardized across the country. In the meantime, patients can be identified using blood bank records or procedure codes.

During the 12 reliability site visits, two Joint Commission staff also found that the abstraction time varied widely based on the method of record retrieval (e.g., paper record, scanned record or electronic information) at each hospital and the amount of blood transfused per case. Based on hospital feedback, measure specifications have been revised to strengthen and provide additional clarity to data element definitions and abstraction guidelines. The timing and frequency of data collection will remain monthly or quarterly as it does for the other Joint Commission measure sets. Maintaining patient confidentiality was not an issue during the pilot test, since blinded hospital and patient identifiers are used on all data received by The Joint Commission staff for data quality reviews.

4e.2 Costs to implement the measure (*costs of data collection, fees associated with proprietary measures*):

The majority of hospitals already have processes in place to abstract measures if the patients are identified using procedure codes. However, some hospitals document total hospital blood use using blood bank records that would have to be cross-referenced by the patient medical record number to determine how much and the type of blood product each patient received which adds to the abstraction burden. There are no Joint Commission fees to abstract the measures, but the abstraction cost for this measure would depend on the amount of blood transfused at each hospital. This measure would evaluate the first three units of plasma regardless of the number of plasma units transfused. Hospitals with Blood Management or conservation programs may have fewer units to review and those with efficient or electronic processes to document blood may have lower abstraction costs.

4e.3 Evidence for costs:

4e.4 Business case documentation: There continues to be considerable unexplained variation in transfusion practices across organizations, products and patient populations. Recent evidence is mounting that demonstrates significant harm from unnecessary blood transfusions. Monitoring transfusions will provide information so hospitals can begin to identify patients who are transfused outside of the guidelines. It has been found that hospitals that track blood use at the patient specific level have a higher percentage of appropriate transfusions than those that do not track blood use at that level. Measuring blood use should decrease the amount of blood transfused and improve patient safety.

TAP/Workgroup: What are the strengths and weaknesses in relation to the subcriteria for *Feasibility*?

4

Steering Committee: Overall, to what extent was the criterion, *Feasibility*, met?
Rationale:

4

C ☐
P ☐
M ☐
N ☐

RECOMMENDATION	
(for NQF staff use) Check if measure is untested and only eligible for time-limited endorsement.	Time-limited <input type="checkbox"/>
Steering Committee: Do you recommend for endorsement? Comments:	Y <input type="checkbox"/> N <input type="checkbox"/> A <input type="checkbox"/>
CONTACT INFORMATION	
Co.1 Measure Steward (Intellectual Property Owner) Co.1 Organization The Joint Commission, One Renaissance Boulevard, Oakbrook Terrace, Illinois, 60181 Co.2 Point of Contact Jerod M., Loeb, PhD, jloeb@jointcommission.org, 630-792-5920-	
Measure Developer If different from Measure Steward Co.3 Organization The Joint Commission, One Renaissance Boulevard, Oakbrook Terrace, Illinois, 60181 Co.4 Point of Contact Harriet, Gammon, MSN, RN, CPHQ, hgammon@jointcommission.org, 630-792-5926-	
Co.5 Submitter If different from Measure Steward POC Harriet, Gammon, MSN, RN, CPHQ, hgammon@jointcommission.org, 630-792-5926-, The Joint Commission	
Co.6 Additional organizations that sponsored/participated in measure development	
ADDITIONAL INFORMATION	
Workgroup/Expert Panel involved in measure development Ad.1 Provide a list of sponsoring organizations and workgroup/panel members' names and organizations. Describe the members' role in measure development. The technical advisory panel determined priority areas in blood management for measure development. They reviewed public comments and were actively involved in all phases of the project to identify and develop the numerator and denominator statements. Measure recommendations for National Quality Forum endorsement were made after careful review of the pilot results and site feedback.	
Ad.2 If adapted, provide name of original measure: Ad.3-5 If adapted, provide original specifications URL or attachment	
Measure Developer/Steward Updates and Ongoing Maintenance Ad.6 Year the measure was first released: Ad.7 Month and Year of most recent revision: 12, 2010 Ad.8 What is your frequency for review/update of this measure? Biannually Ad.9 When is the next scheduled review/update for this measure? 06, 2011	
Ad.10 Copyright statement/disclaimers: No royalty or use fee is required for copying or reprinting this manual, but the following are required as a condition of usage: 1) disclosure that the Specifications Manual is periodically updated, and that the version being copied or reprinted may not be up-to-date when used unless the copier or printer has verified the version to be up-to-date and affirms that, and 2) users participating in Joint Commission accreditation, including performance measures systems, are required to update their software and associated documentation based on the published manual production timelines. Example Acknowledgement: The Specifications Manual for National Hospital Inpatient Quality Measures Patient Blood Management Performance Measure Set is periodically updated by The Joint Commission. Users of the Specifications Manual for National Hospital Inpatient Quality Measures Patient Blood Management Performance Measure Set must update their software and associated documentation based on the published manual production timelines.	
Ad.11 -13 Additional Information web page URL or attachment: Attachment TAPLISTWEBc-	

[634276948397825978.doc](#)Date of Submission (MM/DD/YY): [12/29/2010](#)

Patient Blood Management (PBM)

Set Measures

Set Measure ID	Measure Short Name
<u>PBM-01</u>	Transfusion Consent
<u>PBM-02</u>	RBC Transfusion Indication
<u>PBM-03</u>	Plasma Transfusion Indication
<u>PBM-04</u>	Platelet Transfusion Indication
<u>PBM-05</u>	Blood Administration Documentation
<u>PBM-06</u>	Preoperative Anemia Screening
<u>PBM-07</u>	Preoperative Blood Type Testing and Antibody Screening

Measure Set Specific Data Elements

Element Name	Collected For
<u>Admission From Home</u>	<u>PBM-06</u> ,
<u>Anesthesia Start Date</u>	<u>PBM-06</u> ,
<u>Blood Administration Location</u>	<u>PBM-02</u> , <u>PBM-03</u> , <u>PBM-04</u> , <u>PBM-05</u> ,
<u>Blood Bank Records</u>	<u>PBM-01</u> , <u>PBM-02</u> , <u>PBM-03</u> , <u>PBM-04</u> , <u>PBM-05</u> ,
<u>Blood ID Number</u>	<u>PBM-05</u> ,
<u>Blood Type Testing Ordered</u>	<u>PBM-07</u> ,
<u>Clinical Indication for Plasma</u>	<u>PBM-03</u> ,
<u>Clinical Indication for Platelets</u>	<u>PBM-04</u> ,
<u>Clinical Indication for RBCs</u>	<u>PBM-02</u> ,
<u>Education Addressed Risks, Benefits and Alternatives to Transfusion</u>	<u>PBM-01</u> ,
<u>Patient ID Verification</u>	<u>PBM-05</u> ,
<u>Plasma ID</u>	<u>PBM-03</u> , <u>PBM-05</u> ,
<u>Platelet ID</u>	<u>PBM-04</u> , <u>PBM-05</u> ,
<u>Pre-transfusion Hematocrit</u>	<u>PBM-02</u> ,
<u>Pre-transfusion Hemoglobin</u>	<u>PBM-02</u> ,
<u>Pre-transfusion PT/INR Result</u>	<u>PBM-03</u> ,
<u>Pre-transfusion Platelet Count</u>	<u>PBM-04</u> ,
<u>Preoperative Anemia Screening Date</u>	<u>PBM-06</u> ,
<u>Preoperative Blood Type Testing</u>	<u>PBM-07</u> ,
<u>RBC ID</u>	<u>PBM-02</u> , <u>PBM-05</u> ,
<u>RBC Unit Exclusions</u>	<u>PBM-02</u> , <u>PBM-05</u> ,
<u>Surgery Scheduled Timeframe</u>	<u>PBM-06</u> ,
<u>Transfusion Consent</u>	<u>PBM-01</u> ,
<u>Transfusion Order</u>	<u>PBM-05</u> ,
<u>Transfusion Start Date</u>	<u>PBM-05</u> ,
<u>Transfusion Start Time</u>	<u>PBM-05</u> ,
<u>Vital Sign Monitoring</u>	<u>PBM-05</u> ,

Related Materials

Document Name
<u>z. Appendix E - Miscellaneous Tables</u>

Measure Information Form

Measure Set: Patient Blood Management(PBM)

Set Measure ID: PBM-01

Performance Measure Name: Transfusion Consent

Description: Patients with a signed consent who received information about the risks, benefits and alternatives of transfusion prior to the initial blood transfusion or the initial transfusion was deemed a medical emergency.

Rationale: Planning a discussion with a licensed practitioner regarding the risks, benefits and alternatives of transfusion is an opportunity for the patient to participate in decisions about his or her care. It is a process that takes into consideration, each patient's preferences, clinical needs and provides information in compliance with the regulations and policies of the state and facility. Even though policies related to informed consent may vary among hospitals, all hospitals require some type of consent prior to treatment unless emergency care is needed. The elements of performance for the Joint Commission Standard RI.01.03.01 related to the informed consent process include a discussion about the risks, benefits and alternatives, and a discussion about the risk, if care is not received. This measure is also supported by the Joint Commission's National Patient Safety Goal (NPSG) 13 that encourages patients' active involvement in their own care as a patient safety strategy.

For many years, the American Association of Blood Banks (AABB) organization has supported the consent process for transfusion and has developed several standards such as AABB Standard 5.19.1. AABB requires that at a minimum, a recipient consent for transfusion and that should include; a description of the risks, benefits and treatment alternatives, the opportunity to ask questions and the right to accept or refuse transfusion.

Type of Measure: Process

Improvement Noted As: Increase in the rate

Numerator Statement: Patients with a signed consent who received information about the risks, benefits and alternatives prior to the initial blood transfusion or the initial transfusion was deemed a medical emergency

Included Populations: Not applicable

Excluded Populations: None

Data Elements:

- Education Addressed Risks, Benefits and Alternatives to Transfusion
- Transfusion Consent

Denominator Statement: Patients who received red blood cell, plasma or platelet transfusions

Included Populations: Discharges with an ICD-9-CM Principal or Other Procedure Codes for transfusion as defined in Appendix A, Table 9.3-9.6 or a transfusion documented from Blood Bank Records.

Excluded Populations: None

Data Elements:

- Admission Date
- Blood Bank Records
- Discharge Date
- ICD-9-CM Other Procedure Codes
- ICD-9-CM Principal Procedure Code

Risk Adjustment: No.

Data Collection Approach: Retrospective data collection sources for required data elements include administrative data and medical records. Hospitals that do not use ICD-9-CM procedure codes to document transfusions may use blood bank records to identify the population.

Data Accuracy: Variation may exist in the assignment of ICD-9-CM codes and blood bank records; therefore, coding practices and transfusion documentation may require evaluation to ensure consistency.

Measure Analysis Suggestions: Hospitals may want to evaluate the cases according to medical or surgical designation that were not included in the numerator in order to determine if the consent was signed and/or if all or only part of the educational components were given or if documentation was insufficient. Based on this information, hospitals may assess the barriers impacting this measure that could be improved.

Sampling: Yes. For additional information see the Population and Sampling Specifications Section.

Data Reported As: Aggregate rate generated from count data reported as a proportion.

Selected References:


- Speiss BD, Counts RB, Gould SA. Perioperative Transfusion Medicine, Williams and Wilkins; 1998; 201-204.
- Stowell C, Sazama K. Informed Consent in Blood Transfusion and Cellular Therapies: Patients, Donors and Research Subjects. AABB Press; 2007; ISBN #978-1-56395-254-8.
- Burch JW, Uhl L. Guidelines for Informed Consent in Transfusion Medicine. AABB Press; 2006; ISBN #1-56395-146-0.2008.
- Standards for Blood Banks and Transfusion Services, 25th ed. Bethesda, MD: AABB 2008.
- The Joint Commission: Comprehensive Accreditation Manual for Hospitals, 2009. Oakbrook Terrace, IL. Joint Commission Resources, Inc, 2009.
- The Joint Commission, "National Patient Safety Goals (NPSG)", IN: Comprehensive accreditation manual for hospitals, 2009. Oakbrook Terrace, IL; Joint Commission Resources, Inc., 2009, pp. NPSG 1 – NPSG 4.

Measure Algorithm:

PBM-01: Transfusion Consent

Numerator: Patients with a signed consent who received information about the risks, benefits and alternatives prior to the initial blood transfusion or the initial transfusion was deemed a medical emergency

Denominator: Patients who received red blood cells, platelets or plasma



Related Topics

Measure Information Form

Measure Set: Patient Blood Management(PBM)

Set Measure ID: PBM-02

Performance Measure Name: RBC Transfusion Indication

Description: The number of transfused red blood cell (RBC) units with a pre-transfusion hemoglobin (hgb) or hematocrit (hct) result and clinical indication documented from patients of all ages who received RBCs.

Rationale: Improvement of the safety and quality of care that a hospital provides includes the review of the use of blood and blood products. Despite current evidence and best practice guidelines, clinical practice regarding when to transfuse varies among physicians and institutions even though most would agree that blood products should only be given when the benefits outweigh the harm. Many advocate that transfusion decisions should be based on a clinical assessment and not on laboratory values alone to avoid inappropriate over-or-under transfusion. Measuring whether an “indication for transfusion” and a pre-transfusion laboratory value was documented may improve the utilization of blood components. In addition, implementing such a process may simplify the hospital’s review for appropriateness of the transfusion when auditing records for accreditation and regulatory agencies. In a study by Friedman and Ebrahim, there was a significant correlation between red blood cell transfusions that lacked documentation of the clinical necessity for transfusion and justification of the transfusion.

Type of Measure: Process

Improvement Noted As: Increase in the rate

Numerator Statement: Number of RBC units with pre-transfusion hemoglobin or hematocrit result and clinical indication documented

Included Populations: Not applicable

Excluded Populations: None

Data Elements:

- Clinical Indication for RBCs
- Pre-transfusion Hematocrit
- Pre-transfusion Hemoglobin
- RBC ID

Denominator Statement: Number of transfused red blood cell units evaluated

Included Populations:

- Discharges with an ICD-9-CM Principal or Other Procedure Codes for transfusion as defined in Appendix A, Tables 9.3 or 9.4 or a RBC transfusion documented from Blood Bank Records.
- The first six RBCs units transfused after hospital arrival

Excluded Populations: None

Data Elements:

- Admission Date
- Birthdate
- Blood Administration Location
- Blood Bank Records
- Discharge Date
- ICD-9-CM Other Procedure Codes
- ICD-9-CM Principal Procedure Code
- RBC Unit Exclusions

Risk Adjustment: No.

Data Collection Approach: Retrospective data sources for required data elements include administrative/billing data and medical records. Hospitals that do not use ICD-9-CM procedure codes to document transfusions may use blood bank records to identify the population of patients who received RBCs.

Data Accuracy: Variation may exist in the assignment of ICD-9-CM codes and blood bank records; therefore, coding practices and transfusion documentation may require evaluation to ensure consistency.

Measure Analysis Suggestions: Hospitals may want to use the data to further evaluate the process for determining the need for blood products based on the clinical indications and correlating it with the pre-transfusion value that was documented. This information may assist hospitals to determine if the patients were transfused appropriately or if efforts should be directed toward additional documentation efforts for monitoring blood product usage. Data may be grouped by service designation or by blood products to identify specific areas for staff review.

Sampling: Yes. For additional information see the Population and Sampling Specifications Section.

Data Reported As: Aggregate rate generated from count data reported as a proportion.

Selected References:


- Friedman MT, Ebrahim A. Adequacy of physician documentation of red blood cell transfusion and correlation with assessment of transfusion appropriateness. Arch Pathol Lab Med. 2006;130: 474-79.
- Corwin HL, Parsonnet KC, Gettinger A. RBC transfusion in the ICU: is there a reason? Chest. 1995;108: 767-771.
- Tobin SN, Campbell DA, Boyce NW. Durability of response to a targeted intervention to modify clinician transfusion practices in a major teaching hospital. MJA. 2001;174:445-448.
- Clinical practice guideline: Red blood cell transfusion in adult trauma and critical care. Crit Care Med 2009 Vol.37, No.12.


Measure Algorithm:


PBM-02: RBC Transfusion Indication

Numerator: Number of RBC units (bags) with pre-transfusion hemoglobin or hematocrit result and clinical indication documented

Denominator: Number of transfused red blood cell units evaluated







Related Topics

Measure Information Form

Measure Set: Patient Blood Management(PBM)

Set Measure ID: PBM-03

Performance Measure Name: Plasma Transfusion Indication

Description: The number of transfused plasma units with a pre-transfusion PT/INR result and clinical indication documented from patients of all ages who received plasma.

Rationale: The use of plasma has increased and is disproportionately high compared to other countries with similar levels of health care. Indications for transfusing plasma are very limited, and as a result, published studies often show unjustifiable use of plasma. According to the National Heart Lung and Blood Institute, plasma should be administered only to increase the level of clotting factors in patients with a demonstrated deficiency. If the prothrombin time (PT) and partial thromboplastin time (PTT) are < 1.5 times normal, a plasma transfusion is rarely needed. However, plasma is frequently transfused to patients with mild-to moderate elevations in PT despite numerous studies that have not shown a correlation between the risk of bleeding and mild-to moderate test results. In a study by Wahab et al, transfusion of plasma for mild abnormalities of coagulation values resulted in a partial normalization in a minority of patients, and failed to correct the PT in 99% of the patients. In a 2004 study by Hui, the need to correct prolonged international normalized ratios (INRs) for patients on warfarin emerged as the primary indication for plasma followed by massive transfusions.

Type of Measure: Process

Improvement Noted As: Increase in the rate

Numerator Statement: Number of plasma units with pre-transfusion PT/INR result and clinical indication documented

Included Populations: Not applicable

Excluded Populations: None

Data Elements:

- Clinical Indication for Plasma
- Plasma ID
- Pre-transfusion PT/INR Result

Denominator Statement: Number of transfused plasma units evaluated

Included Populations:

- Discharges with an ICD-9-CM Principal or Other Procedure Codes for transfusion as defined in Appendix A, Table 9.6 or a plasma transfusion documented from Blood Bank Records
- The first three plasma units transfused from hospital arrival

Excluded Populations:

- Discharges with an ICD-9-CM Principal Diagnosis Code of trauma as defined in Appendix A, Table 9.7.

Data Elements:

- Admission Date
- Birthdate
- Blood Administration Location
- Blood Bank Records
- Discharge Date
- ICD-9-CM Other Procedure Codes
- ICD-9-CM Principal Diagnosis Code
- ICD-9-CM Principal Procedure Code

Risk Adjustment: No.

Data Collection Approach: Retrospective data sources for required data elements include administrative/billing data and medical records. Hospitals that do not use ICD-9-CM procedure codes to document transfusions may use blood bank records to identify the population of patients who received plasma.

Data Accuracy: Variation may exist in the assignment of ICD-9-CM codes and blood bank records; therefore, coding practices and transfusion documentation may require evaluation to ensure consistency.

Measure Analysis Suggestions: Data from this measure may be used to review the type of invasive procedures or surgeries that use plasma in order to further evaluate appropriateness of use.

Sampling: Yes. For additional information see the Population and Sampling Specifications Section.

Data Reported As: Aggregate rate generated from count data reported as a proportion.

Selected References:


- Hui C, Williams I, Davis K. Clinical audit of the use of fresh-frozen plasma and platelets in a tertiary teaching hospital and the impact of a new transfusion request form. *Int Med J*. 2005;35:283-288.
- Wallis JP, Dzik S. Is fresh frozen plasma overtransfused in the United States? *Transfusion*. 2004;44:1674-75.
- Ardel-Wahab OI, Healy B, Dzik WH. Effect of fresh-frozen plasma transfusion on prothrombin time and bleeding in patients with mild coagulation abnormalities. *Transfusion*. 2006;46:1479-1285.
- Segal J, Dzik WH; Transfusion Medicine/Hemostasis Clinical Trials Network. Paucity of studies to support that abnormal coagulation test results predict bleeding in the setting of invasive procedures: an evidenced-based review. *Transfusion*. 2005;45:1413-25.


Measure Algorithm:


PBM-03: Plasma Transfusion Indication

Numerator: Number of plasma units with pre-transfusion PT/INR result and clinical indication documented

Denominator: Number of transfused plasma units evaluated







Related Topics

Measure Information Form

Measure Set: Patient Blood Management(PBM)

Set Measure ID: PBM-04

Performance Measure Name: Platelet Transfusion Indication

Description: The number of transfused platelet units with pre-transfusion platelet count and clinical indication documented from patients of all ages who received platelets.

Rationale: Platelets are transfused to treat or prevent bleeding associated with thrombocytopenia and/or platelet dysfunction. Platelets given therapeutically should help stop the bleeding, and if given prophylactically, post transfusion platelet counts should be obtained to monitor the response to determine the effectiveness of the transfusion. Repeated platelet transfusions can cause alloimmunization and cause platelet refractoriness to future transfusions. Multiple infectious risks are associated with platelet transfusions so patients should only be exposed to the least amount needed.

Type of Measure: Process

Improvement Noted As: Increase in the rate

Numerator Statement: Number of platelet units with pre-transfusion platelet count result and clinical indication documented

Included Populations: Not applicable

Excluded Populations: None

Data Elements:

- Clinical Indication for Platelets
- Platelet ID
- Pre-transfusion Platelet Count

Denominator Statement: Number of transfused platelet units evaluated

Included Populations:

- Discharges with an ICD-9-CM Principal or Other Procedure Codes for transfusion as defined in Appendix A, Table 9.5 or a platelet transfusion documented from Blood Bank Records
- The first three platelet units transfused after hospital arrival

Excluded Populations: None

Data Elements:

- Admission Date
- Blood Administration Location
- Blood Bank Records

- Discharge Date
- ICD-9-CM Other Procedure Codes
- ICD-9-CM Principal Procedure Code

Risk Adjustment: No.

Data Collection Approach: Retrospective data sources for required data elements include administrative/billing data and medical records. Hospitals that do not use ICD-9-CM procedure codes to document transfusions may use blood bank records to identify the population of patients who received platelets.

Data Accuracy: Variation may exist in the assignment of ICD-9-CM codes and blood bank records; therefore, coding practices and transfusion documentation may require evaluation to ensure consistency.

Measure Analysis Suggestions: Data from this measure may be used to evaluate the utilization and appropriateness of platelets used by an organization.

Sampling: Yes. For additional information see the Population and Sampling Specifications.

Data Reported As: Aggregate rate generated from count data reported as a proportion.

Selected References:


- Garrioch M, Sandbach J, Pirie E, Morrison A, Todd A, Green R. Reducing red cell transfusion by audit, education and a new guideline in a large teaching hospital. *Transfusion Med.* 2004;14:25-31.
- Petrides M. Red cell transfusion “trigger”: A review. *Southern Med J.* 2003; 96:664-667.
- Roback JD, ed. Technical manual. 16th ed, Bethesda, MD: AABB, 2008.
- BR J Haematol 1998, 101:609 - 617.


Measure Algorithm:


PBM-04: Platelet Transfusion Indication

Numerator: Number of platelet doses with pre-transfusion platelet count result and clinical indication documented

Denominator: Number of transfused platelet units evaluated







Related Topics

Measure Information Form

Measure Set: Patient Blood Management(PBM)

Set Measure ID: PBM-05

Performance Measure Name: Blood Administration Documentation

Description: The number of transfused red blood cells, plasma or platelet transfusion units/doses (bags) that had documentation of the following: patient identification and an order to transfuse (Blood ID Number) confirmed prior to the initiation of transfusion, transfusion start date and time, and blood pressure, pulse and temperature recorded at specific intervals.

Rationale: Since the majority of blood units are transfused in hospitals, specific policies and procedures have been developed by each hospital to address documentation of blood administration standards in accordance with their state and federal regulations. Though documentation components vary among organizations, identification of the patient and confirmation of the order to transfuse are common indicators used for all blood products since incomplete patient identification could result in an adverse outcome. Prior to administering blood or blood products, patient identification by two identifiers is required by numerous organizations including the AABB Standard 5.19.3, and the Joint Commission National Patient Safety Goal (NPSG) 1. In addition, numerous organizations require or advise that the licensed staff confirm that there is a transfusion order as directed by the AABB Standard 5.19.6 and the elements of performance for the Joint Commission NPSG.01.01.01.

Patient monitoring during the transfusion is an important component related to patient safety. The first 10 to 15 minutes of the transfusion are considered the most critical to assess for a potential transfusion reaction and close observation during this time is recommended in the AABB Primer. Monitoring of vital signs at baseline, during and at the completion of the transfusion in addition to observation are used to assess the patient's condition for any changes.

Type of Measure: Process

Improvement Noted As: Increase in the rate

Numerator Statement: Number of units/doses (bags) with documentation for all of the following:

- patient identification and transfusion order (Blood ID Number) confirmed prior to the initiation of transfusion
- transfusion start date and time
- blood pressure, pulse and temperature recorded pre, during and post transfusion

Included Populations: Not applicable

Excluded Populations: None

Data Elements:

- Blood ID Number
- Patient ID Verification
- Plasma ID

- Platelet ID
- RBC ID
- Transfusion Order
- Transfusion Start Date
- Transfusion Start Time
- Vital Sign Monitoring

Denominator Statement: Number of transfused red blood cells, plasma or platelet units/doses (bags) evaluated

Included Populations:

- Discharges with an ICD-9-CM Principal or Other Procedure Codes for transfusion as defined in Appendix A, Table 9.3-9.6 or a transfusion documented from Blood Bank Records

Excluded Populations:

- Units used in massive transfusion protocols
- Uncrossmatched units
- Units used to prime equipment

Data Elements:

- Admission Date
- Birthdate
- Blood Administration Location
- Blood Bank Records
- Discharge Date
- ICD-9-CM Other Procedure Codes
- ICD-9-CM Principal Procedure Code
- RBC Unit Exclusions

Risk Adjustment: No.

Data Collection Approach: Retrospective data sources for required data elements include administrative/billing data and medical records. Hospitals that do not use ICD-9-CM procedure codes to document transfusions may use blood bank records to identify the population.

Data Accuracy: Variation may exist in the assignment of ICD-9-CM codes and blood bank records; therefore, coding practices and transfusion documentation may require evaluation to ensure consistency.

Measure Analysis Suggestions: The data from this measure may be used to evaluate the adherence to organizational policies and procedures for blood administration for each of the blood products. Data could be evaluated by unit or service in order to identify areas for staff education. The data could also be used during accreditation surveys to document the hospital's efforts to improve the accuracy of patient identification when administering blood related to the Joint Commission National Patient Safety Goal #1.

Sampling: Yes. For additional information see the Population and Sampling Specifications.

Data Reported As: Aggregate rate generated from count data reported as a proportion.

Selected References:

- Whitsett CF, Robichaux MG. Assessment of blood administration procedures: problems identified by direct observation and administrative incident reporting. *Transfusion*. 2001;41:581-86.
- Saxena S, Ramer L, Shulman IA. A comprehensive assessment program to improve blood-administering practices using the FOCUS-PDCA model. *Transfusion*. 2004; 44:1350-56.
- Novis DA, Miller KA, Howanitz PJ, Renner SW, Walsh MK; College of American Pathologists. Audit of transfusion procedures in 660 hospitals. A College of American Pathologists Q-Probes study of patient identification and vital sign monitoring frequencies in 16494 transfusions. *Arch Pathol Lab Med*. 2003;127:541-8.
- Roback JD, ed. Technical manual. 16th ed, Bethesda, MD: AABB, 2008.
- The Joint Commission: Comprehensive Accreditation Manual for Hospitals, 2009. Oakbrook Terrace, IL; Joint Commission Resources, Inc., 2009.
- The Joint Commission, "National Patient Safety Goals (NPSG)", IN: Comprehensive accreditation manual for hospitals, 2009. Oakbrook Terrace, IL; Joint Commission Resources, Inc., 2009, pp. NPSG 1 – NPSG 4.
- AABB Primer of Blood Administration. Revised August 2008. Bethesda, Maryland. [Available at http://www.aabb.org/Content/Professional_Development/Education_and_Training_Material/edtr (accessed November 2009).]


Measure Algorithm:


PBM-05: Blood Administration Documentation


Numerator: Number of blood transfusion units (bags) or doses with documentation for all of the following:

- patient identification (ID) and transfusion order (blood ID number) confirmed prior to the initiation of blood
- date and time of transfusion
- blood pressure, pulse and temperature recorded pre, during and post transfusion

Denominator: Number of transfused red blood cells, plasma and platelet units (bags) or doses evaluated







Related Topics

Measure Information Form

Measure Set: Patient Blood Management(PBM)

Set Measure ID: PBM-06

Performance Measure Name: Preoperative Anemia Screening

Description: Selected elective orthopedic, cardiac and hysterectomy surgical patients with documentation of preoperative anemia screening date 14 – 45 days before surgery start date for procedures scheduled 14 or more days before surgery.

Rationale: Development of formal protocols for preoperative testing of hemoglobin (hgb) for potential high-blood loss elective surgeries could be used to identify and intervene for optimal management of blood resources. Preoperative anemia often goes unrecognized and untreated unless tests are ordered in advance of a planned surgery. Early recognition of anemia offers patients an opportunity to receive the most appropriate transfusion-sparing strategy, and avoid the risk of a potential transfusion. Researchers have shown that preoperative hgb and hematocrit can be used as predictors of outcome for specific types of patients such as cardiac artery bypass graft or orthopedic surgery. In a study by Salido, orthopedic patients with a preoperative hemoglobin <13 g/dL had four times the risk of transfusion than those with a hemoglobin level between 13 g/dL and 15 g/dL.

Type of Measure: Process

Improvement Noted As: Increase in the rate

Numerator Statement: Patients with preoperative anemia screening 14 - 45 days before Anesthesia Start Date

Included Populations: Not applicable

Excluded Populations: None

Data Elements:

- Preoperative Anemia Screening Date

Denominator Statement: Selected elective surgical patients

Included Populations:

- Discharges with an ICD-9-CM Principal Procedure Codes of selected surgeries as defined in Appendix A, Tables 2.2, 5.01, 5.02, 5.08, 5.11, 5.22, 5.23, 9.1 or 9.2.

Excluded Populations:

- Patients less than 18 years of age
- Patients with surgery scheduled less than 14 days before Anesthesia Start Date
- Patients not admitted from home

Data Elements:

- Admission Date
- Admission From Home
- Birthdate
- Discharge Date
- ICD-9-CM Principal Procedure Code
- ICD-9-CM Principal Procedure Date
- Surgery Scheduled Timeframe

Risk Adjustment: No.

Data Collection Approach: Retrospective data sources for required data elements include administrative data and medical records.

Data Accuracy: Variation may exist in the assignment of ICD-9-CM codes; therefore, coding practices may require evaluation to ensure consistency.

Measure Analysis Suggestions: These data may be used to evaluate specific patient groups at high risk for a blood transfusion that did not have their pre-operative hemoglobin and/or transfusion testing completed and/or documented prior to surgery. The data could be further analyzed based on physician or type of procedure. Patients who are not included in the numerator could be tracked to see if there were any adverse outcomes due to the lack of preoperative anemia screening.

Sampling: Yes. For additional information see the Population and Sampling Specifications Section.

Data Reported As: Aggregate rate generated from count data reported as a proportion.

Selected References: * Roback JD, ed. Technical manual. 16th ed, Bethesda, MD: AABB, 2008.


- Salido JA, Martin LA, Gomez LA, et al. Preoperative hemoglobin levels and the need for transfusion after prosthetic hip and knee surgery; analysis of predictive factors. J Bone Joint Surg. 2002;84: 216-20.
- Rady MY, Ryan T, Starr NJ. Perioperative determinants of morbidity and mortality in elderly patients undergoing cardiac surgery. Crit Care Med. 1998;26: 225-235.
- Magovern JA, Sakert T, Magovern GJ et al. A model that predicts morbidity and mortality after coronary artery bypass graft surgery. J Am Coll Cardiol. 1996;28: 1147-1153.
- Campbell DA, Henderson WG, Englesbe, MJ, Hall BL, O'Reilly M, Bratzler D et al. Surgical site infection prevention: the importance of operative duration and blood transfusion-results of the first american college of surgeons –national surgical quality improvement program best practices initiative. J AM Coll Surg 2008;207:810-820.


Measure Algorithm:

PBM-06: Preoperative Anemia Screening

Numerator: Patients with documentation of preoperative anemia screening 14 - 45 days before Anesthesia Start Date

Denominator: Selected elective surgical patients





Related Topics

Measure Information Form

Measure Set: Patient Blood Management(PBM)

Set Measure ID: PBM-07

Performance Measure Name: Preoperative Blood Type Testing and Antibody Screening

Description: Selected elective orthopedic, cardiac and hysterectomy surgical patients who had preoperative blood type testing and antibody screening (type and screen or type and crossmatch) completed prior to surgery start time if ordered preoperatively.

Rationale: Hospitals need to ensure that sufficient compatible blood is available for each scheduled procedure. Since about 3% of specimens have a serologic finding that requires further investigation that may cause a delay in the availability of the blood, patient screening of ABO group and Rh type should be collected in sufficient time to complete all pretransfusion testing before surgery begins. According to the Joint Commission's Pre-publication National Patient Safety Goal UP.01.01.01 for 2010, a preprocedure verification process should be conducted to identify items that must be available for the procedure and use a standardized list to verify their availability. Documentation of any required blood products for the procedure is required. Development of formal protocols to ensure that patients have blood testing completed prior to surgery start time for potential high-blood loss elective surgeries may optimize management of blood resources and maximize patient safety.

Type of Measure: Process

Improvement Noted As: Increase in the rate

Numerator Statement: Patients with preoperative type and crossmatch or type and screen completed prior to surgery start time

Included Populations: Not applicable

Excluded Populations: None

Data Elements:

- Preoperative Blood Type Testing

Denominator Statement: Selected elective surgical patients

Included Populations:

- Discharges with an ICD-9-CM Principal Procedure Code of selected surgeries as defined in Appendix A, Tables 2.2, 5.01, 5.02, 5.08, 5.11, 5.22, 5.23, 9.1 or 9.2.

Excluded Populations:

- Patients less than 18 years of age
- Patients with type and screen or type and crossmatch ordered preoperatively

Data Elements:

- Admission Date
- Birthdate
- Blood Type Testing Ordered
- Discharge Date
- ICD-9-CM Principal Procedure Code

Risk Adjustment: No.

Data Collection Approach: Retrospective data collection sources for required data elements include administrative data and medical records.

Data Accuracy: Variation may exist in the assignment of ICD-9-CM codes; therefore, coding practices may require evaluation to ensure consistency.

Measure Analysis Suggestions: These data may be used to evaluate specific patient groups at high risk for a blood transfusion that did not have pre-operative transfusion testing completed and/or documented prior to surgery start time. The data could be further analyzed based on physician or type of procedure. Patients who are not included in the numerator could be tracked to see if there were any adverse outcomes due to the lack of preoperative testing.

Sampling: Yes. For additional information see the Population and Sampling Specifications.

Data Reported As: Aggregate rate generated from count data reported as a proportion.

Selected References: * Saxena S, Nelson JM, Osby M, Shah M, Kempf R, Shulman IA. Ensuring timely completion of type and screen testing and the verification of ABO/Rh status for elective surgical patients. Arch Pathol Lab Med. 2007;131:576-81.


- Friedberg RC, Jones BA, Walsh MK. Type and screen completion for scheduled surgical procedures. A College of American Pathologists Q-Probes study of 8941 type and screen tests in 108 institutions. Arch Pathol Lab Med. 2003;127:533-40.
- Roback JD, ed. Technical manual. 16th ed, Bethesda, MD: AABB, 2008.
- Magovern JA, Sakert T, Magovern GJ et al. A model that predicts morbidity and mortality after coronary artery bypass graft surgery. J Am Coll Cardiol. 1996;28: 1147-1153.
- The Joint Commission 2010 National Patient Safety Goals, Oakbrook Terrace, IL [Available at http://www.jointcommission.org/NR/rdonlyres/868C9E07-037F-433D-8858-0D5FAA4322F2/0/RevisedChapter_HAP_NPSG_20090924.pdf (accessed January 27, 2010).]

Measure Algorithm:

PBM-07: Preoperative Blood Type Testing and Antibody Screening

Numerator: Patients with documentation of preoperative type and crossmatch or type and screen completed prior to Anesthesia Start Time

Denominator: Selected elective surgical patients



Related Topics

Data Element Name: *Admission From Home*

Collected For: PBM-06,

Definition: Patient was admitted for the pre-scheduled elective surgery procedure from home.

Suggested Data Collection Question: Was the patient admitted from home?

Format: **Length:** 1
 Type: Alphanumeric
 Occurs: 1

Allowable Values:

- 1 Patient was admitted from home.
- 2 Patient was not admitted from home or unable to determine from medical record documentation.

Notes for Abstraction:

- Patients who have to stay overnight at a location other than their primary residence due to long distance travel for procedure are considered admitted from home.

Suggested Data Sources:

- Face sheet
- Nursing admission assessment
- Physician's notes
- Preop checklist

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name:	<i>Anesthesia Start Date</i>
Collected For:	<u>PBM-06</u> ,
Definition:	The date the anesthesia for the procedure started.
Suggested Data Collection Question:	On what date did the anesthesia for the procedure start?
Format:	Length: 10 – MM-DD-YYYY (includes dashes) Type: Date Occurs: 1
Allowable Values:	MM-DD-YYYY MM = Month (01-12) DD = Day (01-31) YYYY = Year (2001-Current Year) Leave Blank if Unable to Determine
Notes for Abstraction:	<p>If the Anesthesia Start Date cannot be determined from medical record documentation, enter UTD. When the date documented is obviously invalid (not a valid format/range [12-39-20xx] or after the Discharge Date or Anesthesia End Date) and no other documentation can be found that provides the correct information, the abstractor should select “UTD.”</p> <p>Example: Patient expires on 02-12-20xx and documentation indicates the Anesthesia Start Date was 03-12-20xx. Other documentation in the medical record supports the date of death as being accurate, but no other documentation of the Anesthesia Start Date can be found. Since the Anesthesia Start Date is outside of the parameter for care (after the Discharge Date [death]) and no other documentation is found, the abstractor should leave blank.</p> <p>If the Anesthesia Start Date is incorrect (in error) but it is a valid date and the correct date can be supported with other documentation in the medical record, the correct date may be entered. If supporting documentation of the correct date cannot be found, the medical record must be abstracted as documented or at “face value.”</p> <p>Examples: The anesthesia form is dated 12-10-2007, but other documentation in the medical record supports that the correct date was 12-10-2009. Enter the correct date of 12-10-2009 as the Anesthesia Start Date.</p> <p>An Anesthesia End Date of 11-20-20xx is documented but the Anesthesia Start Date is documented as 11-10-20xx. If no other documentation can be found to support another Anesthesia Start Date, then it must be abstracted as 11-10-20xx because the date is not considered invalid or outside the parameter of care.</p>

Suggested Data Sources:

Other Suggested Sources:

- Intraoperative record
- Circulator record
- Post-anesthesia evaluation record
- Operating room notes

Additional Notes: Suggested Data Sources:

Note: The anesthesia record is the priority data source for this data element, if a valid Anesthesia Start Date is found on the anesthesia record, use that date. If a valid date is not on the anesthesia record, other suggested data sources may be used in no particular order to determine the Anesthesia Start Date.

Priority Source:

- Anesthesia record

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Blood Administration Location*

Collected For: PBM-02, PBM-03, PBM-04, PBM-05,

Definition: The hospital setting (intraoperative or non-intraoperative) where the blood product began infusing.

Suggested Data Collection Question: In what setting did the blood product begin infusing?

Format:
Length: 1
Type: Alphanumeric
Occurs: 1-12

Allowable Values:

- 1 Intraoperative setting
- 2 Non-intraoperative setting
- 3 Unable to determine

Notes for Abstraction:

- Select setting for each unit transfused based on the physical location of the patient.
- Intraoperative setting is anytime during the operation.
- Non-intraoperative setting is any area outside of the operating room. For example, setting such as the intensive care unit, surgical floor or emergency room.

Suggested Data Sources:

- Anesthesia record
- Emergency department record
- Nursing notes
- Nursing flow sheet
- Nursing admission assessment
- Progress notes
- Physician's notes
- Operative notes
- Operating room notes
- Operative report
- Procedure notes
- ICU notes
- PACU/recovery room record

Blood Administration Documentation Sheet

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Blood Bank Records*

Collected For: PBM-01, PBM-02, PBM-03, PBM-04, PBM-05,

Definition: Documentation that the patient received red blood cells (RBCs), plasma or platelets after hospital arrival.

Suggested Data Collection Question: Was there documentation that the patient received RBCs, plasma or platelets after hospital arrival?

Format:
Length: 1
Type: Alphanumeric
Occurs: 1-12

Allowable Values:

- Select all that apply: 1 RBCs
2 Plasma
3 Platelets
4 None of the above or unable to determine from medical record documentation

Notes for Abstraction:

- Include transfusions given in the emergency room or observation area.

Suggested Data Sources: Blood Bank Records

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion

Data Element Name: *Blood ID Number*

Collected For: PBM-05,

Definition: Documentation of the actual blood bank identification number in the intraoperative record for the unit that was transfused.

Suggested Data Collection Question: Was there documentation of a blood bank identification number for the unit or dose of blood transfused during surgery?

Format:
Length: 1
Type: Alphanumeric
Occurs: 1

Allowable Values:

- 1 There is documentation of a blood bank identification number for the unit that was transfused.
- 2 There is no documentation of a blood bank identification number for the unit that was transfused or unable to determine from medical record documentation.

Notes for Abstraction:

Suggested Data Sources:

- Anesthesia record
- Operative report

Blood administration record

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Blood Type Testing Ordered*

Collected For: PBM-07,

Definition: A type and screen and/or type and crossmatch was ordered preoperatively for the elective surgery.

Suggested Data Collection Question: Was a type and screen and/or type and crossmatch ordered preoperatively?

Format:
Length: 1
Type: Alphanumeric
Occurs: 1

Allowable Values:

- 1 A type and screen and/or type and crossmatch was ordered preoperatively.
- 2 A type and screen and/or type and crossmatch was not ordered preoperatively or unable to determine

Notes for Abstraction:

Suggested Data Sources:

- Physician orders
- Preop checklist

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Clinical Indication for Plasma*

Collected For: PBM-03,

Definition: Documentation by the physician/advance practice nurse/physician assistant or (physician/APN/PA) of the clinical indication for the plasma transfusion unit.

Suggested Data Collection Question: Was there a clinical indication documented by the physician/APN/PA for the transfused plasma unit?

Format:
Length: 1
Type: Numeric
Occurs: 1 - 3

Allowable Values:

- 1 There was a clinical indication documented by the physician/APN/PA for the transfused plasma unit.
- 2 There was no documentation of a clinical indication for the transfusion or unable to determine from the medical record.

Notes for Abstraction:

- The clinical indication for the transfusion must be documented within 24 hours after the start of the transfusion.
- Select the first four plasma transfusion units closest to hospital arrival for abstraction.

Suggested Data Sources:

ONLY PHYSICIAN/APN/PA DOCUMENTATION OF THE CLINICAL INDICATION FOR ADMINISTERING BLOOD:

- Anesthesia record
- Consultation notes
- Emergency department record
- Physician orders
- Progress notes

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Clinical Indication for Platelets*

Collected For: PBM-04,

Definition: Documentation by the physician/advance practice nurse/physician assistant (physician/APN/PA) of the clinical indication for the transfused platelet unit.

Suggested Data Collection Question: Was there a clinical indication documented by the physician/APN/PA for the transfused platelet unit?

Format:
Length: 1
Type: Numeric
Occurs: 1 - 3

Allowable Values:

- 1 There was a clinical indication documented by the physician/APN/PA for the transfused platelet unit.
- 2 There was no documentation of clinical indication for the platelet transfusion or unable to determine from the medical record

Notes for Abstraction:

- The clinical indication for the transfusion must be documented within 24 hours after the start of the transfusion.
- Select the first three units transfused after hospital arrival for abstraction.

Suggested Data Sources:

ONLY PHYSICIAN/APN/PA DOCUMENTATION OF THE CLINICAL INDICATION FOR ADMINISTERING PLASMA:

- Anesthesia record
- Consultation notes
- Emergency department record
- Physician orders
- Progress notes

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Clinical Indication for RBCs*

Collected For: PBM-02,

Definition: Documentation by the physician/advance practice nurse/physician assistant (physician/APN/PA) of the clinical indication for the transfused red blood cell (RBCs) unit.

Suggested Data Collection Question: Was there a clinical indication documented by the physician/APN/PA for the transfused RBC unit?

Format:
Length: 1
Type: Numeric
Occurs: 1 - 6

Allowable Values:

- 1 There was a clinical indication documented by the physician/APN/PA for the transfused RBC unit.
- 2 There was no clinical indication documented by the physician/APN/PA for the transfused RBC unit or unable to determine from medical record documentation.

Notes for Abstraction:

- The clinical indication for the transfusion must be documented within 24 hours after the start of the transfusion.
- Select the first six RBC transfusion units after hospital arrival for abstraction.

Suggested Data Sources:

ONLY PHYSICIAN/APN/PA DOCUMENTATION OF THE CLINICAL INDICATION FOR ADMINISTERING RBCs:

- Anesthesia record
- Consultation notes
- Emergency department record
- Operative notes
- Physician orders
- Progress notes

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name:	<i>Education Addressed Risks, Benefits and Alternatives to Transfusion</i>
Collected For:	<u>PBM-01</u> ,
Definition:	Documentation that information addressing risks, benefits and alternatives to transfusion was given to the patient/caregiver prior to the initial transfusion or the initial transfusion was deemed a medical emergency after hospital arrival.
Suggested Data Collection Question:	Was there documentation that information regarding risks, benefits and alternatives to transfusion was given to the patient/caregiver prior to the initial transfusion event or was the initial transfusion deemed a medical emergency after hospital arrival?
Format:	Length: 1 Type: Numeric Occurs: 1
Allowable Values:	<ol style="list-style-type: none"> 1 Information addressing the risks, benefits and alternatives to transfusion was given to the patient/caregiver prior to the initial transfusion after hospital arrival. 2 Information addressing the risks, benefits and alternatives to transfusion was not given to the patient/caregiver prior to the initial transfusion after hospital arrival or unable to determine from medical record documentation.
Notes for Abstraction:	<ul style="list-style-type: none"> • Use only documentation provided in the medical record. • If the patient refused information about risks, benefits and alternatives to transfusion, select “1.” • The caregiver is defined as the patient’s family or any other person (e.g., guardian) who will be responsible for care of the patient.
Suggested Data Sources:	<ul style="list-style-type: none"> • Consultation notes • Emergency department record • History and physical • Nursing notes • Progress notes • Operative notes • Admission forms • Consent form • Emergency department record • Progress notes • Nursing notes
Additional Notes:	

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name:	<i>Patient ID Verification</i>
Collected For:	<u>PBM-05</u> ,
Definition:	Documentation that two unique patient identifiers were checked during a two-person verification process (or the use of automated identification technology may be used in place of one of the individuals) prior to the administration of the transfusion unit/dose (bag).
Suggested Data Collection Question:	Was there documentation that two unique patient identifiers were checked or automated identification was used in place of one person during the verification process prior to the administration of the blood transfusion unit/dose (bag)?
Format:	Length: 1 Type: Numeric Occurs: 1 - 12
Allowable Values:	<ol style="list-style-type: none"> 1 There was documentation that two unique patient identifiers were checked during the two person verification process or an automated identification system was used in place of one of the individuals prior to the administration of the transfusion unit/dose (bag). 2 There was no documentation that two unique patient identifiers or automated identification were used during the two-person identification check prior to the administration of the transfusion unit/dose (bag) or unable to determine from medical record documentation.
Notes for Abstraction:	<ul style="list-style-type: none"> • <i>Patient ID Verification</i> must be associated with the blood product and RBC ID that was selected for abstraction. • <i>Patient ID Verification</i> can be documented by the signature of two persons that attest that two unique patient identifiers were checked to verify the identification of the patient prior to the transfusion or the signature of one person and an automated identification device. • Patient identifiers that could be used include; name, date of birth, patient identification number or unique identifier given at the time the crossmatch was drawn. • The patient room number should not be used to identify the patient.
Suggested Data Sources:	<ul style="list-style-type: none"> • Anesthesia record • Emergency department record • Nursing notes • Progress notes • Physician's notes • Operative notes • Operative report • Procedure notes • PACU/recovery room record

- Blood administration form

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Plasma ID*

Collected For: PBM-03, PBM-05,

Definition: The number assigned to designate whether the plasma unit was the first, second or third unit transfused after hospital arrival.

Suggested Data Collection Question: What number was assigned to the plasma unit selected for abstraction?

Format:
Length: 1
Type: Numeric
Occurs: 1 - 3

Allowable Values:

- 1 First Plasma Unit
- 2 Second Plasma Unit
- 3 Third Plasma Unit

Notes for Abstraction:

- The abstractor assigns a plasma identification (ID) number for each unit evaluated.
- Each allowable value is only used one time and is determined by the order in which it was administered.
- Abstract up to three plasma transfusion units per patient.
- Include plasma transfusions administered after hospital arrival.

Suggested Data Sources:

- Anesthesia record
- Emergency department record
- Progress notes
- Operative notes
- Blood administration form
- Blood bank records

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Platelet ID*

Collected For: PBM-04, PBM-05,

Definition: The number assigned to designate whether the platelet unit was the first, second or third unit that was transfused after hospital arrival.

Suggested Data Collection Question: What number was assigned to the platelet unit selected for abstraction?

Format:
Length: 2
Type: Numeric
Occurs: 1 - 3

Allowable Values:

- 1 First Platelet Unit
- 2 Second Platelet Unit
- 3 Third Platelet Unit

Notes for Abstraction:

- The abstractor assigns a platelet identification (ID) number for each unit evaluated.
- Each allowable value is only used one time and is determined by the order in which it was administered.
- Abstract up to three platelet units per patient
- Include platelet transfusions administered after hospital arrival.

Suggested Data Sources:

- Anesthesia record
- Emergency department record
- Progress notes
- Operative notes
- Blood administration form
- Blood bank records

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Pre-transfusion Hematocrit*

Collected For: PBM-02,

Definition: Documentation of the closest hematocrit (hct) completed prior to the RBC transfusion.

Suggested Data Collection Question: What was documented as the closest pre-transfusion hct prior to the RBC transfusion?

Format:
Length: 4
Type: Alphanumeric
Occurs: 1 - 6

Allowable Values:

Enter the patient's closest hematocrit result (number only, reported in percent) performed prior to each RBC transfusion.

UTD = Unable to Determine

- For abstraction, select either the pre-transfusion hematocrit or the hemoglobin result; both are not required.
- Select the result associated with the RBC ID selected for abstraction.
- When recording the allowable value for hematocrit, input 23.00 if the patient's hematocrit is 23%.

Notes for Abstraction:

Suggested Data Sources:

- Consultation notes
- Emergency department record
- History and physical
- Laboratory report
- Progress notes
- Operative report
- Blood administration form

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Pre-transfusion Hemoglobin*

Collected For: PBM-02,

Definition: Documentation of the closest hemoglobin (hgb) completed prior to the RBC transfusion.

Suggested Data Collection Question: What was documented as the closest pre-transfusion hgb prior to the RBC transfusion?

Format:
Length: 4
Type: Alphanumeric
Occurs: 1 - 6

Allowable Values:

Enter the patient's closest hemoglobin result reported in g/dL performed prior to transfusion.

UTD = Unable to Determine

- For abstraction, select either the pre-transfusion hematocrit or the hemoglobin result; both are not required.
- Select the hemoglobin result that is associated with the RBC ID selected for abstraction.
- If the hemoglobin result is 9.9 g/dL, enter 9.9.

Notes for Abstraction:

Suggested Data Sources:

- Consultation notes
- Emergency department record
- History and physical
- Laboratory report
- Progress notes
- Operative report
- Blood administration form

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Pre-transfusion PT/INR Result*

Collected For: PBM-03,

Definition: Documentation of PT/INR result completed prior to the plasma transfusion.

Suggested Data Collection Question: What was the PT/INR result completed prior to the plasma transfusion.

Format:
Length: 1 - 5
Type: Alphanumeric
Occurs: 1 - 3

Allowable Values:
Enter the closest PT/INR result to the plasma transfusion.

UTD = Unable to determine

Notes for Abstraction:

- Enter the PT/INR result that is associated with the plasma ID selected for abstraction.
- An allowable value should be entered with one decimal. For example, a PT/INR of 1.5 should be entered as written. INR values over 10 should be entered as 10.00.

Suggested Data Sources:

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Pre-transfusion Platelet Count*

Collected For: PBM-04,

Definition: Documentation of the closest platelet count completed prior to the platelet transfusion.

Suggested Data Collection Question: What was the closest platelet count documented prior to the platelet transfusion?

Format:
Length: 1 - 5
Type: Alphanumeric
Occurs: 1 - 3

Allowable Values:

Enter the patient's closest platelet count result, in $10^9/\mu\text{L}$ performed prior to the platelet transfusion selected for abstraction.

UTD = Unable to Determine

Note:

- Select the platelet count result that is associated with the Platelet ID selected for abstraction.
- An allowable value for a platelet count result should be entered as '11.00' for a platelet count of 11,000.

Notes for Abstraction:

Suggested Data Sources:

- Anesthesia record
- Consultation notes
- Emergency department record
- History and physical
- Laboratory report
- Progress notes
- Operative report
- Blood administration form

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Preoperative Anemia Screening Date*

Collected For: PBM-06,

Definition: The date that preoperative anemia screening or a hemoglobin (hgb) or hematocrit (hct) result was completed.

Suggested Data Collection Question: What date was preoperative anemia screening or a hgb or hct result completed?

Format: **Length:** 10 - MM-DD-YYYY (includes dashes)
Type: Date
Occurs: 1

Allowable Values:

MM-DD-YYYY

MM = Month (01-12)
DD = Day (01-31)
YYYY = Year (2001-Current Year)
UTD

Notes for Abstraction:

- Select the *Preoperative Anemia Screening Date* associated with the elective surgical procedure selected for abstraction. *Preoperative Transfusion Testing*.
- The medical record must be abstracted as documented (taken at "face value"). When the date documented is obviously in error (not a valid date/format) and no other documentation is found that provides this information, the abstractor should select UTD.
- Example: Documentation indicates the Preoperative Anemia Screening Date was 03-42-2008. No other documentation in the medical record provides a valid date. Since the Preoperative Anemia Screening Date is outside of the range listed in the Allowable Values for "Day," it is not a valid date, and the abstractor should select UTD.

Suggested Data Sources:

- Nursing notes
- Progress notes
- Preop checklist
- Pre-arrival laboratory reports

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Preoperative Blood Type Testing*

Collected For: PBM-07,

Definition: Documentation that a type and screen or type and crossmatch was completed prior to anesthesia start time.

Suggested Data Collection Question: Was there documentation of a type and screen or type and crossmatch completed prior to anesthesia start time?

Format:
Length: 1
Type: Numeric
Occurs: 1

Allowable Values:

- 1 There is documentation that a type and screen or type and crossmatch was completed prior to anesthesia start time.
- 2 There is no documentation that a type and screen or type and crossmatch was completed prior to anesthesia start time or unable to determine from medical record documentation.

Notes for Abstraction:

- If type and screen and type and crossmatch were completed prior to the surgical procedure, select “1”.
- Anesthesia Start Time is the same as surgery start time.

Suggested Data Sources:

- Consultation notes
- History and physical
- Progress notes
- Preop checklist
- Pre-arrival laboratory reports

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *RBC ID*

Collected For: PBM-02, PBM-05,

Definition: The number assigned to designate whether the RBC transfusion was the first through the sixth RBC transfusion unit that was transfused after hospital arrival.

Suggested Data Collection Question: What RBC unit was selected for abstraction?

Format:
Length: 1
Type: Numeric
Occurs: 1 - 6

Allowable Values:

- 1 First RBC Unit
- 2 Second RBC Unit
- 3 Third RBC Unit
- 4 Fourth RBC Unit
- 5 Fifth RBC Unit
- 6 Sixth RBC Unit

Notes for Abstraction:

- The abstractor assigns a RBC identification (ID) number for each unit evaluated.
- Each allowable value is used only one time and is determined by the order in which it was administered.
- Abstract up to six RBC transfusion units per patient.
- Include RBC transfusions administered after hospital arrival.

Suggested Data Sources:

- Anesthesia record
- Emergency department record
- Progress notes
- Operative notes
- Operative report
- Medication administration record (MAR)
- Blood administration form
- Blood bank records

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
-----------	-----------

None	None
------	------

Data Element Name:	<i>RBC Unit Exclusions</i>
Collected For:	<u>PBM-02</u> , <u>PBM-05</u> ,
Definition:	Red blood cell (RBC) units that are excluded from abstraction. The following RBC units excluded from abstraction are; units used for a massive transfusion protocol or documentation of hemorrhagic shock, uncrossmatched units given during an emergency situation and units used to prime equipment for treatment.
Suggested Data Collection Question:	Was this unit transfused for a massive transfusion protocol, hemorrhagic shock, uncrossmatched or used to prime equipment?
Format:	Length: 1
	Type: Alphanumeric
	Occurs: 1-6
Allowable Values:	<ol style="list-style-type: none"> 1. There was documentation that this unit was transfused for a massive transfusion protocol, hemorrhagic shock, uncrossmatched or used to prime equipment 1. There was no documentation that this unit was transfused for a massive transfusion protocol, hemorrhagic shock, uncrossmatched or used to prime equipment or unable to determine from medical record documentation.
Notes for Abstraction:	<ul style="list-style-type: none"> • If the initial six units transfused are excluded due to the exclusion criteria, abstract the next six units that were transfused. If the patient only received RBC units that are excluded, then no RBC units should be abstracted.
Suggested Data Sources:	<ul style="list-style-type: none"> • Anesthesia record • Circulation record • Emergency department record • Laboratory report • Nursing notes • Nursing flow sheet • Progress notes • Physician orders • Physician's notes • Operative notes • Operating room notes • Operative report • Procedure notes • ICU notes

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Surgery Scheduled Timeframe*

Collected For: PBM-06,

Definition: The elective surgery was scheduled in less than 14 days from the planned surgery start date.

Suggested Data Collection Question: Was the elective surgery scheduled in less than 14 days from the planned surgery?

Format:
Length: 1
Type: Alphanumeric
Occurs: 1

Allowable Values:

- 1 There was documentation that the elective surgery was scheduled in less than 14 days from the planned surgery.
- 2 There was no documentation that the elective surgery was scheduled in less than 14 days from the planned surgery or unable to determine from medical record documentation.

Notes for Abstraction:

Suggested Data Sources:

- Preop checklist

Preoperative paperwork

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Transfusion Consent*

Collected For: PBM-01,

Definition: Documentation of a signed consent **prior** to the first transfusion of RBCs, platelets or plasma.

Suggested Data Collection Question: Was there documentation of a signed consent **prior** to the first blood transfusion?

Format:
Length: 1
Type: Numeric
Occurs: 1

Allowable Values:

- 1 There was documentation of a signed consent prior to the first blood transfusion.
- 2 The first blood transfusion was deemed a medical emergency.
- 3 There was no documentation of a blood transfusion consent prior to the first blood transfusion or unable to determine from medical record documentation.

Notes for Abstraction:

- The consent may be signed by the patient or caregiver.
- If organizations require a consent prior to every transfusion, then review the record for the first transfusion to answer this data element.
- For hospitals that use a general consent for treatment that includes transfusions, select “Yes”.
- If a patient receives chronic transfusions and a previous consent is acceptable for a defined timeframe within the institution, select “1” if the consent is valid.

Suggested Data Sources:

- Emergency department record
- History and physical
- Nursing notes
- Progress notes
- Operative notes
- Consent form

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Transfusion Order*

Collected For: PBM-05,

Definition: An order to transfuse was written by the physician/advance practice nurse/physician assistant (physician/APN/PA) **prior** to the initiation of the transfusion.

Suggested Data Collection Question: Was there documentation of an order to transfuse **prior** to the transfusion?

Format:
Length: 1
Type: Numeric
Occurs: 1 - 12

Allowable Values:

- 1 There was documentation of an order to transfuse prior to transfusion.
- 2 There was no documentation of an order to transfuse prior to transfusion or unable to determine from medical record documentation.

Notes for Abstraction:

- A verbal or telephone order that was written prior to the transfusion is acceptable.
- The Transfusion Order must be associated with the blood product unit ID that was selected for abstraction.
- Note: Transfusion Order may apply to more than one unit/dose (bag). For example: An order written to "Transfuse two doses of platelets" would apply to both bags that were administered.

Suggested Data Sources:

ONLY PHYSICIAN/APN/PA DOCUMENTATION OF THE ORDER TO TRANSFUSE:

- Anesthesia record
- Consultation notes
- Emergency department record
- Operative notes
- Physician orders
- Progress notes

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name:	<i>Transfusion Start Date</i>
Collected For:	<u>PBM-05</u> ,
Definition:	The date that the blood transfusion unit/dose (bag) was administered.
Suggested Data Collection Question:	What is the date that the blood transfusion unit/dose (bag) was administered?
Format:	Length: 10 – MM-DD-YYYY (includes dashes)
	Type: Date
	Occurs: 1 - 12
Allowable Values:	MM-DD-YYYY MM = Month (01-12) DD = Day (01-31) YYYY = Year (2001-Current Year) UTD
Notes for Abstraction:	<ul style="list-style-type: none"> • Abstract the Transfusion Date associated with the Transfusion Start Time of the unit/dose (bag) from the blood product ID selected for abstraction. • Some of the dates of the transfusion units may be the same date. Record a transfusion date for each unit abstracted up to three units for plasma or platelets or up to six units for RBCs. • The medical record must be abstracted as documented (taken at “face value”). When the date documented is obviously in error (not a valid date/format) and no other documentation is found that provides this information, the abstractor should select UTD. Example: Documentation indicates the Transfusion Start Date was 03-42-2008. No other documentation in the medical record provides a valid date. Since the Transfusion Start Date is outside of the range listed in the Allowable Values for “Day,” it is not a valid date and the abstractor should select UTD.
Suggested Data Sources:	<ul style="list-style-type: none"> • Anesthesia record • Emergency department record • Nursing notes • Progress notes • Operative notes • Blood administration record
Additional Notes:	

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name:	<i>Transfusion Start Time</i>
Collected For:	<u>PBM-05</u> ,
Definition:	The start time (military time) of the unit/dose (bag) of RBCs, plasma or platelets that was administered.
Suggested Data Collection Question:	What was the start time of the blood unit/dose (bag) administration?
Format:	Length: 5 - HH:MM (with or without colon) or UTD Type: Time Occurs: 1 - 12
Allowable Values:	<p>Select the Transfusion Start Time associated with the Transfusion Start Date of the unit/dose (bag) from the associated blood product ID being abstracted.</p> <p>HH = Hour (00-23) MM = Minutes (00-59) UTD = Unable to Determine</p>
Notes for Abstraction:	<p>Time must be recorded in military time format. With the exception of Midnight and Noon:</p> <ul style="list-style-type: none"> • If the time is in the a.m., conversion is not required • If the time is in the p.m., add 12 to the clock time hour <p>Examples: Midnight - 00:00 Noon - 12:00 5:31 am - 05:31 5:31pm - 17:31 11:59 am - 11:59 11:59pm - 23:59</p> <ul style="list-style-type: none"> • For times that include “seconds,” remove the seconds and record the time as is. Example: 15:00:35 would be recorded as 15:00 • If more than one Transfusion Start Time is documented, use the earliest time documented. • The medical record must be abstracted as documented (taken at “face value”). When the time documented is obviously in error (not a valid format/range) and no other documentation is found that provides this information, the abstractor should select “UTD.” • Example: Documentation indicates the Transfusion Start Time was 3300. Since the Transfusion Start Time is outside of the range in the Allowable Values for “Hour,” it is not a valid time and the abstractor should select “UTD.”
Suggested Data Sources:	<ul style="list-style-type: none"> • Anesthesia record

- Emergency department record
- Nursing notes
- Operative notes
- Operative report
- Blood administration form

Additional Notes:

Select the Transfusion Start Time associated with the Transfusion Start Date of the unit/dose (bag) from the blood product ID identified for abstraction.

Time must be recorded in military time format.
With the exception of Midnight and Noon:

- If the time is in the a.m., conversion is not required
- If the time is in the p.m., add 12 to the clock time hour.

The medical record must be abstracted as documented (taken at “face value”). When the time documented is obviously in error (not a valid format/range) and no other documentation is found that provides this information, the abstractor should select “UTD.”

Example:

Documentation indicates the Transfusion Start Time was 3300. Since the Transfusion Start Time is outside of the range in the Allowable Values for “Hour,” it is not a valid time and the abstractor should select “UTD.”

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Data Element Name: *Vital Sign Monitoring*

Collected For: PBM-05,

Definition: Documentation of blood pressure (BP), pulse and temperature monitored at specific intervals for the transfusion. The intervals are:

- Pre-transfusion, within 15 minutes of the initiation of the transfusion and within one hour of transfusion completion

Suggested Data Collection Question: Was there documentation of BP and temperature monitored for all of the specified intervals for the transfusion?

Format: **Length:** 2
Type: Numeric
Occurs: 1 -12

Allowable Values:

- 1 There was documentation for all of the BP, pulse and temperature monitoring intervals for the transfusion.
- 2 There was no documentation for all of the blood pressure, pulse and temperature monitoring intervals for the transfusion or unable to determine from medical record documentation.

Notes for Abstraction:

- All vital signs must be recorded at the following times: pre-transfusion, within 15 minutes of the initiation of the transfusion and within one hour of transfusion completion. To select "1", all recordings must be documented.
- The pre-transfusion BP, pulse and temperature must be within one hour of the Transfusion Start Time. Vitals documented at the start of the transfusion are considered "within one hour of transfusion initiation".
- For blood that may be transfused within 15 minutes, select "1" if the pre-transfusion and the within one hour of transfusion completion vitals are documented.
- Vitals documented at the completion of the transfusion are considered "within one hour of transfusion completion".
- The "unit" or "dose" information for the Vital Sign Monitoring data element must be associated with the blood product ID that was selected for abstraction.

Suggested Data Sources:

- Anesthesia record
- Consultation notes
- Emergency department record
- Nursing notes
- Progress notes
- Operative notes

Additional Notes:

Guidelines for Abstraction:

Inclusion	Exclusion
None	None

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

Index		
Number	Name	Page
Table 2.2	Left Ventricular Assistive Device (LVAD) and Heart Transplant	
Table 5.01	Coronary Artery Bypass Graft (CABG)	
Table 5.02	Other Cardiac Surgery	
Table 5.08	Vascular Surgery	
Table 5.11	Cardiac Surgery	
Table 5.22	Elective Hip Replacement	
Table 5.23	Elective Total Knee Replacement	
Table 9.1	Elective Cardiac Surgery	
Table 9.2	Elective Hysterectomy	
Table 9.3	Previously Donated Autologous Transfusion	
Table 9.4	Packed Red Blood Cell Transfusion	
Table 9.5	Platelet Transfusion	
Table 9.6	Plasma (Serum) Transfusion	
Table 9.7	Trauma	

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

Table 2.2 Left Ventricular Assistive Device (LVAD) and Heart Transplant		
Code	ICD-9-CM Description	Shortened Description
33.6	Combined heart-lung transplantation	COMB HEART/LUNG TRANSPLA
37.51	Heart transplantation	HEART TRANSPLANTATION
37.52	Implantation of total replacement heart system	IMPLANT TOT REP HRT SYS
37.53	Replacement or repair of thoracic unit of total replacement heart system	REPL/REP THORAC UNIT HRT
37.54	Replacement or repair of other implantable component of total replacement heart system	REPL/REP OTH TOT HRT SYS
37.62	Insertion of non-implantable heart assist system	INS NON-IMPL HRT ASSIST
37.63	Repair of heart assist system	REPAIR HEART ASSIST SYS
37.64	Removal of heart assist system	REMOVE HEART ASSIST SYS
37.65	Implant of external heart assist system	IMP EXT HRT ASSIST SYST
37.66	Insertion of implantable heart assist system	IMPLANTABLE HRT ASSIST
37.68	Insertion of percutaneous external heart assist device	PERCUTAN HRT ASSIST SYST

Table 5.01 Coronary Artery Bypass Graft (CABG)		
Code	ICD-9-CM Description	Shortened Description
36.10	Aortocoronary bypass for heart revascularization, not otherwise specified	AORTOCORONARY BYPASS NOS
36.11	(Aorto)coronary bypass of one coronary artery	(AORTO)COR BYPAS-1 COR ART
36.12	(Aorto)coronary bypass of two coronary arteries	(AORTO)COR BYPAS-2 COR ART
36.13	(Aorto)coronary bypass of three coronary arteries	(AORTO)COR BYPAS-3 COR ART
36.14	(Aorto)coronary bypass of four coronary arteries	(AORT)COR BYPAS-4+ COR ART
36.15	Single internal mammary-coronary artery bypass	1 INT MAM-COR ART BYPASS
36.16	Double internal mammary-coronary artery bypass	2 INT MAM-COR ART BYPASS
36.17	Abdominal-coronary artery bypass	ABD-CORON ARTERY BYPASS
36.19	Other bypass anastomosis for heart revascularization	HRT REVAS BYPS ANAS NEC

Table 5.02 Other Cardiac Surgery		
Code	ICD-9-CM Description	Shortened Description
35.10	Open heart valvuloplasty, without replacement, unspecified valve	OPEN VALVULOPLASTY NOS
35.11	Open heart valvuloplasty of aortic valve without replacement	OPN AORTIC VALVULOPLASTY
35.12	Open heart valvuloplasty of mitral valve without replacement	OPN MITRAL VALVULOPLASTY
35.13	Open heart valvuloplasty of pulmonary valve without replacement	OPN PULMON VALVULOPLASTY
35.14	Open heart valvuloplasty of tricuspid valve without	OPN TRICUS

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

	replacement	VALVULOPLASTY
35.20	Replacement of unspecified heart valve	REPLACE HEART VALVE NOS
35.21	Replacement of aortic valve with tissue graft	REPLACE AORT VALV-TISSUE
35.22	Other replacement of aortic valve	REPLACE AORTIC VALVE NEC
35.23	Replacement of mitral valve with tissue graft	REPLACE MITR VALV-TISSUE
35.24	Other replacement of mitral valve	REPLACE MITRAL VALVE NEC
35.25	Replacement of pulmonary valve with tissue graft	REPLACE PULM VALV-TISSUE
35.26	Other replacement of pulmonary valve	REPLACE PULMON VALVE NEC
35.27	Replacement of tricuspid valve with tissue graft	REPLACE TRIC VALV-TISSUE
35.28	Other replacement of tricuspid valve	REPLACE TRICUSP VALV NEC
35.31	Operations on papillary muscle	PAPILLARY MUSCLE OPS
35.32	Operations on chordae tendineae	CHORDAE TENDINEAE OPS
35.33	Annuloplasty	ANNULOPLASTY
35.34	Infundibulectomy	INFUNDIBULECTOMY
35.35	Operations on trabeculae carneae cordis	TRABECUL CARNEAE CORD OP
35.39	Operations on other structures adjacent to valves of heart	TISS ADJ TO VALV OPS NEC
35.42	Creation of septal defect in heart	CREATE SEPTAL DEFECT
35.50	Repair of unspecified septal defect of heart with prosthesis	PROSTH REP HRT SEPTA NOS
35.51	Repair of atrial septal defect with prosthesis, open technique	PROS REP ATRIAL DEF-OPN
35.53	Repair of ventricular septal defect with prosthesis, open technique	PROS REP VENTRIC DEF- OPN
35.54	Repair of endocardial defect with prosthesis	PROS REP ENDOCAR CUSHION
35.60	Repair of unspecified septal defect with tissue graft	GRFT REPAIR HRT SEPT NOS
35.61	Repair of atrial septal defect with tissue graft	GRAFT REPAIR ATRIAL DEF
35.62	Repair of ventricular septal defect with tissue graft	GRAFT REPAIR VENTRIC DEF
35.63	Repair of endocardial cushion defect with tissue graft	GRFT REP ENDOCAR CUSHION
35.70	Other and unspecified repair of unspecified septal defect of heart	HEART SEPTA REPAIR NOS
35.72	Other and unspecified repair of ventricular septal defect	VENTR SEPTA DEF REP NEC
35.73	Other and unspecified repair of endocardial cushion defect	ENDOCAR CUSHION REP NEC
35.81	Total repair of tetralogy of Fallot	TOT REPAIR TETRAL FALLOT
35.82	Total repair of total anomalous pulmonary venous connection	TOTAL REPAIR OF TAPVC
35.83	Total repair of truncus arteriosus	TOT REP TRUNCUS ARTERIOS
35.84	Total correction of transposition of great vessels, not elsewhere classified	TOT COR TRANSPOS GRT VES
35.91	Interatrial transposition of venous return	INTERAT VEN RETRN TRANSP

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

35.92	Creation of conduit between right ventricle and pulmonary artery	CONDUIT RT VENT-PUL ART
35.93	Creation of conduit between left ventricle and aorta	CONDUIT LEFT VENTR-AORTA
35.94	Creation of conduit between atrium and pulmonary artery	CONDUIT ARTIUM-PULM ART
35.98	Other operations on septa of heart	OTHER HEART SEPTA OPS
35.99	Other operations on valves of heart	OTHER HEART VALVE OPS

Table 5.08 Vascular Surgery		
Code	ICD-9-CM Description	Shortened Description
38.14	Endarterectomy, aorta	ENDARTERECTOMY OF AORTA
38.16	Endarterectomy, abdominal arteries	ABDOMINAL ENDARTERECTOMY
38.18	Endarterectomy, lower limb arteries	LOWER LIMB ENDARTERECT
38.34	Resection of vessel with anastomosis, aorta	AORTA RESECTION & ANAST
38.36	Resection of vessel with anastomosis, abdominal arteries	ABD VESSEL RESECT/ANAST
38.37	Resection of vessel with anastomosis, abdominal veins	ABD VEIN RESECT & ANAST
38.44	Resection of vessel with replacement, aorta, abdominal	RESECT ABDOM
38.48	Resection of vessel with replacement, lower limb arteries	LEG ARTERY RESEC W REPLA
38.49	Resection of vessel with replacement, lower limb veins	LEG VEIN RESECT W REPLAC
38.64	Other excision of vessels, aorta, abdominal	EXCISION OF AORTA
39.25	Aorta-iliac-femoral bypass	AORTA-ILIAC-FEMOR BYPASS
39.26	Other intra-abdominal vascular shunt or bypass	INTRA-ABDOMIN SHUNT NEC
39.29	Other (peripheral) vascular shunt or bypass	VASC SHUNT & BYPASS NEC

Table 5.11 Cardiac Surgery		
Code	ICD-9-CM Description	Shortened Description
35.10	Open heart valvuloplasty without replacement, unspecified valve	OPEN VALVULOPLASTY NOS
35.11	Open heart valvuloplasty of aortic valve without replacement	OPN AORTIC VALVULOPLASTY
35.12	Open heart valvuloplasty of mitral valve without replacement	OPNMITRAL VALVULOPLASTY
35.13	Open heart valvuloplasty of pulmonary valve without replacement	OPN PULMON VALVULOPLASTY
35.14	Open heart valvuloplasty of tricuspid valve without replacement	OPN TRICUS VALVULOPLASTY
35.20	Replacement of unspecified heart valve	REPLACE HEART VALVE NOS
35.21	Replacement of aortic valve with tissue graft	REPLACE AORT VALVE-TISSUE
35.22	Other replacement of aortic valve	REPLACE AORT VALVE NEC

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

35.23	Replacement of mitral valve with tissue graft	REPLACE MITR VALVE-TISSUE
35.24	Other replacement of mitral valve	REPLACE MITRAL VALVE NEC
35.25	Replacement of pulmonary valve with tissue graft	REPLACE PULM VALV-TISSUE
35.26	Other replacement of pulmonary valve	REPLACE PULMON VALVE NEC
35.27	Replacement of tricuspid valve with tissue graft	REPLACE TRICUSP VALV NEC
35.28	Other replacement of tricuspid valve	REPLACE TRICUSP VALV NEC
35.31	Operations on papillary muscle	PAPILLARY MUSCLE OPS
35.32	Operations on chordae tendineae	CHORDAE TENDINEAE OPS
35.33	Annuloplasty	ANNULOPLASTY
35.34	Infundibulectomy	INFUNDIBULECTOMY
35.35	Operations of trabeculae carneae cordis	TRABECUL CARNEAE CORD OP
35.39	Operations on other structures adjacent to valves of heart	TISS ADJ TO VALV OPS NEC
35.42	Creation of septal defect in heart	CREATE SEPTAL DEFECT
35.50	Repair of unspecified septal defect of heart with prosthesis	PROSTH REP HRT SEPTA NOS
35.51	Repair of atrial septal defect with prosthesis, open technique	PROS REP ATRIAL DEF-OPN
35.53	Repair of ventricular septal defect with prosthesis, open technique	PROS REP VENTRIC DEF-OPN
35.54	Repair of endocardial cushion defect with prosthesis	PROS REP ENDOCAR CUSHION
35.60	Repair of unspecified septal defect of heart with tissue graft	GRFT REPAIR HRT SEPT NOS
35.61	Repair of atrial septal defect with tissue graft	GRAFT REPAIR ATRIAL DEF
35.62	Repair of ventricular septal defect with tissue graft	GRAFT REPAIR VENTRIC DEF
35.63	Repair of endocardial cushion defect with tissue graft	GRFT REP ENDOCAR CUSHION
35.70	Other and unspecified repair of unspecified septal defect of heart	HEART SEPTA REPAIR NOS
35.71	Other and unspecified repair of atrial septal defect	ATRIA SEPTA DEF REP NEC
35.72	Other and unspecified repair of ventricular septal defect	VENTR SEPTA DEF REP NEC
35.73	Other and unspecified repair of endocardial cushion defect	ENDOCAR CUSHION REP NEC
35.81	Total repair of tetralogy of Fallot	TOT REPAIR TETRAL FALLOT
35.82	Total repair of total anomalous pulmonary venous connection	TOTAL REPAIR OF TAPVC
35.83	Total repair of truncus arteriosus	TOT REP TRUNCUS ARTERIOS

Table 5.11 Cardiac Surgery (cont.)		
Code	ICD-9-CM Description	Shortened Description

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

35.84	Total connection of transposition of great vessels, not elsewhere classified	TOT COR TRANSPOS GRT VES
35.91	Interatrial transposition of venous return	INTERAT VEN RETRN TRANSP
35.92	Creation of conduit between right ventricle and pulmonary artery	CONDUIT RT VENT-PUL ART
35.93	Creation of conduit between left ventricle and aorta	CONDUIT LEFT VENTR-AORTA
35.94	Creation of conduit between atrium and pulmonary artery	CONDUIT ARTIUM-PULM ART
35.98	Other operations on septa of heart	OTHER HEART SEPTA OPS
35.99	Other operations on valves of heart	OTHER HEART VALVE OPS
36.03	Open chest coronary artery angioplasty	OPEN CORONRY ANGIOPLASTY
36.10	Aortocoronary bypass for heart revascularization, not otherwise specified	AORTOCORONARY BYPASS NOS
36.11	Aortocoronary bypass of one coronary artery	AORTOCOR BYPASS-1 COR ART
36.12	Aortocoronary bypass of two coronary arteries	AORTOCOR BYPASS-2 COR ART
36.13	Aortocoronary bypass of three coronary arteries	AORTOCOR BYPASS-3 COR ART
36.14	Aortocoronary bypass of four or more coronary arteries	AORTOCOR BYPASS-4+ COR ART
36.15	Single internal mammary-coronary artery bypass	1 INT MAM-COR ART BYPASS
36.16	Double internal mammary-coronary artery bypass	2 INT MAM-COR ART BYPASS
36.17	Abdominal-coronary artery bypass	ABD-CORON ARTERY BYPASS
36.19	Other bypass anastomosis for heart revascularization	HRT REVAS BYPS ANAS NEC
36.31	Open chest transmyocardial revascularization	OPEN CHEST TRANS REVASC
36.32	Other transmyocardial revascularization	OTH TRANSMYO REVASCULAR
36.39	Other heart revascularization	OTH REVASCULAR
36.91	Repair of aneurysm of coronary vessel	CORON VESS ANEURYSM REP
36.99	Other operations on vessels of heart	HEART VESSEL OP NEC
37.10	Incision of heart, not otherwise specified	INCISION OF HEART NOS
37.11	Cardiotomy	CARDIOTOMY
37.31	Pericardiectomy	PERICARDIECTOMY
37.32	Excision of aneurysm of heart	HEART ANEURYSM EXCISION
37.33	Excision or destruction of other lesion or tissue of heart, open approach	EXC/DEST HRT LESION OPEN
37.35	Partial ventriculectomy	PARTIAL VENTRICULECTOMY
37.41	Implantation of prosthetic cardiac support device around the heart	IMPL CARDIAC SUPPORT DEV
37.49	Other repair of heart and pericardium	HEART/PERICARD REPR NEC
37.51	Heart transplantation	HEART TRANSPLANTATION

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

37.52	Implantation of total replacement heart system	IMPLANT TOT REP HRT SYS
37.53	Replacement or repair of thoracic unit of total replacement heart system	REPL/REP THORAC UNIT HRT
37.54	Replacement or repair of other implants component of total replacement heart system	REPL/REP OTH TOT HRT SYS
37.62	Insertion of non-implantable heart assist system	INS NON-IMPL HRT ASSIST
37.63	Repair of heart assist system	REPAIR HEART ASSIST SYS
37.64	Removal of heart assist system	REMOVE HEART ASSIST SYS
37.66	Insertion of implantable heart assist system	IMPLANTABLE HRT ASSIST
37.67	Implantation of cardiomyostimulation system	IMP CARDIOMYOSTIMUL SYS

Table 5.22 Elective Hip Replacement		
Code	ICD-9-CM Description	Shortened Description
00.70	Revision of hip replacement, both acetabular and femoral components	REV HIP REPL-ACETAB/FEM
00.71	Revision of hip replacement, acetabular component	REV HIP REPL-ACETAB COMP
00.72	Revision of hip replacement, femoral component	REV HIP REPL-FEM COMP
00.73	Revision of hip replacement, acetabular liner and/or femoral head only	REV HIP REPL-LINER/HEAD
00.77	Hip bearing surface, ceramic-on-polyethylene	HIP SURFACE, CERMC/POLY
00.85	Resurfacing hip, total, acetabulum and femoral head	RESRF HIP,TOTAL-ACET/FEM
00.86	Resurfacing hip, partial, femoral head	RESRF HIP,PART-FEM HEAD
00.87	Resurfacing hip, partial, acetabulum	RESRF HIP,PART-ACETABLUM
81.51	Total hip replacement	TOTAL HIP REPLACEMENT
81.52	Partial hip replacement	PARTIAL HIP REPLACEMENT
81.53	Revision of hip replacement	REVISE HIP REPLACEMENT

Table 5.23 Elective Total Knee Replacement		
Code	ICD-9-CM Description	Shortened Description
00.80	Revision of knee replacement, total (all components)	REV KNEE REPLACENT-TOTAL
00.81	Revision of knee replacement, tibial component	REV KNEE REPL-TIBIA COMP
00.82	Revision of knee replacement, femoral component	REV KNEE REPL-FEMUR COMP
00.83	Revision of knee replacement, patellar component	REV KNEE REPLACE-PATELLA
00.84	Revision of total knee replacement, tibial insert (liner)	REV KNEE REPL-TIBIA LIN
81.54	Total knee replacement	TOTAL KNEE REPLACEMENT
81.55	Revision of knee replacement	REVISE KNEE REPLACEMENT

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

Table 9.1 Elective Cardiac Surgery (Selected Codes from Table 5.25)		
Code	ICD-9-CM Description	Shortened Description
35.71	Other and unspecified repair of atrial septal defect	ATRIA SEPTA DEF REP NEC
36.03	Open chest coronary artery angioplasty	OPEN CORONRY ANGIOPLASTY
36.31	Open chest transmyocardial revascularization	OPEN CHEST TRANS REVASC
36.32	Other transmyocardial revascularization	OTH TRANSMYO REVASCULAR
36.39	Other heart revascularization	OTH HEART REVASCULAR
36.91	Repair of aneurysm of coronary vessel	CORON VESS ANEURYSM REP
36.99	Other operations on vessels of heart	HEART VESSEL OP NEC
37.10	Incision of heart, not otherwise specified	INCISION OF HEART NOS
37.11	Cardiotomy	CARDIOTOMY
37.32	Excision of aneurysm of heart	HEART ANEURYSM EXCISION
37.33	Excision or destruction of other lesion or tissue of heart, open approach	EXC/DEST HRT LESION OPEN
37.35	Partial ventriculectomy	PARTIAL VENTRICULECTOMY
37.36	Excision or destruction of left atrial appendage (LAA)	EXC LEFT ATRIAL APPENDAG
37.41	Implantation of prosthetic cardiac support device around the heart	IMPL CARDIAC SUPPORT DEV
37.49	Other repair of heart and pericardium	HEART/PERICARD REPR NEC
37.51	Heart transplantation	HEART TRANSPLANTATION
37.52	Implantation of total internal biventricular heart replacement system	IMP TOT INT BI HT RP SYS
37.53	Replacement or repair of thoracic unit of (total) replacement heart system	REPL/REP THR UNT TOT HRT
37.54	Replacement or repair of other implantable component of (total) replacement heart system	REPL/REP OTH TOT HRT SYS
37.55	Removal of internal biventricular heart replacement system	REM INT BIVENT HRT SYS
37.60	Implantation or insertion of biventricular external heart assist system	IMP BIVN EXT HRT AST SYS
37.62	Insertion of temporary non-implantable extracorporeal circulatory assist device	INSRT NON-IMPL CIRC DEV
37.63	Repair of heart assist system	REPAIR HEART ASSIST SYS
37.64	Removal of external heart assist system(s) or device(s)	REMVE EXT HRT ASSIST SYS
37.66	Insertion of implantable heart assist system	IMPLANTABLE HRT ASSIST
37.67	Implantation of cardiomyostimulation system	IMP CARDIOMYOSTIMUL SYS

Table 9.2 Elective Gynecological		
Code	ICD-9-CM Description	Shortened Description
68.31	Other incision and excision of uterus, subtotal abdominal hysterectomy, other incision and excision of uterus, laparoscopic supracervical hysterectomy [LSH]	Lap scervic hysterectomy
68.39	Other incision and excision of uterus, subtotal abdominal hysterectomy, other incision and excision of uterus, other and unspecified subtotal	Subtotl abd hyst NEC/NOS

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

	abdominal hysterectomy	
68.41	Other incision and excision of uterus, total abdominal hysterectomy, laparoscopic total abdominal hysterectomy	Lap total abdominal hyst
68.49	Other incision and excision of uterus, total abdominal hysterectomy, other and unspecified total abdominal hysterectomy	Total abd hyst NEC/NOS
68.51	Vaginal hysterectomy, laparoscopically assisted vaginal hysterectomy [LAVH]	Lap ast vag hysterectomy
68.59	Vaginal hysterectomy, other and unspecified vaginal hysterectomy	Vag hysterectomy NEC/NOS
68.61	Radical abdominal hysterectomy, laparoscopic radical abdominal hysterectomy	Lap radical abdomnl hyst
68.69	Radical abdominal hysterectomy, other and unspecified radical abdominal hysterectomy	Radical abd hyst NEC/NOS
68.71	Radical vaginal hysterectomy, laparoscopic radical vaginal hysterectomy [LRVH]	Lap radical vaginal hyst
68.79	Radical vaginal hysterectomy, other and unspecified radical vaginal hysterectomy	Radical vag hyst NEC/NOS
68.9	Other and unspecified hysterectomy	Hysterectomy NEC/NOS

Table 9.3 Previously Donated Autologous Transfusion		
Code	ICD-9-CM Description	Shortened Description
99.02	Other nonoperative procedures, transfusion of blood and blood components, transfusion of previously collected autologous blood	TRANSFUS PREV AUTO BLOOD

Table 9.4 Packed Red Blood Cell Transfusion		
Code	ICD-9-CM Description	Shortened Description
99.04	Other nonoperative procedures, transfusion of blood and blood components, transfusion of packed cells	PACKED CELL TRANSFUSION

Table 9.5 Platelet Transfusion		
Code	ICD-9-CM Description	Shortened Description
99.05	Other nonoperative procedures, transfusion of blood and blood components, transfusion of platelets	PLATELET TRANSFUSION

Table 9.6 Plasma Transfusion		
Code	ICD-9-CM Description	Shortened Description
99.07	Other nonoperative procedures, transfusion of blood and blood components, transfusion of other serum	SERUM TRANSFUSION NEC

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

Table 9.7 Trauma		
Code	ICD-9-CM Description	Shortened Description
800	Fracture of vault of skull	CLOSED SKULL VAULT FX
801	Fracture of base of skull	CLOS SKULL BASE FRACTURE
802	Fracture of face bones	NASAL BONE FX-CLOSED
803	Other and unqualified skull fractures	CLOSE SKULL FRACTURE NEC
804	Multiple fractures involving skull or face with other bones	CL SKUL FX W OTH BONE FX
805	Fracture of vertebral column without mention of spinal cord injury	FX CERVICAL VERT NOS-CL
806	Fracture of vertebral column with spinal cord injury	C1-C4 FX-CL/CORD INJ NOS
807	Fracture of rib(s), sternum, larynx, and trachea	FRACTURE RIB NOS-CLOSED
808	Fracture of pelvis	FRACTURE ACETABULUM-CLOS
809	Ill-defined fractures of bones of trunk	FRACTURE TRUNK BONE-CLOS
810	Fracture of clavicle	FX CLAVICLE NOS-CLOSED
811	Fracture of scapula	FX SCAPULA NOS-CLOSED
812	Fracture of humerus	FX UP END HUMERUS NOS-CL
813	Fracture of radius and ulna	FX UPPER FOREARM NOS-CL
814	Fracture of carpal bones(s)	FX CARPAL BONE NOS-CLOSE
815	Fracture of metacarpal bones(s)	FX METACARPAL NOS-CLOSED
816	Fracture of one or more phalanges of hands	FX PHALANX, HAND NOS-CL
817	Multiple fractures of hand bones	MULTIPLE FX HAND-CLOSED
818	Ill-defined fractures of upper limb	FX ARM MULT/NOS-CLOSED
819	Multiple fractures involving both upper limbs, and upper limb with rib(s) and sternum	FX ARMS W RIB/STERNUM-CL
820	Fracture of neck of femur	FX FEMUR INTRCAPS NOS-CL
821	Fracture of other and unspecified parts of femur	FX FEMUR NOS-CLOSED
822	Fracture of patella	FRACTURE PATELLA-CLOSED
823	Fracture of tibia and fibula	FX UPPER END TIBIA-CLOSE
824	Fracture of ankle	FX MEDIAL MALLEOLUS-CLOS
825	Fracture of one or more tarsal and metatarsal bones	FRACTURE CALCANEUS-CLOSE
826	Fracture of one or more phalanges of foot	FX PHALANX, FOOT-CLOSED
827	Other, multiple, and ill-defined fractures of lower limb	FX LOWER LIMB NEC-CLOSED
828	Multiple fractures involving both lower limbs, lower with upper limb, and lower limb(s) with rib(s) and sternum	FX LEGS W ARM/RIB-CLOSED
829	Fracture of unspecified bones	FRACTURE NOS-CLOSED
830	Dislocation of jaw	DISLOCATION JAW-CLOSED
831	Dislocation of shoulder	DISLOC SHOULDER NOS-

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

		CLOS
832	Dislocation of elbow	DISLOCAT ELBOW NOS-CLOSE
833	Dislocation of wrist	DISLOC WRIST NOS-CLOSED
834	Dislocation of finger	DISL FINGER NOS-CLOSED
835	Dislocation of hip	DISLOCAT HIP NOS-CLOSED
836	Dislocation of knee	TEAR MED MENISC KNEE-CUR
837	Dislocation of ankle	DISLOCATION ANKLE-CLOSED
838	Dislocation of foot	DISLOCAT FOOT NOS-CLOSED
839	Other, multiple, and ill-defined dislocations	DISLOC CERV VERT NOS-CL
840	Sprains and strains of shoulder and upper arm	SPRAIN ACROMIOCLAVICULAR
841	Sprains and strains of elbow and forearm	SPRAIN RADIAL COLLAT LIG
842	Sprains and strains of wrist and hand	SPRAIN OF WRIST NOS
843	Sprains and strains of hip and thigh	SPRAIN ILIOFEMORAL
844	Sprains and strains of knee and leg	SPRAIN LATERAL COLL LIG
845	Sprains and strains of ankle and foot	SPRAIN OF ANKLE NOS
846	Sprains and strains of sacroiliac region	SPRAIN LUMBOSACRAL
847	Sprains and strains of other and unspecified parts of back	SPRAIN OF NECK
848	Other and ill-defined sprains and strains	SPRAIN OF NASAL SEPTUM
850	Concussion	CONCUSSION W/O COMA
851	Cerebral laceration and contusion	CEREBRAL CORTX CONTUSION
852	Subarachnoid, subdural, and extradural hemorrhage, following injury	TRAUM SUBARACHNOID HEM
853	Other and unspecified intracranial hemorrhage following injury	TRAUMATIC BRAIN HEM NEC
854	Intracranial injury of other and unspecified nature	BRAIN INJURY NEC
860	Traumatic pneumothorax and hemothorax	TRAUM PNEUMOTHORAX-CLOSE
861	Injury to heart and lung	HEART INJURY NOS-CLOSED
862	Injury to other and unspecified intrathoracic organs	DIAPHRAGM INJURY-CLOSED
863	Injury to gastrointestinal tract	STOMACH INJURY-CLOSED
864	Injury to liver	LIVER INJURY NOS-CLOSED
865	Injury to spleen	SPLEEN INJURY NOS-CLOSED
866	Injury to kidney	KIDNEY INJURY NOS-CLOSED
867	Injury to pelvic organs	BLADDER/URETHRA INJ-CLOS
868	Injury to other intra-abdominal organs	INTRA-ABDOM INJ NOS-CLOS
869	Internal injury to unspecified or ill-defined organs	INTERNAL INJ NOS-CLOSED
870	Open wound of ocular adnexa	LAC EYELID SKN/PERIOCLUR
871	Open wound of eyeball	OCULAR LAC W/O PROLAPSE
872	Open wound of ear	OPN WOUND EXTERN EAR

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

		NOS
873	Other open wound of head	OPEN WOUND OF SCALP
874	Open wound of neck	OPN WND LARYNX W TRACHEA
875	Open wound of chest (wall)	OPEN WOUND OF CHEST
876	Open wound of back	OPEN WOUND OF BACK
877	Open wound of buttock	OPEN WOUND OF BUTTOCK
878	Open wound of genital organs (external), including traumatic amputation	OPEN WOUND OF PENIS
879	Open wound of other and unspecified sites, except limbs	OPEN WOUND OF BREAST
880	Open wound of shoulder and upper arm	OPEN WOUND OF SHOULDER
881	Open wound of elbow, forearm, and wrist	OPEN WOUND OF FOREARM
882	Open wound of hand except finger(s) alone	OPEN WOUND OF HAND
883	Open wound of finger(s)	OPEN WOUND OF FINGER
884	Multiple and unspecified open wound of upper limb	OPEN WOUND ARM MULT/NOS
885	Traumatic amputation of thumb (complete) (partial)	AMPUTATION THUMB
886	Traumatic amputation of other finger(s) (complete) (partial)	AMPUTATION FINGER
887	Traumatic amputation of arm and hand (complete) (partial)	AMPUT BELOW ELB, UNILAT
890	Open wound of hip and thigh	OPEN WOUND OF HIP/THIGH
891	Open wound of knee, leg [except thigh], and ankle	OPEN WND KNEE/LEG/ANKLE
892	Open wound of foot except toe(s) alone	OPEN WOUND OF FOOT
893	Open wound of toe(s)	OPEN WOUND OF TOE
894	Multiple and unspecified open wound of lower limb	OPEN WOUND OF LEG NEC
895	Traumatic amputation of toe(s) (complete) (partial)	AMPUTATION TOE
896	Traumatic amputation of foot (complete) (partial)	AMPUTATION FOOT, UNILAT
897	Traumatic amputation of leg(s) (complete) (partial)	AMPUT BELOW KNEE, UNILAT
900	Injury to blood vessels of head and neck	INJUR CAROTID ARTERY NOS
901	Injury to blood vessels of thorax	INJURY THORACIC AORTA
902	Injury to blood vessels of abdomen and pelvis	INJURY ABDOMINAL AORTA
903	Injury to blood vessels of upper extremity	INJ AXILLARY VESSEL NOS
904	Injury to blood vessels of lower extremity and unspecified sites	INJ COMMON FEMORAL ARTER
905	Late effects of musculoskeletal and connective tissue injuries	LATE EFFEC SKULL/FACE FX
906	Late effects of injuries to skin and subcutaneous tissues	LT EFF OPN WND HEAD/TRNK
907	Late effects of injuries to the nervous system	LT EFF INTRACRANIAL INJ
908	Late effects of other and unspecified injuries	LATE EFF INT INJUR CHEST
909	Late effects of other and unspecified external causes	LATE EFF DRUG POISONING
910	Superficial injury of face, neck, and scalp except eye	ABRASION HEAD
911	Superficial injury of trunk	ABRASION TRUNK
912	Superficial injury of shoulder and upper arm	ABRASION SHOULDER/ARM

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

913	Superficial injury of elbow, forearm, and wrist	ABRASION FOREARM
914	Superficial injury of hand(s) except finger(s) alone	ABRASION HAND
915	Superficial injury of finger(s)	ABRASION FINGER
916	Superficial injury of hip, thigh, leg, and ankle	ABRASION HIP & LEG
917	Superficial injury of foot and toe(s)	ABRASION FOOT & TOE
918	Superficial injury of eye and adnexa	SUPERFIC INJ PERIOCLAR
919	Superficial injury of other, multiple, and unspecified sites	ABRASION NEC
920	Contusion of face, scalp, and neck except eye(s)	CONTUSION FACE/SCALP/NCK
921	Contusion of eye and adnexa	BLACK EYE NOS
922	Contusion of trunk	CONTUSION OF BREAST
923	Contusion of upper limb	CONTUSION SHOULDER REG
924	Contusion of lower limb and of other and unspecified sites	CONTUSION OF THIGH
925	Crushing injury of face, scalp, and neck	
926	Crushing injury of trunk	CRUSH INJ EXT GENITALIA
927	Crushing injury of upper limb	CRUSH INJ SHOULDER REG
928	Crushing injury of lower limb	CRUSHING INJURY THIGH
929	Crushing injury of multiple and unspecified sites	CRUSH INJ MULT SITE NEC
930	Foreign body on external eye	CORNEAL FOREIGN BODY
931	Foreign body in ear	FOREIGN BODY IN EAR
932	Foreign body in nose	FOREIGN BODY IN NOSE
933	Foreign body in pharynx and larynx	FOREIGN BODY IN PHARYNX
934	Foreign body in trachea, bronchus, and lung	FOREIGN BODY IN TRACHEA
935	Foreign body in mouth, esophagus, and stomach	FOREIGN BODY IN MOUTH
936	Foreign body in intestine and colon	FB IN INTESTINE & COLON
937	Foreign body in anus and rectum	FOREIGN BODY ANUS/RECTUM
938	Foreign body in digestive system, unspecified	FOREIGN BODY GI NOS
939	Foreign body in genitourinary tract	FB BLADDER & URETHRA
940	Burn confined to eye and adnexa	CHEMICAL BURN PERIOCLAR
941	Burn of face, head, and neck	BURN NOS HEAD-UNSPEC
942	Burn of trunk	BURN NOS TRUNK-UNSPEC
943	Burn of upper limb, except wrist and hand	BURN NOS ARM-UNSPEC
944	Burn of wrist(s) and hand(s)	BURN NOS HAND-UNSPEC
945	Burn of lower limb(s)	BURN NOS LEG-UNSPEC
946	Burns of multiple specified sites	BURN NOS MULTIPLE SITE
947	Burn of internal organs	BURN OF MOUTH & PHARYNX
948	Burns classified according to extent of body surface involved	BDY BRN < 10%/3D DEG NOS
949	Burn, unspecified	BURN NOS
950	Injury to optic nerve and pathways	OPTIC NERVE INJURY
951	Injury to other cranial nerve(s)	INJURY OCULOMOTOR NERVE
952	Spinal cord injury without evidence of spinal bone injury	C1-C4 SPIN CORD INJ NOS

Appendix A

ICD-9-CM Diagnosis and Procedure Code Tables

953	Injury to nerve roots and spinal plexus	CERVICAL ROOT INJURY
954	Injury to other nerve(s) of trunk, excluding shoulder and pelvic girdles	INJ CERV SYMPATH NERVE
955	Injury to peripheral nerve(s) of shoulder girdle and upper limb	INJURY AXILLARY NERVE
956	Injury to peripheral nerve(s), of pelvic girdle and lower limb	INJURY SCIATIC NERVE
957	Injury to other and unspecified nerves	INJ SUPERF NERV HEAD/NCK
958	Certain early complications of trauma	AIR EMBOLISM
959	Injury, other and unspecified	
960	Poisoning by antibiotics	POISONING-PENICILLINS
961	Poisoning by other anti-infectives	POISONING-SULFONAMIDES
962	Poisoning by hormones and synthetic substitutes	POIS-CORTICOSTEROIDS
963	Poisoning by primarily systemic agents	POIS-ANTIALLRG/ANTIEMET
964	Poisoning by agents primarily affecting blood constituents	POISONING-IRON/COMPOUNDS
965	Poisoning by analgesics, antipyretics, and antirheumatics	POISONING-OPIUM NOS
966	Poisoning by anticonvulsants and anti-Parkinsonism drugs	POISON-OXAZOLIDINE DERIV
967	Poisoning by sedatives and hypnotics	POISONING-BARBITURATES
968	Poisoning by other central nervous system depressants and anesthetics	POIS-CNS MUSCLE DEPRESS
969	Poisoning by psychotropic agents	POISON-ANTIDEPRESNT NOS
970	Poisoning by central nervous system stimulants	POISONING-ANALEPTICS
971	Poisoning by drugs primarily affecting the autonomic nervous system	POIS-PARASYMPATHOMIMETIC
972	Poisoning by agents primarily affecting the cardiovascular system	POIS-CARD RHYTHM REGULAT
973	Poisoning by agents primarily affecting the gastrointestinal system	POIS-ANTACID/ANTIGASTRIC
974	Poisoning by water, mineral, and uric acid metabolism drugs	POIS-MERCURIAL DIURETICS
975	Poisoning by agents primarily acting on the smooth and skeletal muscles and respiratory system	POISONING-OXYTOCIC AGENT
976	Poisoning by agents primarily affecting skin and mucous membrane, ophthalmological, otorhinolaryngological, and dental drugs	POIS-LOCAL ANTI-INFECT
977	Poisoning by other and unspecified drugs and medicinal substances	POISONING-DIETETICS
978	Poisoning by bacterial vaccines	POISONING-BCG VACCINE
979	Poisoning by other vaccines and biological substances	POISON-SMALLPOX VACCINE
980	Toxic effect of alcohol	TOXIC EFF ETHYL ALCOHOL
981	Toxic effect of petroleum products	TOXIC EFF PETROLEUM PROD
982	Toxic effect of solvents other than petroleum-based	TOXIC EFFECT BENZENE
983	Toxic effect of corrosive aromatics, acids, and caustic alkalis	TOX EFF CORROSIVE AROMAT

Appendix A


ICD-9-CM Diagnosis and Procedure Code Tables

984	Toxic effect of lead and its compounds (including fumes)	TX EFF INORG LEAD COMPND
985	Toxic effect of other metals	TOXIC EFFECT MERCURY
986	Toxic effect of carbon monoxide	TOX EFF CARBON MONOXIDE
987	Toxic effect of other gases, fumes, or vapors	TOXIC EFF LIQ PETROL GAS
988	Toxic effect of noxious substances eaten as food	TOXIC EFF FISH/SHELLFISH
989	Toxic effect of other substances, chiefly nonmedicinal as to source	TOXIC EFFECT CYANIDES
990	Effects of radiation, unspecified	EFFECTS RADIATION NOS
991	Effects of reduced temperature	FROSTBITE OF FACE
992	Effects of heat and light	HEAT STROKE & SUNSTROKE
993	Effects of air pressure	BAROTRAUMA, OTITIC
994	Effects of other external causes	EFFECTS OF LIGHTNING
995	Certain adverse effects not elsewhere classified	ANAPHYLACTIC SHOCK
996	Complications peculiar to certain specified procedures	MALFUNC CARD DEV/GRF NOS
997	Complications affecting specified body systems, not elsewhere classified	NERVOUS SYST COMPLC NOS
998	Other complications of procedures, not elsewhere classified	POSTOPERATIVE SHOCK
999	Complications of medical care, not elsewhere classified	GENERALIZED VACCINIA

Navigating the Blood Management Project Data Collection Tool


How to Log In and Get Started

- 1) Once you have registered and received your confirmation to submit data for the Blood Management Project, you may access the project website at:
<http://manual.jointcommission.org>
- 2) Click on “Login” in the upper right hand corner.



The screenshot shows the homepage of The Joint Commission. The header includes the logo and the text "The Joint Commission" with links for "Login", "Register", and "Print". A sidebar on the left contains a "HOME" section with a link to "Site Home". The main content area features a "Welcome to the Performance Measurement Network Q&A Forum" message, followed by a "Published Manuals" section. This section contains two columns of links: "Joint Commission Only Measures" and "CMS and Joint Commission Aligned Measures". The "Joint Commission Only Measures" column lists an "UPDATED Hospital Based Psychiatric Inpatient Services (HBIPS) and Perinatal Care (PC) Measures (version 2010A2)" with sub-links for "Original release (version 2010A)" and "1st update (version 2010A1)". The "CMS and Joint Commission Aligned Measures" column lists "Current Specification Manual for National Hospital Quality Measures", "Future Specification Manual for National Hospital Quality Measures", and "Historical Specification Manuals for National Hospital Quality Measures". Below this, an "Important publications:" section mentions a contribution by Dr. Mark Chassin to the publication of "Cottage Industry to Postindustrial Care — The Revolution in Health Care Delivery" in the New England Journal of Medicine.


- 3) Enter your Login and Password and click “ok”.



The login form is titled "Welcome to the Performance Measurement Network" and asks the user to "Please enter your username and password." It features two input fields: "Login:" with the text "testuser50" and a red double asterisk (**) to its right, and "Password:" with a masked password "*****" and a red double asterisk (**) to its right. Below the fields are three buttons: "OK" (green), "Clear" (blue), and "Cancel" (red). At the bottom, there is a link "See also: [Create Login/Register](#), [Forgot password?](#)" and a contact instruction: "Contact SWilliams@jointcommission.org if you have any questions."

Navigating the Blood Management Project Data Collection Tool

- 4) Welcome to the Performance Measurement Network. Select the “Blood Mgmt Project” link from the left hand navigation bar.



- 5) You are now on the Blood Management Project Page. You will see your hospital(s) listed here. In the Project Help section, you will find a link to the measure specifications, an example of the import file template, and other material intended to assist you with your participation in this project. Please click on the hospital name to enter blood management data.



Navigating the Blood Management Project Data Collection Tool

6) You are now on your hospital page. From this page, you can:

- update your hospital demographic information
- enter new records
- import new records
- view and update existing records
- add RBC, Plasma and Platelet events
- mark records as “complete”
- review records that have been completed
- view import attachments

Each function will be discussed in detail below.

**The Joint Commission**

Test User 50 Logout Print
Edit Attach Move Raw Diffs More

BLOODMGMTPROJ

- Site Home
- Post a question
- Blood Mgmt Project

Sample Staff Hospital

333 Somewhere Place, Smalltown, NC 28605
Health Care Organization ID: 44444
Contact Person: Pleasant Contact
Phone: (828) 260-5555
Email: someone@smalltown.us

Submitted Data

[Show all Records \(including complete\)](#)

UBCI	Birthdate	Admitted	Discharged	Completed 
333333	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
333331	05-01-2001	01-01-2010	01-10-2010	<input type="checkbox"/>
333334	05-01-2001	01-01-2010	01-10-2010	<input type="checkbox"/>
333332	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
333335	05-01-2001	01-01-2010	01-10-2010	<input type="checkbox"/>
1234567	12-30-2008	01-26-2010	02-02-2010	<input type="checkbox"/>
2223	05/01/01	01/01/10	01/10/10	<input type="checkbox"/>
333336	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>

 [Enter New Client Record](#)

Import Data

Steps for importing base data set using a properly formatted Excel spreadsheet:

1. Save the file that is to be imported with the EXACT Name: "import.xls".
2. Click the link:  [Import](#) and follow the instructions to select and upload your "import.xls" file.
3. Once you have uploaded the file,  [Click here](#) to finish the upload process.

Navigating the Blood Management Project Data Collection Tool

Updating your Hospital Demographic Information

a) To update your hospital's demographic information, click the "Edit" link, Fill out the form that appears, and click the "Save" button at the bottom of the form.




Test User 50 Logout Print
Edit Attach Move Raw Diffs More

it Commission

Sample Staff Hospital

333 Somewhere Place, Smalltown, NC 28605
Health Care Organization ID: 44444
Contact Person: Pleasant Contact
Phone: (828) 260-5555
Email: someone@smalltown.us

You will be directed to the Edit form, and you can change your hospital's contact details here. Click "Save" to save your changes, or "Cancel" to exit without saving.



Address: 333 Somewhere Place

City: Smalltown

State: NC

Zip Code: 28605

Contact Person: Pleasant Contact

Contact Phone: (828) 260-5555

Contact Email: someone@smalltown.us

Save Save and Continue Preview Change form Cancel



Navigating the Blood Management Project Data Collection Tool

Importing Records

a) To import data, click on the “Import” link on your hospital home page. The template for this import file can be found on the project home page.


Import Data

Steps for importing base data set using a properly formatted Excel spreadsheet:

1. Save the file that is to be imported with the EXACT Name: “import.xls”.
2. Click the link  [Import](#) and follow the instructions to select and upload your “import.xls” file.
3. Once you have uploaded the file,  [Click here](#) to finish the upload process.
 - a. Once the import has been completed, you will need to click your web browser's "Back" button and then "Refresh" the web page before you will see your new data records.

b) Click on “browse” to find and select your import file (which must be named “import.xls”), and click on “Upload File”. You do not need to check the checkboxes, but you may want to add a comment to keep track of your imports (e.g., April 2010 discharges; 51 records)



Attach file to Sample Staff Hospital



c) Once you have uploaded your file, you will need to click on the “Click here” link to finish the upload process. You’ll then need to click your browser’s “Back” button and “Refresh” your hospital page.

Import Data


Steps for importing base data set using a properly formatted Excel spreadsheet:

1. Save the file that is to be imported with the EXACT Name: “import.xls”.
2. Click the link:  [Import](#) and follow the instructions to select and upload your “import.xls” file.
3. Once you have uploaded the file,  [Click here](#) to finish the upload process.
 - a. Once the import has been completed, you will need to click your web browser's "Back" button and then "Refresh" the web page before you will see your new data records.


d) You may notice a form at the bottom of your hospital page. It displays the most recently imported file. This area will only be used to verify that your import was successful (note the date, time and comments to ensure that it represents the file you imported).

Navigating the Blood Management Project Data Collection Tool


Attachments ▾

Attachment	Action	Size	Date	Who
 import.xls	props, move	55.0 K	22 Feb 2010 - 08:20	ScottWilliams
Monday 2/22 test of import				

e) Your uploaded records are shown here (in a rather unappealing format!) and you will need to click on your browser's "Back" button to return to your hospital home page.




f) You are now back on your hospital's home page. Please click on your browser's "Refresh" button to view the records you just imported. Your records have been imported, but you will not be able to see them until the page is refreshed (or you navigate away from it and then back to it).



g) Your uploaded files should now be viewable in the "Submitted Data" section of your hospital home page.

Navigating the Blood Management Project Data Collection Tool

[Show all Records \(including complete\)](#)

UBCI	Birthdate	Admitted	Discharged	Completed 
333333	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
333331	05-01-2001	01-01-2010	01-10-2010	<input type="checkbox"/>
555555	04-04-1974	07-04-2009	07-07-2009	<input type="checkbox"/>
333332	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
333335	05-01-2001	01-01-2010	01-10-2010	<input type="checkbox"/>
1234567	12-30-2008	01-26-2010	02-02-2010	<input type="checkbox"/>
2223	05/01/01	01/01/10	01/10/10	<input type="checkbox"/>
333336	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
555556	12-09-1970	08-08-2009	08-12-2009	<input type="checkbox"/>

Navigating the Blood Management Project Data Collection Tool


Enter New Records (without using the file import)


a) To enter a new record, click on the “Enter New Client Record” link (right below the data record table).





b) You are now viewing the data collection tool for Blood Management. Enter data for the client record. Note: hovering over the green “i” next to a data element will show you the question and allowable values associated with that data element as well as a link to the data element page.


Draft Data Collection Tool


Unique Blinded Case Identifier: 


Admission Date: MM-DD-YYYY 

Birthdate: MM-DD-YYYY 


Discharge Date: MM-DD-YYYY 

Discharge Status: 


Sex: ☐ M ☐ F ☐ U 


ICD-9-CM Principal Diagnosis Code: 

ICD-9-CM Other Diagnosis Codes


ICD-9-CM Other Diagnosis Codes: 


[Add another response](#)

ICD-9-CM Principal Procedure Code: 


ICD-9-CM Principal Procedure Date: 


ICD-9-CM Other Procedures Codes/Dates


ICD-9-CM Other Procedure Codes: 


ICD-9-CM Other Procedure Dates: 


[Add another response](#)


Elective Surgery: ☐ 1 ☐ 2 


Transfusion Consent: ☐ 1 ☐ 2 ☐ 3 

Education Addressed Risks, Benefits And Alternatives To Transfusion: ☐ 1 ☐ 2 

Preoperative Blood Type Testing: ☐ Y ☐ N 

Preoperative Anemia Screening: ☐ 1 ☐ 2 ☐ 3 

Preoperative Anemia Screening Data: MM-DD-YYYY or UTD 

Anesthesia Start Date: 











c) Once you have completed data entry for this record, click on “Save Data Record”.

**Navigating the Blood Management Project
Data Collection Tool
To View and Update Existing Records**

a) There are two ways to view the list of submitted records. The default view is of all incomplete records. If you would like to view all records, including completed (locked) records, click the link “Show all Records (including complete)”.

View of the default setting showing a list of only incomplete records:
















[Show all Records \(including complete\)](#)

UBCI	Birthdate	Admitted	Discharged	Completed 
333333	03-03-1983	02-02-2010	02-05-2010	
333331	05-01-2001	01-01-2010	01-10-2010	
555555	04-04-1974	07-04-2009	07-07-2009	
333332	03-03-1983	02-02-2010	02-05-2010	
333335	05-01-2001	01-01-2010	01-10-2010	
1234567	12-30-2008	01-26-2010	02-02-2010	
2223	05/01/01	01/01/10	01/10/10	
333336	03-03-1983	02-02-2010	02-05-2010	
555556	12-09-1970	08-08-2009	08-12-2009	

View of alternate setting showing list of all records (both incomplete and complete). To return the default setting, click the link “Show Incomplete Records Only”

Navigating the Blood Management Project Data Collection Tool

[Show incomplete Records Only](#)

UBCI	Birthdate	Admitted	Discharged	Completed 
333333	03-03-1983	02-02-2010	02-05-2010	
333331	05-01-2001	01-01-2010	01-10-2010	
555555	04-04-1974	07-04-2009	07-07-2009	
333332	03-03-1983	02-02-2010	02-05-2010	
1234567	12-30-2008	01-26-2010	02-02-2010	
333335	05-01-2001	01-01-2010	01-10-2010	
333336	03-03-1983	02-02-2010	02-05-2010	
2223	05/01/01	01/01/10	01/10/10	
555556	12-09-1970	08-08-2009	08-12-2009	
333334	05-01-2001	01-01-2010	01-10-2010	
99999999	01-01-1901	11-11-2010	11-15-2010	
4445	03/03/83	02/02/10	02/05/10	
444555	03/03/83	02/02/10	02/05/10	
2224	05/01/01	01/01/10	01/10/10	

b) To view or update data in an existing record, click on the UBCI number. This will create a drop down that includes all of the information for that client record. You can contract the drop down by clicking on the “-“ or expand by clicking on the “+” before the different sections.

Navigating the Blood Management Project Data Collection Tool

1234567	12-30-2008	01-26-2010	02-02-2010
<div> <input type="checkbox"/> General and other patient-level data elements </div>			
Discharge Status 01			
Sex M			
ICD-9-CM Principal Diagnosis Code 49301			
ICD-9-CM Other Diagnosis Codes			
ICD-9-CM Principal Procedure Code 7301			
ICD-9-CM Principal Procedure Date 01-25-2010			
ICD-9-CM Other Procedure Codes			
ICD-9-CM Other Procedure Dates			
Transfusion Consent			
Education Addressed Risks, Benefits and Alternatives to Transfusion			
Elective Surgery			
Anesthesia Start Date			
Preoperative Anemia Screening Date			
Preoperative Anemia Screening			
Preoperative Blood Type Testing			
<div> <input type="checkbox"/> Measure Set Specific Data Elements </div>			
<div> <input type="checkbox"/> RBC Event(s) </div>			
<div> Add RBC Event record (3 left) </div>			
<div> <input type="checkbox"/> Plasma Event(s) </div>			
<div> Add Plasma Event record (3 left) </div>			
<div> <input type="checkbox"/> Platelet Event(s) </div>			
<div> Add Platelet Event record (3 left) </div>			

c) To edit the “General and other patient-level data elements”, click on the pencil icon.


1234567	12-30-2008	01-26-2010	02-02-2010
<div> <input type="checkbox"/> General and other patient-level data elements </div>			
Discharge Status 01			
Sex M			
ICD-9-CM Principal Diagnosis Code 49301			
ICD-9-CM Other Diagnosis Codes			
ICD-9-CM Principal Procedure Code 7301			
ICD-9-CM Principal Procedure Date 01-25-2010			
ICD-9-CM Other Procedure Codes			


d) Make changes to the “General and other patient-level data elements” and click “Save” when you are done.


Navigating the Blood Management Project Data Collection Tool


▼ Form Data ► Permissions


Draft Data Collection Tool


Unique Blinded Case Identifier 


Admission Date MM-DD-YYYY 

Birthdate 


Discharge Date 

Discharge Status 

Sex ☒ M ☐ F ☐ U 

ICD-9-CM Principal Diagnosis Code 

ICD-9-CM Other Diagnosis Codes

ICD-9-CM Other Diagnosis Codes 

Save Save and Continue Preview Change form Cancel ☐ New Revision

Navigating the Blood Management Project Data Collection Tool

Add RBC Events and BM Unit Level Data Elements

a) To add a RBC event (NOTE: you can add up to three RBC events), click on the “Add RBC Event Record” Link.

333331 05-01-2001 01-01-2010 01-10-2010

- [-] General and other patient-level data elements
- [-] Measure Set Specific Data Elements
 - [-] RBC Event(s)
 - [Add RBC Event record \(3 left\)](#)
 - [-] Plasma Event(s)
 - [Add Plasma Event record \(3 left\)](#)
 - [-] Platelet Event(s)
 - [Add Platelet Event record \(3 left\)](#)

b) Enter data for RBC Event 1 and click “Save Data Record”

RBC Event

RBC Event ID ☒ 1 ☐ 2 ☐ 3

RBC Event Total Doses

Clinical Indication For RBCs

Pre-transfusion Hemoglobin

Pre-transfusion Hematocrit

Surgical Procedure ☐ 1 ☐ 2

c) Data for “RBC Event 1” is now included with this client record. To edit the RBC Event data that you just entered, click on the pencil icon next to the event. To add unit level data for RBC Event 1, click on the “Add BM Unit Level Data Elements Record” link. (NOTE: you can add up to three BM Unit Level Records)

Navigating the Blood Management Project Data Collection Tool

333331 05-01-2001 01-01-2010 01-10-2010

- ⊕ General and other patient-level data elements
- ⊖ Measure Set Specific Data Elements
 - ⊖ RBC Event(s)
 - ⊖ RBC Event 1

RBC Event ID	1
RBC Event Total Doses	2
Clinical Indication for RBCs	1
Pre-transfusion Hemoglobin	8
Pre-transfusion Hematocrit	21
Surgical Procedure	1
 - ⊖ BM Unit Level Data Elements(s)
 - [Add BM Unit Level Data Elements record \(3 left\)](#)
 - [Add RBC Event record \(2 left\)](#)
 - ⊖ Plasma Event(s)
 - [Add Plasma Event record \(3 left\)](#)
 - ⊖ Platelet Event(s)
 - [Add Platelet Event record \(3 left\)](#)

d) Enter data for the BM Unit Level Record for RBC Event 1 and click “Save Data Record”

BM Unit Level Data Elements

Transfusion Start Date

Transfusion Start Time

Transfusion Order ☐ Y ☐ N

Patient ID Verification ☐ 1 ☐ 2

Vital Sign Monitoring ☐ 1 ☐ 2

[Save Data Record](#)

e) Data for “BM Unit 1” for “RBC Event 1” is now included with this client record. To edit the BM unit data that you just entered, click on the pencil icon. To add another BM Unit for RBC Event 1, click on “Add BM Unit Level Data Elements Record” link. To add another RBC Event, click on “Add RBC Event Record”.

Navigating the Blood Management Project Data Collection Tool

333331	05-01-2001	01-01-2010	01-10-2010
--------	------------	------------	------------


- ⊕ General and other patient-level data elements
- ⊖ Measure Set Specific Data Elements
 - ⊖ RBC Event(s)
 - ⊖ RBC Event 1
 - RBC Event ID 1
 - RBC Event Total Doses 2
 - Clinical Indication for RBCs 1
 - Pre-transfusion Hemoglobin 8
 - Pre-transfusion Hematocrit 21
 - Surgical Procedure 1
 - ⊖ BM Unit Level Data Elements(s)
 - ⊖ BM Unit Level Data Elements 1
 - Transfusion Start Date 03-03-2010
 - Transfusion Start Time 11:00
 - Transfusion Order Y
 - Patient ID Verification 1
 - Vital Sign Monitoring 1
 - [Add BM Unit Level Data Elements record \(2 left\)](#)
 - [Add RBC Event record \(2 left\)](#)
 - ⊖ Plasma Event(s)
 - [Add Plasma Event record \(3 left\)](#)
 - ⊖ Platelet Event(s)
 - [Add Platelet Event record \(3 left\)](#)

Navigating the Blood Management Project Data Collection Tool

Add Plasma Events and BM Unit Level Data Elements


a) To add a Plasma event, click on the “Add Plasma Event Record” Link


333331	05-01-2001	01-01-2010	01-10-2010
--------	------------	------------	------------


- General and other patient-level data elements 
- Measure Set Specific Data Elements
 - RBC Event(s)
 - [Add RBC Event record \(3 left\)](#)
 - Plasma Event(s)
 - [Add Plasma Event record \(3 left\)](#)
 - Platelet Event(s)
 - [Add Platelet Event record \(3 left\)](#)


b) Enter data for Plasma Event 1 and click “Save Data Record”

Plasma Event

Plasma Event ID  ☐ 1 ☐ 2 ☐ 3

Plasma Event Total Doses 

Clinical Indication For Plasma 

Pre-transfusion Laboratory Testing  ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5

[Save Data Record](#)

Navigating the Blood Management Project Data Collection Tool

c) Data for “Plasma Event 1” is now included with this client record. To edit the Plasma Event data that you just entered, click on the pencil icon next to the event. To add unit level data for Plasma Event 1, click on the “Add BM Unit Level Data Elements Record” link. (NOTE: you can add up to three BM Unit Level Records)

333331	05-01-2001	01-01-2010	01-10-2010
+ General and other patient-level data elements			
[-] Measure Set Specific Data Elements			
+ RBC Event(s)			
[-] Plasma Event(s)			
[-] Plasma Event 1			
Plasma Event ID			1
Plasma Event Total Doses			2
Clinical Indication for Plasma			1
Pre-transfusion Laboratory Testing			2
[-] BM Unit Level Data Elements(s)			
Add BM Unit Level Data Elements record (3 left)			
Add Plasma Event record (2 left)			
[-] Platelet Event(s)			
Add Platelet Event record (3 left)			

d) Enter data for the BM Unit Level Record for Plasma Event 1 and click “Save Data Record”

BM Unit Level Data Elements

Transfusion Start Date

Transfusion Start Time

Transfusion Order ☐ Y ☐ N

Patient ID Verification ☐ 1 ☐ 2

Vital Sign Monitoring ☐ 1 ☐ 2

Navigating the Blood Management Project Data Collection Tool

e) Data for “BM Unit Level 1” for “Plasma Event 1” is now included with this client record. To edit the BM unit data that you just entered, click on the pencil icon. To add another BM Unit for Plasma Event 1, click on “Add BM Unit Level Data Elements Record” link. To add another Plasma Event, click on “Add Plasma Event Record”.

333331	05-01-2001	01-01-2010	01-10-2010
--------	------------	------------	------------

- General and other patient-level data elements
- Measure Set Specific Data Elements
 - RBC Event(s)
 - Plasma Event(s)
 - Plasma Event 1

Plasma Event ID	1
Plasma Event Total Doses	2
Clinical Indication for Plasma	1
Pre-transfusion Laboratory Testing	2
 - BM Unit Level Data Elements(s)
 - BM Unit Level Data Elements 1

Transfusion Start Date	03-03-2010
Transfusion Start Time	11:00
Transfusion Order	Y
Patient ID Verification	1
Vital Sign Monitoring	1
 - [Add BM Unit Level Data Elements record \(2 left\)](#)
 - [Add Plasma Event record \(2 left\)](#)
- Platelet Event(s)
 - [Add Platelet Event record \(3 left\)](#)

Navigating the Blood Management Project Data Collection Tool

Add Platelet Events and BM Unit Level Data Elements

a) To add a Platelet event, click on the “Add Platelet Event Record” Link

333331	05-01-2001	01-01-2010	01-10-2010
--------	------------	------------	------------

- General and other patient-level data elements
- Measure Set Specific Data Elements
 - RBC Event(s)
 - [Add RBC Event record \(3 left\)](#)
 - Plasma Event(s)
 - [Add Plasma Event record \(3 left\)](#)
 - Platelet Event(s)
 - [Add Platelet Event record \(3 left\)](#)

b) Enter data for Platelet Event 1 and click “Save Data Record”

Platelet Event

Platelet Event ID ☐ 1 ☐ 2 ☐ 3

Platelet Event Total Doses

Clinical Indication For Platelets

Pre-transfusion Platelet Count

Pre-transfusion Platelet Testing ☐ 1 ☐ 2

[Save Data Record](#)

Navigating the Blood Management Project Data Collection Tool

c) Data for “Platelet Event 1” is now included with this client record. To edit the Platelet Event data that you just entered, click on the pencil icon next to the event. To add unit level data for Platelet Event 1, click on the “Add BM Unit Level Data Elements Record” link. (NOTE: you can add up to three BM Unit Level Records)

333331	05-01-2001	01-01-2010	01-10-2010
--------	------------	------------	------------

General and other patient-level data elements

Measure Set Specific Data Elements

- RBC Event(s)**
- Plasma Event(s)**
- Platelet Event(s)**
 - Platelet Event 1

Platelet Event ID	1
Platelet Event Total Doses	3
Clinical Indication for Platelets	1
Pre-transfusion Platelet Count	100
Pre-transfusion Platelet Testing	1
- BM Unit Level Data Elements(s)**
 - [Add BM Unit Level Data Elements record \(3 left\)](#)
 - [Add Platelet Event record \(2 left\)](#)

d) Enter data for the BM Unit Level Record for Platelet Event 1 and click “Save Data Record”

BM Unit Level Data Elements

Transfusion Start Date

Transfusion Start Time

Transfusion Order

☐ Y ☐ N

Patient ID Verification

☐ 1 ☐ 2

Vital Sign Monitoring

☐ 1 ☐ 2

Save Data Record

Navigating the Blood Management Project Data Collection Tool

e) Data for “BM Unit Level 1” for “Platelet Event 1” is now included with this client record. To edit the BM unit data that you just entered, click on the pencil icon. To add another BM Unit for Platelet Event 1, click on “Add BM Unit Level Data Elements Record” link. To add another Platelet Event, click on “Add Platelet Event Record”.

333331	05-01-2001	01-01-2010	01-10-2010
<div> <div>+</div> General and other patient-level data elements <div></div> </div>			
<div> <div>-</div> Measure Set Specific Data Elements </div>			
<div> <div>+</div> RBC Event(s) </div>			
<div> <div>+</div> Plasma Event(s) </div>			
<div> <div>-</div> Platelet Event(s) </div>			
<div> <div>-</div> Platelet Event 1 <div></div> </div>			
Platelet Event ID			1
Platelet Event Total Doses			3
Clinical Indication for Platelets			1
Pre-transfusion Platelet Count			100
Pre-transfusion Platelet Testing			1
<div> <div>-</div> BM Unit Level Data Elements(s) </div>			
<div> <div>-</div> BM Unit Level Data Elements 1 <div></div> </div>			
Transfusion Start Date			03-03-2010
Transfusion Start Time			11:00
Transfusion Order			Y
Patient ID Verification			1
Vital Sign Monitoring			1
<div> <div></div> Add BM Unit Level Data Elements record (2 left) </div>			
<div> <div></div> Add Platelet Event record (2 left) </div>			

Navigating the Blood Management Project Data Collection Tool

Marking Records As “Complete”

a) Once you are done entering and editing data for a record, you will need to mark the record as complete. **Please note: Once you check the box for a record under “Complete” you are BOTH marking the record as complete AND locking that record for any further editing.** When you click on the checkbox, the record will “disappear” from view. Do not be alarmed. The default view of the table is to only show incomplete records. To view the record you just completed, click on the link to “Show all Records (including complete)”

Show all Records (including complete)				
UBCI	Birthdate	Admitted	Discharged	Completed 
333333	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
333331	05-01-2001	01-01-2010	01-10-2010	<input type="checkbox"/>
555555	04-04-1974	07-04-2009	07-07-2009	<input type="checkbox"/>
333332	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
333335	05-01-2001	01-01-2010	01-10-2010	<input checked="" type="checkbox"/>
1234567	12-30-2008	01-26-2010	02-02-2010	<input type="checkbox"/>
2223	05/01/01	01/01/10	01/10/10	<input type="checkbox"/>
333336	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
555556	12-09-1970	08-08-2009	08-12-2009	<input type="checkbox"/>

Navigating the Blood Management Project Data Collection Tool

Reviewing Records That Have Been Completed

a) To review a record that has been marked complete, switch the view on your hospital home page by clicking on the “Show all Records (including complete)” link.

Submitted Data

Show all Records (including complete)

b) In this view you can see all records both complete and incomplete. Completed records are now LOCKED and can not be edited.

[Show incomplete Records Only](#)

UBCI	Birthdate	Admitted	Discharged	Completed
333333	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
333331	05-01-2001	01-01-2010	01-10-2010	<input type="checkbox"/>
555555	04-04-1974	07-04-2009	07-07-2009	<input type="checkbox"/>
333332	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
1234567	12-30-2008	01-26-2010	02-02-2010	<input type="checkbox"/>
333335	05-01-2001	01-01-2010	01-10-2010	<input type="checkbox"/>
333336	03-03-1983	02-02-2010	02-05-2010	<input type="checkbox"/>
2223	05/01/01	01/01/10	01/10/10	<input type="checkbox"/>
555556	12-09-1970	08-08-2009	08-12-2009	<input type="checkbox"/>
333334	05-01-2001	01-01-2010	01-10-2010	<input checked="" type="checkbox"/>
99999999	01-01-1901	11-11-2010	11-15-2010	<input checked="" type="checkbox"/>
4445	03/03/83	02/02/10	02/05/10	<input checked="" type="checkbox"/>
444555	03/03/83	02/02/10	02/05/10	<input checked="" type="checkbox"/>
2224	05/01/01	01/01/10	01/10/10	<input checked="" type="checkbox"/>

b) If, for any reason, you need to unlock a record, you will need to send an e-mail to the project leader, Harriet Gammon. To send your e-mail request, click on the “lock” icon, and an e-mail form should appear. It will be addressed to Harriet, and the subject line will contain a reference to the specific record.

To...

Gammon, Harriet

Cc...

Subject:

Request to unlock record BloodMgmtProject/RecBmpHco003C333334L0D40188

c) In your e-mail, please briefly explain why the record needs to be unlocked (e.g., Accidentally clicked the “Complete” checkbox).

<p>PATIENT BLOOD MANAGEMENT PERFORMANCE MEASURES PROJECT - Technical Advisory Panel</p>
--

David J. Ballard MD, MSPH, PhD, FACP, Co-Chair
Baylor Health Care System
Dallas, TX

Jonathan H. Waters, MD, Co-Chair
Magee Women's Hospital
University of Pittsburgh
Pittsburgh, PA

Neil Bangs, MS, MT (ASCP) SBB
Virginia Commonwealth University Medical Systems
Richmond, Virginia

Richard J. Benjamin, MD, PhD, FRCPath, MS
American Red Cross, National Headquarters
Washington, DC

Laurence Bilfield, MD
Cleveland Clinic HS - Lutheran
Cleveland, OH

Victor A. Ferraris, MD, PhD
Division of Cardiovascular & Thoracic Surgery
University of Kentucky Chandler Medical Center
Lexington, KY

John Freedman, MD, FPCPC
St. Michael's Hospital
University of Toronto
Toronto, Ontario, Canada

Jonathan C. Goldsmith, MD
Division of Blood Diseases and Resources
National Heart, Lung, and Blood Institute
National Institutes of Health
Bethesda, MD

Lawrence Tim Goodnough, MD
Stanford University Medical Center
Stanford, CA

Penny S. Gozia, MD, FACOG, MBA
St. Joseph's Hospital,
Breese, IL

<p>PATIENT BLOOD MANAGEMENT PERFORMANCE MEASURES PROJECT - Technical Advisory Panel</p>

Jerry Holmberg, PhD., MT (ASCP) SBB
Department of Health and Human Services
Rockville, MD

Joseph E. Kiss, MD
The Institute for Transfusion Medicine
University of Pittsburgh Medical Center
Pittsburgh, PA

Harvey G. Klein, MD
National Institutes of Health
Bethesda, MD

Mark T. Lucas, MPS, RCS, CCP
Denver Cardiovascular Perfusionists
Denver, CO

Vijay K. Maker, MD, FACCS
Advocate Illinois Masonic Hospital
Chicago, IL

John (Jeffrey) McCullough, MD
University of Minnesota
Minneapolis, MN

Aryeh Shander, MD, FCCM, FCCP
Englewood Hospital and Medical Center
Englewood, NJ

Bruce D. Spiess, MD, FAHA
Virginia Commonwealth University Medical Center
Richmond, Virginia

Lynne Uhl, MD
Beth Israel Deaconess Medical Center
Boston, MA

Jeffrey Wagner, BSN, RN
Puget Sound Blood Center
Seattle, WA

Rosalyn Yomtovian, MD
Department of Veterans Affairs, Louis Stokes Medical Center
Case Western Reserve University School of Medicine
Cleveland, OH