



# Trauma Outcomes: Final Environmental Scan Report

---

*FINAL ENVIRONMENTAL SCAN REPORT*

*October 30, 2018*

*This report is funded by the Department of Health and Human Services under contract HHSM-500-2017-00060I HHSM-500-0005.*

## Contents

Introduction .....	3
Project Purpose, Scope, and Approach .....	3
Environmental Scan Strategy .....	4
Purpose and Limitations of Measure Concepts .....	4
Characteristics of Good Measures .....	5
Findings: Measures, Concepts, Tools .....	5
Measures .....	5
Measure Concepts .....	7
Instruments, Scales, and Tools .....	8
Findings: Risk Adjustment .....	8
Risk Adjustment in Measures .....	8
Covariates Used and Discussion .....	9
Findings: Shared Accountability and Attribution .....	10
Findings: Data Sources .....	11
References .....	12
Appendix A: Committee Members, NQF Staff, and Federal Liaisons .....	16
Appendix B: Measure and Measure Concept Inventory .....	19
Appendix C: Environmental Scan Strategy .....	30

## Introduction

Traumatic injuries (both non-intentional and intentional) are the third leading cause of death in the United States, and in the aggregate, these injuries cause more premature death than any other illness or disability.<sup>1</sup> Traumatic injuries are a major public health concern accounting for 39 million emergency department visits and 12.3 million hospital admissions annually, and they were among the highest condition-related expenditure among adults ages 18-64 in 2012 and were also associated with approximately \$670 billion in medical expenses and lost productivity in 2013.<sup>2,3</sup> Further, studies have found that trauma disproportionately affects the young and estimated that 20 percent of trauma deaths were survivable.<sup>4</sup> In addition to loss of life, rehabilitation, quality of life, and mental health status are also key outcomes of interest in trauma care.

The 2016 report by a committee of the National Academies of Medicine (NAM), *A National Trauma Care System: Integrating Military and Civilian Trauma Systems to Achieve Zero Preventable Deaths After Injury*, offered 10 recommendations to help achieve high-quality trauma care on a national level.<sup>1</sup> Two recommendations that are relevant to this work called for a “designated locus of responsibility and authority” to be held accountable for developing a national approach to improve care for trauma patients. The committee also called for governmental, private, and academic partners to work together to collect and share data across the trauma continuum of care to help identify measures that assess the quality of trauma care.

Despite the magnitude and expense of traumatic injuries, few performance measures address the quality of trauma care. While gains have been made in trauma care including better patient stabilization, rapid transfer to appropriate care, resuscitation, and management of hemorrhage, how trauma systems perform as a whole is understudied.<sup>1</sup> Performance measures provide an opportunity to assess key aspects of care for specific conditions or settings of care and identify levers and areas where focused attention can promote improvement in the quality of care. The 2016 NAM Committee noted the absence of standard, national metrics for trauma care, and called for further development of measures in this area.

Measurement related to trauma care presents unique challenges, including risk adjustment and attributing performance across the trauma care continuum, including prehospital care (e.g., emergency medical services and coordination of patient transport) and post-acute care (e.g., rehabilitation). Accountability for trauma care is challenging since leadership can be assigned to states, counties, and cities with minimal federal oversight.<sup>1</sup> Responsibility for patient care and patient outcomes is distributed among multiple stakeholders, including regional and community entities. Measures that promote shared accountability, such as population-level measures, may help to drive greater integration of care and system-wide improvement.

## Project Purpose, Scope, and Approach

The National Quality Forum (NQF), with funding from the Department of Health and Human Services (HHS), convened a multistakeholder Trauma Outcomes Committee ([Appendix A](#)) to provide input and guidance on the identification of developed measures and concepts addressing population-based trauma outcomes. Results of the scan will be used to produce a measurement framework to help

identify areas for measure development and gaps in trauma care. This measurement framework will help to conceptualize measurement strategies related to the quality of trauma care and to address related issues, such as level of analysis, attribution, and risk adjustment. NQF will also issue a final report summarizing the results of the final environmental scan, the measurement framework, the multistakeholder committee's discussion on population-based measurement of trauma care and strategies for future quality measurement efforts related to trauma care.

This work will be accomplished over the course of 12 months through one in-person meeting, and up to seven web meetings with the Committee. Key informant interviews may also be used to obtain additional expert insight not otherwise identified in publicly available sources. This project will add to the existing body of knowledge around trauma measurement and associated challenges, and spur action in areas of measurement that need additional research and development.

## Environmental Scan Strategy

With parameters established in consultation with the HHS Government Task Lead (GTL), the Contracting Office Representative (COR), and the Committee, NQF staff completed an environmental scan of measures and measure concepts to address population-based trauma outcomes. Upon completion of the environmental scan, NQF staff gathered the information and used it to identify measurement gaps. The Committee used the analysis to: (1) provide input and direction on the development of a conceptual framework for analyzing measures to improve the quality of trauma care and (2) identify measurement gaps.

NQF used the search terms outlined in the subsection below and the search parameters ([Appendix C](#)). Note that search words were combined with terms like “measure,” “measurement,” “survey,” “scale,” etc. to help identify relevant measures. Information sources were identified through various resources such as PubMed, as well as grey literature and web searches through Google Scholar to identify reports, white papers, and other documentation related to trauma care and traumatic injuries.

NQF staff initially reviewed abstracts and articles that were relevant to the operational definition and research questions, synthesized the sources, and compiled a list of measure concepts related to trauma care ([Appendix B](#)).

For the environmental scan, NQF staff identified 90 measures from the NQF Quality Positioning System, the Centers for Medicare & Medicaid Services Measures Inventory, HEDIS, the National Trauma Databank, and several other measure inventories. Out of the 90 identified measures, 49 measures were included in the scan. Measures were excluded due to duplication, irrelevance to trauma outcomes, or listing of traumatic injuries as exclusions in the measure specifications. With input from the Committee and NQF members, 238 measure concepts were identified.

## Purpose and Limitations of Measure Concepts

NQF distinguishes between a measure and a measure concept. A **measure** is defined as a fully developed metric that includes detailed specifications and may have undergone scientific testing. A fully developed measure identifies what should happen (what is being measured), who should be measured

(population), where measurement should happen (setting), when it should happen (time), and how it should occur. It is important to note that the Committee is not recommending specific measures for immediate implementation and use. A **measure concept** is an idea for a measure that includes a description of the measure, ideally including planned target and population. Note that some measure concepts are rooted in current work, and others are more forward-thinking ideas with little or no existing research.

## Characteristics of Good Measures

To receive NQF endorsement, measures must meet four criteria: important to measure and report, scientific acceptability, usability and use, and feasibility. As measures related to trauma outcomes are developed, these criteria may guide measure developers as they work to specify and test measures. The first criterion, **important to measure and report**, aims to keep measurement focused on high-priority areas with strong evidence that measurement can have a positive impact on healthcare quality. The **scientific acceptability** criterion assesses whether the measure, when implemented, will produce consistent (reliable) and credible (valid) results about the quality of care. Measures are also assessed for whether they are **usable and relevant**—that is whether the intended users of the measure can understand the measure results and use them in a meaningful way. Finally, the **feasibility** criterion assesses whether data needed for the measure are readily available and retrievable without undue burden.

## Findings: Measures, Concepts, Tools

As previously mentioned, the NAM report called on government, private, and academic agencies to work together to collect and share data to develop measures that address the continuum of trauma care. Measures should include structure, processes, outcomes, access and patient experiences from the point of injury, the emergency department, in-patient care, through rehabilitation. A thorough review of more than 300 measures and concepts identified in the scan resulted in the final list included in [Appendix B](#). Measures and concepts were removed from the inventory if they were not specifically focused on trauma (e.g., general fall prevention measures or care coordination measures) and if the measure specifications included trauma as an exclusion.

## Measures

The environmental scan included 49 measures, including seven NQF-endorsed measures. Measures were found in the Centers for Medicare & Medicaid Services measure inventory and in registries maintained by the American College of Surgeons and the American College of Emergency Physicians and several others. Of the 49 measures identified, 25 are process measures, 12 are outcome measures, nine are structural measures, and three are efficiency measures. Measures focused on the timeliness of transport to trauma centers, timeliness of intervention for hemorrhage and venous thromboembolism prophylaxis, pain assessment for the injured patient, utilization of emergency department resources, and information sharing between hospitals. Eight patient safety indicators were also included at the request of the Committee. While certain process measures are key to assess critical processes like transfer of a patient to a trauma center in a timely manner, this assessment does not guarantee that the

patient received high-quality care. More outcome measures are needed to assess the quality of trauma care.

Table 1 and Table 2 detail measures by data source and level of analysis, respectively. As expected, most of the measures identified are computed via registry, followed by electronic health record and electronic health data, and paper records. Multiple data sources marked “other” included survey data, hospital licensure and regulation data, and management data. With regard to level of analysis, most measures were specified at the facility level, followed by states and EMS organizations in a state or region. Level of analysis was unknown or unavailable for five measures.

**Table 1. Measures by Data Source**

Data source	# of measures
Registry	14
Claims	12
Electronic health record/ electronic health data	7
Paper records	6
Other	11

**Table 2. Measures by Level of Analysis**

Level of analysis	# of measures
Facility	22
State	9
EMS system	4
Clinician: group/practice	10
Unknown	5

Measures were also categorized using the conceptual model used by Stelfox et al. where measures were segmented into one of four categories: prehospital, hospital, posthospital, and secondary prevention (Table 3).<sup>5</sup> Note that the structural measures identified were not included in this table. Similar to findings by Stelfox et al., a majority of the measures identified in this project were for processes and outcomes occurring in the hospital setting.<sup>5</sup> Topic areas included in the hospital category included efficiency, resource utilization, and appropriate use or avoidance of use.

Three measures addressed patient preferences such as palliative care and advance care plans. Five previously endorsed NQF measures described the timing of information shared from the transferring hospital to another healthcare facility. Two mortality measures, one imaging measure, one timeliness measure, and one nonoperative management measure were also included in the inventory. The prehospital measures focused on pain assessment (two measures), pain management (one) and patients transported to trauma centers (one). The sole injury prevention measure addressed alcohol screening in the emergency room.

**Table 3. Measures by Where Measurement Occurs**

Where measurement/intervention occurs	# of measures
Prehospital	4
Hospital	35
Posthospital	0
Secondary prevention	1

## Measure Concepts

More than 200 concepts were identified in the scan. Most concepts contained a proposed topic or title, but many lacked detailed descriptions and detailed information on target populations, how to construct and calculate the measure, and other pertinent information. Concepts were pulled from various sources including from the Eastern Association for the Surgery of Trauma (EAST), the American College of Surgeons Trauma Quality Improvement Program (ACS TQIP), and the literature. Of the 238 concepts identified, 39 were duplicative of other concepts, or there was a fully developed measure based on the concept. From the ACS TQIP guidelines, 56 concepts were identified and 46 were based on EAST guidelines. Note that some guidelines encompassed several protocols that could be developed into individual concepts, but they are not listed in the inventory. Twelve concepts were pulled from the Victorian State Trauma System, and 11 concepts were based on indicators outlined in HHS Health Resources and Services Administration (HRSA) Model Trauma System Planning and Evaluation handbook.<sup>6</sup> Although some guidelines documented varying levels of evidence, they may still provide a path forward to measure and improve trauma outcomes.

Six additional concepts were gathered using data elements collected in the National Trauma Data Bank, five from deliberations at the Hartford Consensus meetings, five from the American Association of Blood Banks (AABB), and two from the Tactical Combat Casualty Care (TCCC) guidelines. The Hartford Consensus meetings were held after the mass shooting at Sandy Hook Elementary School to develop strategies to increase survivability following these events.<sup>7</sup> Some of the metrics to evaluate the effectiveness of their strategy, THREAT (threat suppression; hemorrhage control; rapid extrication to safety; assessment by medical providers; transport to definitive care), were included in this scan as applicable. Patient blood management standards from the AABB were included since they can help guide the transfusion decision making process, decrease blood loss, and reduce the waste of blood products.<sup>8</sup> The TCCC prehospital guidelines used in the military were also consulted and included as applicable.<sup>9</sup>

Concepts addressed indicators that could evaluate the status of trauma systems and specific populations, such as geriatric and pediatric patients, and for specific injuries or conditions including orthopedic trauma, traumatic brain injury, massive transfusion, and hepatic injuries among others. Similar to the findings of a mixed methods study for trauma indicators, the majority of concepts found focused on specific injuries or populations which limits the broad application of the concepts for population-based trauma outcomes.<sup>10</sup> General palliative care concepts and those that addressed specific injuries (e.g., spinal cord injury) and special populations (e.g., pediatric) were also included.

In the literature, 19 concepts (four were removed as they were duplicative of other concepts) were identified in Gruen et al. addressing elements across the trauma care continuum including dispatch criteria, prehospital deaths, triage and transfer, activation of trauma protocols, screening for alcohol and drug use for alcohol or drug related injuries, access to rehabilitation facilities, and return to work.<sup>11</sup> Newer concepts were also discussed including trauma team efficacy,<sup>12</sup> missed injuries,<sup>13</sup> and compliance with specific protocols including trauma team activation,<sup>14</sup> hemorrhage control and anticoagulation, and compliance to trauma alerts.<sup>15</sup> Concepts identified from the international community included hospital survival rates and probability of survival,<sup>16</sup> and destination compliance, discharge status, and major trauma ICU mechanical ventilation hours.<sup>17</sup>

As stated previously, most of the measure concepts had varying levels of evidence or had no evidence base, and the scan did not identify any measures or concepts that assess population-level outcomes for regional trauma systems. Further research is needed to determine the concepts that should be further developed.

## Instruments, Scales, and Tools

Although this work specified that performance measures and concepts should be included, the Committee also identified 61 instruments, tools, or scales that measured other outcomes important for trauma survivors including quality of life, disability, and mental health. Several tools measured quality of life (e.g., SF-12, SF-36, and trauma-specific quality of life (T-QoL), patient-reported outcomes (e.g., Beck's Depression Inventory, Spinal Cord Independence Measure), and longer-term outcomes (e.g., Glasgow Outcome Scale – Extended, European Quality of Life 5D). Scales for populations (e.g., pediatric) and clinical areas (e.g., musculoskeletal) and for patients with ongoing rehabilitation needs were also included (e.g., Rehabilitation Complexity Scale). Although NQF does not consider these individual instruments and scales to be standalone measures, measures could be developed from them. For instance, NQF 0712 *Depression Utilization of the PHQ-9 Tool*, is an example of a measure developed based on a validated instrument.

## Findings: Risk Adjustment

### Risk Adjustment in Measures

Of the measures identified in the environmental scan, five included a risk-adjustment model:

- ACSTrauma2 Mortality Rate Following Blunt Traumatic Injury to the Chest and/or Abdomen
- ACSTrauma3 Mortality Rate Following Penetrating Traumatic Injury to the Chest and/or Abdomen
- ACSTrauma4 Splenic Salvage Rate
- PSI 14 Postoperative Wound Dehiscence

For the ACS trauma measures, the risk adjustment includes race, comorbidities, and transfer status, in addition to a variety of clinical factors such as blood pressure, pulse, or the presence of a spinal injury. PSI 14 and one other untested measure concept of cost of treating intracranial hemorrhage or cerebral infarction, included a risk-adjustment methodology with unspecified parameters.



The National Surgical Quality Improvement Program and Trauma Quality Improvement Program use a risk-adjustment methodology to identify differences between hospitals, generally based on outcome measures of mortality, complications, and cost. Recent publications have suggested applying a similar framework to the assessment of emergency general surgery patients.<sup>18</sup>

Gruen et al.<sup>11</sup> identified a risk-adjusted mortality measure for head injuries as a potential measure concept, while highlighting that such a measure does not yet exist. Gruen et al. also draws a distinction between measures used internally for quality improvement, where risk adjustment may not be needed, and measures used externally for benchmarking or other comparative purposes, including assessing population-based trauma outcomes where “detailed knowledge of factors that affect the outcome of interest, (such as patient age, injury severity, and co-morbid status)” are important components of a valid performance measure.<sup>11</sup>

## Covariates Used and Discussion

In some aspects of trauma care, such as hip fracture mortality assessment, efforts such as Jiang et al.<sup>19</sup> to create a risk-adjustment model were successful, identifying factors such as age, gender, and some co-morbidities as strong components of a model of mortality. Other published approaches to risk adjustment for mortality follow a similar approach,<sup>18,20,21</sup> including a seminal effort based on the National Trauma Data Bank Registry.<sup>22</sup> After reviewing 106 possible covariates, the researchers found that just six covariates—age, hypotension, pulse, the total Glasgow Coma Scale (tGCS), Injury Severity Score, and need for ventilator use—were sufficient to develop a multivariate model with strong predictive power of mortality. However, a review of 286 publications using data from the NTDB indicated that as many as 43 percent did not follow these best practices in evaluating trauma outcomes.<sup>23</sup>

Gruen et al.<sup>11</sup> and other works critiquing extant risk-adjustment approaches concentrate on risk adjustment based on clinical factors, but do not address risk adjustment on the basis of sociodemographic status. Other critiques of existing risk-adjustment models are based on the use of the GCS motor component (mGCS) at admission; Gomez et al.<sup>24</sup> note that while it is extensively used as a covariate for risk-adjustment models used in external benchmarking programs,<sup>25</sup> it does not reflect the effect of possible confounders, and they suggest the highest mGCS score as a supplemental covariate. Their research shows incorporating this new factor results in stronger model performance. Others have suggested increasing the statistical sophistication of the modeling approaching to address volume concerns and the discriminatory power of models.<sup>26</sup>

Other criticism has been directed at the Injury Severity Score (ISS) and the trauma and injury severity score (TRISS), reproaching the score for insufficient predictive power of mortality. Alternatives have been proposed, including the Trauma Risk Adjustment Model (TRAM)<sup>27</sup> and the TMPM (Trauma Mortality Prediction Model).<sup>28</sup> The TRAM includes adjustments such as inclusion of body region and counting the number of comorbidities. The TMPM relies on an empirical analysis of the severity of injuries classified in the Abbreviated Injury Scale (AIS), rather than expert opinion.

## Findings: Shared Accountability and Attribution

Measurement in healthcare has traditionally focused on the performance of individual facilities or practices, seeking to hold providers responsible for adhering to best practices and achieving positive outcomes for patients who are in their direct care. However, with the emergence of accountable care organizations and an increasing focus on population health, there has been a movement toward shared accountability in measurement. Shared accountability is the notion that a wide range of stakeholders may contribute to patient outcomes, and that performance measures should encourage integration and coordination across settings and providers to ensure that high-value, patient-centered care is being delivered across the continuum of care.

Trauma care is well suited to shared accountability approaches, given the distribution of responsibility across various groups and the importance of system-wide planning and coordination to ensure the optimal use of resources and capabilities.<sup>1</sup> Stakeholders are increasingly recognizing that “regionalization”—establishing organized networks to deliver care to populations within defined geographical areas—is likely to be critical for quality improvement in emergency and trauma care.<sup>29,30</sup> Measures that assess the quality and efficiency of care at the regional level may incentivize greater integration of trauma systems and collaboration across sectors.

Community viral load (CVL) is an example of a population-based measure aimed at understanding the quality of care in a defined community. To calculate CVL, viral loads of all HIV-infected persons are aggregated and then used in the monitoring of HIV treatment and its impact on HIV transmission.<sup>31</sup> Low CVL may indicate good uptake of HIV treatment and can provide estimates on HIV incidence. CVL serves as a model metric for measuring population-based outcomes, although it has its limitations, including selection and measurement of viral load, among others.<sup>31</sup> In addition, efforts by the Resuscitation Outcomes Consortium (ROC) have been successful in studying regional variations in survival of out-of-hospital cardiac arrests (OOHCA) across EMS systems.<sup>32</sup> The ROC effort used the Epistry Cardiac Arrest registry to study all out-of-hospital cardiac arrests from 11 sites in the U.S. and Canada and found significant variation in the incidence of OOHCA and associated outcomes. While both CVL and the research conducted by ROC have limitations, both of these approaches can serve as examples of how to approach measurement and improvement of population-based trauma outcomes.

The measures identified in the environmental scan are focused largely on individual aspects of the trauma care continuum (e.g., EMS agencies, hospitals/trauma centers, etc.). Several measures and concepts can be applied at the state level, and may serve as potential examples of measuring system capacity at a population level, but these measures do not allow for evaluation, comparison, and benchmarking between regional trauma networks. The scan did not identify any measures that assess population-level outcomes for regional trauma systems.

Data limitations pose a significant challenge to population-level measurement of outcomes; the ability to aggregate data across the prehospital, hospital, and posthospital settings and link those data to individual patients to track quality over episodes of care remains extremely limited.<sup>33</sup>

Moreover, there is currently no standard way of defining regional trauma networks and attributing patients to those networks for measurement purposes.<sup>33,34</sup> Attribution refers to the methodology used to assign patients and their quality outcomes to providers, clinicians, or other accountable entities.<sup>35</sup> Glickman and colleagues suggest potential approaches to defining emergency care networks for measurement purposes, including the use of geographical boundaries, market-based health referral regions, or government-defined hospital service areas.<sup>34</sup> Carr and colleagues also propose a method for “clustering” hospitals together based on patient use patterns, and suggest that these groupings could serve to define and attribute geographical populations for measurement of emergency care, including trauma.<sup>36</sup> In addition, other initiatives focused on regional coordination of emergency care, such as the federal Hospital Preparedness Program (HPP)—which supports the development of regional healthcare coalitions to improve preparedness for public health emergencies—could provide examples of how regional trauma systems could be defined for measurement purposes.<sup>37</sup> The Health Preparedness Program assesses participating healthcare coalitions on structural and process measures focused on system readiness and adequacy of planning, some of which could potentially be modified or adapted to assess regional trauma system readiness.<sup>38</sup>

Advancing models of attribution that promote improved planning and coordination within regional trauma networks could help promote shared accountability across relevant stakeholders and accelerate quality improvement in trauma care.

## Findings: Data Sources

As noted in the above section, “Findings: Measures, Concepts, Tools,” current measures use a variety of different data sources to measure trauma outcomes. These variations create a foundation to evaluate outcomes by a variety of different, yet complementary, methods. Loosely speaking, the datasets can be grouped into population-level data sets, event-based sets/registries, and patient-level/patient-identifiable data sets. Each of these different data sources has inherent strengths.

At the broadest level, population -level datasets contain large amounts of representative data, often at the national level. Most notably, vital and health statistics datasets can be used for items such as geospatial analysis and sociodemographic factors. The data provided by population-level datasets are often the least specific to trauma outcomes, but are broadly and consistently available. Additionally, population-level datasets often lag and are slow to be updated.

Event-based datasets move towards a finer level of detail that are specific to trauma. Two examples of these are the National Trauma Data Bank (NTDB) and the National Emergency Medicine Services Information System (NEMSIS). Both datasets contain information specific to trauma. Additionally, these datasets collate data from many institutions across the country. The data are stored at the level of the event. Patient identifiable information is not included, and it is not possible to differentiate between a single patient with multiple events as opposed to multiple patients with a single event. Despite the specificity of the data, it is subject to bias as not all institutions submit all cases.

At the most granular level, patient-level data is available in the forms of electronic health record (EHR) data, claims data, and even paper records. These data are commonly used to evaluate outcomes and

calculate quality measures. Patient-level data provide the greatest opportunity for meaningful improvement. On the other hand, patient-level data are not interoperable, and lack standardization across EHR vendors or even within the same EHR vendor.

In summary, while the environmental scan highlighted different data sources that have been used to evaluate trauma outcomes, it is important to consider the utility of different sources when assessing the universe of potential data.

## References

- 1 National Academies of Sciences, Engineering, and Medicine, Health and Medicine Division, National Academies of Sciences, Engineering, and Medicine. *A National Trauma Care System: Integrating Military and Civilian Trauma Systems to Achieve Zero Preventable Deaths after Injury*. (Downey A, Berwick D, Berwick D, et al., eds.). Washington, DC: National Academies Press; 2016. <http://www.ncbi.nlm.nih.gov/books/NBK390327/>. Last accessed September 2018.
- 2 Centers for Disease Control and Prevention (CDC). FastStats. Emergency department visits website. <https://www.cdc.gov/nchs/fastats/emergency-department.htm>. Last accessed October 2018.
- 3 Florence C, Thomas S, Haegerich T, et al. Estimated lifetime medical and work-loss costs of fatal injuries — United States, 2013. *Morbidity and Mortality Weekly Report (MMWR)*. <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm6438a4.htm>. Published October 2, 2015. Last accessed October 2018.
- 4 Kwon AM, Garbett NC, Kloecker GH. Pooled preventable death rates in trauma patients: Meta analysis and systematic review since 1990. *Eur J Trauma Emerg Surg*. 2014;40(3):279-285.
- 5 Stelfox HT, Bobranska-Artiuch B, Nathens A, et al. Quality indicators for evaluating trauma care: a scoping review. *Arch Surg*. 2010;145(3):286-295.
- 6 *Model Trauma System Planning and Evaluation*. United States: Department of Health and Human Services; 2006. <https://www.hsdl.org/?view&did=463554>. Last accessed August 2018.
- 7 Jacobs LM, Wade D, McSwain NE, et al. Hartford Consensus: A call to action for THREAT, a medical disaster preparedness concept. *J Am Coll Surg*. 2014;218(3):467-475. [https://www.journalacs.org/article/S1072-7515\(13\)01295-7/fulltext](https://www.journalacs.org/article/S1072-7515(13)01295-7/fulltext). Last accessed September 2018.
- 8 AABB. *AABB White Paper - Building a Better Patient Blood Management Program*. Bethesda, MD: American Association of Blood Banks; 2015:18.
- 9 The Department of Defense Center of Excellence for Trauma. Tactical Combat Casualty Care Guidelines. <http://jts.amedd.army.mil/index.cfm/committees/cotccc/guidelines>. Last accessed October 2018.
- 10 Santana MJ, Stelfox HT. Development and evaluation of evidence-informed quality indicators for adult injury care. *Ann Surg*. 2014;259(1):186-192.
- 11 Gruen RL, Gabbe BJ, Stelfox HT, et al. Indicators of the quality of trauma care and the performance of trauma systems. *Br J Surg*. 2012;99 Suppl 1:97-104.

- 12 Easton R, Sisak K, Balogh ZJ. Time to computed tomography scanning for major trauma patients: the Australian reality. *ANZ J Surg.* 2012;82(9):644-647.
- 13 Strudwick K, Nelson M, Martin-Khan M, et al. Quality indicators for musculoskeletal injury management in the emergency department: a systematic review. *Acad Emerg Med.* 2015;22(2):127-141.
- 14 Tignanelli CJ, Vander Kolk WE, Mikhail JN, et al. Noncompliance with American College of Surgeons Committee on Trauma recommended criteria for full trauma team activation is associated with undertriage deaths. *J Trauma Acute Care Surg.* 2018;84(2):287.  
[https://journals.lww.com/jtrauma/Abstract/2018/02000/Noncompliance\\_with\\_American\\_College\\_of\\_Surgeons.8.aspx](https://journals.lww.com/jtrauma/Abstract/2018/02000/Noncompliance_with_American_College_of_Surgeons.8.aspx). Last accessed October 2018.
- 15 Bradburn EH, Gross BW, Jammula S, et al. Improved outcomes in elderly trauma patients with the implementation of two innovative geriatric-specific protocols—final report. *J Trauma Acute Care Surg.* 2018;84(2):301.  
[https://journals.lww.com/jtrauma/Abstract/2018/02000/Improved\\_outcomes\\_in\\_elderly\\_trauma\\_patients\\_with.10.aspx](https://journals.lww.com/jtrauma/Abstract/2018/02000/Improved_outcomes_in_elderly_trauma_patients_with.10.aspx). Last accessed October 2018.
- 16 The Trauma Audit & Research Network. TARN - Home. <https://www.tarn.ac.uk/>. Last accessed October 2018.
- 17 Victorian State Trauma System. Victorian Trauma System-Key Messages | Trauma Victoria. <https://trauma.reach.vic.gov.au/guidelines/victorian-trauma-system/key-messages>. Last accessed October 2018.
- 18 Ogola GO, Crandall ML, Shafi S. Variations in outcomes of emergency general surgery patients across hospitals: a call to establish emergency general surgery quality improvement program. *J Trauma Acute Care Surg.* 2018;84(2):280-286.
- 19 Jiang HX, Majumdar SR, Dick DA, et al. Development and initial validation of a risk score for predicting in-hospital and 1-year mortality in patients with hip fractures. *J Bone Miner Res.* 2005;20(3):494-500. <http://europepmc.org/abstract/med/15746995>. Last accessed October 2018.
- 20 Shafi S, Parks J, Ahn C, et al. Centers for Medicare and Medicaid services quality indicators do not correlate with risk-adjusted mortality at trauma centers. *J Trauma.* 2010;68(4):771-777.
- 21 Moore L, Stelfox HT, Turgeon AF, et al. Derivation and validation of a quality indicator for 30-day unplanned hospital readmission to evaluate trauma care. *J Trauma Acute Care Surg.* 2014;76(5):1310-1316.
- 22 Haider AH, Hashmi ZG, Zafar SN, et al. Developing best practices to study trauma outcomes in large databases: an evidence-based approach to determine the best mortality risk adjustment model. *J Trauma Acute Care Surg.* 2014;76(4):1061-1069.
- 23 Haider AH, Saleem T, Leow JJ, et al. Influence of the National Trauma Data Bank on the study of trauma outcomes: is it time to set research best practices to further enhance its impact? *J Am Coll Surg.* 2012;214(5):756-768.

- 24 Gomez D, Byrne JP, Alali AS, et al. Inclusion of highest Glasgow coma scale motor component score in mortality risk adjustment for benchmarking of trauma center performance. *J Am Coll Surg*. 2017;225(6):755-762.
- 25 Beskind DL, Stolz U, Gross A, et al. A comparison of the prehospital motor component of the Glasgow coma scale (mGCS) to the prehospital total GCS (tGCS) as a prehospital risk adjustment measure for trauma patients. *Prehosp Emerg Care*. 2014;18(1):68-75.
- 26 Clark DE, Hannan EL, Wu C. Predicting risk-adjusted mortality for trauma patients: logistic versus multilevel logistic models. *J Am Coll Surg*. 2010;211(2):224-231.
- 27 Moore L, Hanley JA, Turgeon AF, et al. A new method for evaluating trauma centre outcome performance: TRAM-adjusted mortality estimates. *Ann Surg*. 2010;251(5):952-958.
- 28 Cook A, Weddle J, Baker S, et al. A comparison of the injury severity score and the trauma mortality prediction model. *J Trauma Acute Care Surg*. 2014;76(1):47-52; discussion 52-53.
- 29 National Quality Forum (NQF). *Regionalized Emergency Medical Care Services: Emergency Department Crowding and Boarding, Healthcare System Preparedness and Surge Capacity - Performance Measurement Gap Analysis and Topic Prioritization*. Washington, DC: National Quality Forum; 2012.  
[http://www.qualityforum.org/Publications/2012/12/REMCS\\_Emergency\\_Department\\_Crowding\\_and\\_Boarding\\_Healthcare\\_System\\_Preparedness\\_and\\_Surge\\_Capacity.aspx](http://www.qualityforum.org/Publications/2012/12/REMCS_Emergency_Department_Crowding_and_Boarding_Healthcare_System_Preparedness_and_Surge_Capacity.aspx). Last accessed September 2017.
- 30 Institute of Medicine. Regionalizing emergency care: workshop summary. In: *The National Academies Planning Committee on Regionalizing Emergency Care Workshop*. Washington, DC: The National Academies Press; 2010.
- 31 Miller WC, Powers KA, Smith MK, et al. Community viral load as a measure for assessment of HIV treatment as prevention. *Lancet Infect Dis*. 2013;13(5):459-464.  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4512165/>.
- 32 Nichol G, Thomas E, Callaway CW, et al. Regional variation in out-of-hospital cardiac arrest incidence and outcome. *JAMA*. 2008;300(12):1423-1431.  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3187919/>.
- 33 Haut ER, Mann NC, Kotwal RS. *Military Trauma Care's Learning Health System: The Importance of Data Driven Decision Making*. Washington, DC: The National Academies Press; 2016.
- 34 Glickman SW, Delgado MK, Hirshon JM, et al. Defining and measuring successful emergency care networks: a research agenda. *Soc Acad Emerg Med*. 2010;(17):1297-1305.
- 35 National Quality Forum (NQF). *Attribution: Principles and Approaches*. Washington, DC: National Quality Forum; 2016. [http://www.qualityforum.org/Publications/2016/12/Attribution\\_-\\_Principles\\_and\\_Approaches.aspx](http://www.qualityforum.org/Publications/2016/12/Attribution_-_Principles_and_Approaches.aspx). Last accessed May 2016.
- 36 Carr B, Kilaru AS, Karp DN, et al. Quality through coopetition: an empiric approach to measure population outcomes for emergency care - sensitive conditions. *Ann Emerg Med*. 2018;72(3):237-245.

- 37 HHS. *2017-2022 Health Care Preparedness and Response Capabilities*. Office of the Assistant Secretary for Preparedness and Response  
<https://www.phe.gov/Preparedness/planning/hpp/reports/Documents/2017-2022-healthcare-pr-capabilities.pdf>.
- 38 HHS. *2017-2022 Hospital Preparedness Program, Performance Measures Implementation Guidance*. Office of the Assistant Secretary for Preparedness and Response  
<https://www.phe.gov/Preparedness/planning/hpp/reports/Documents/hpp-pmi-guidance-2017.pdf>.

## Appendix A: Committee Members, NQF Staff, and Federal Liaisons

### Committee Members

**Carol Immermann, BSN, RN**

Nursing Manager, Trauma, Critical Care and General Surgery, Mayo Clinic  
Rochester, MN

**Avery Nathens, MD, MPH, PhD, FACS**

Medical Director, ACS Trauma Quality Programs, American College of Surgeons  
Chicago, IL

**Robert Bass, MD, FACEP**

Director Emeritus, Maryland Institute for EMS Systems  
Annapolis, MD

**Derek Bergsten, Paramedic, CFO, CEMSO, CTO**

Chief, Rockford Fire Department  
Rockford, IL

**Bryan Collier, DO**

Section Chief of Acute Care Surgery; Medical Director of Trauma and Surgical Critical Care, Carilion Clinic  
Roanoke, VA

**Joseph Cuschieri, MD**

Professor of Surgery, Harborview Medical Center, University of Washington  
Seattle, WA

**James Eubanks, MD, FACS**

Chief, Division of Pediatric Surgery, Le Bonheur Children's Hospital  
Memphis, TN

**Alexander Garza, MD, MPH**

Chief Quality Officer, SSM Health  
St. Louis, MO

**Michael Gonzalez, MD, FACEP, FAAEM**

Associate Medical Director, Emergency Medical Services, Houston Fire Department  
Houston, TX

**Adil Haider, MD, MPH, FACS**

Kessler Director, Center for Surgery and Public Health at Brigham and Women's Hospital, Harvard Medical School  
Boston, MA

**Kurt Hoppe, MD**

Consultant, Mayo Clinic  
Rochester, MN



**Elliott Haut, MD, PhD, FACS**

Associate Professor of Surgery, Anesthesiology & Critical Care Medicine, and Emergency Medicine, Johns Hopkins University School of Medicine  
Baltimore, MD

**Gregory Hawryluk, MD, PhD, FRCSC**

Assistant Professor of Neurosurgery, University of Utah  
Salt Lake City, UT

**David Livingston, MD**

Wesley J. Howe Professor, Chief of Trauma and Surgical Critical Care, Rutgers New Jersey Medical School  
Newark, NJ

**Barry Markman, MD, MBA, FACS**

Senior Medical Director Medicaid, Aetna  
Las Vegas, NV

**Linda Melillo, MA, MS, CPHRM, CPXP**

Olanta, PA

**Anna Miller, MD, FACS**

Associate Professor, Washington University Orthopaedics  
St. Louis, MO

**Sage Myers, MD, MSCE**

Assistant Professor of Pediatrics, Children's Hospital of Pennsylvania  
Philadelphia, PA

**Craig Newgard, MD, MPH**

Professor of Emergency Medicine, Oregon Health & Science University  
Portland, OR

**Jack Sava, MD**

Regional Vice Chair for Education; Regional Chief, Trauma/Burns; Program Director, General Surgery, MedStar Washington Hospital Center  
Washington, DC

**Andrew Schrag, MBA, MA, LPCS**

Regional Director of Community Operations, Partners Behavioral Health Management  
Hickory, NC

**David Seidenwurm, MD, FACP**

Quality & Safety Director, Sutter Health Group  
Sacramento, CA

**Theresa Snavely, MSN, RN**

Manager of Performance Improvement, Pennsylvania Trauma Systems Foundation  
Camp Hill, PA

**Peter Thomas, JD**

Principal, Powers Law Firm  
Washington, DC

**Garth Utter, MD, MSc**

Professor, University of California, Davis School of Medicine  
Sacramento, CA

**NQF Staff****Elisa Munthali, MPH**

Senior Vice President, Quality Measurement

**John Bernot**

Vice President, Quality Measurement Initiatives

**Andrew Lyzenga, MPP**

Senior Director

**Christy Skipper, MS**

Project Manager

**Jean-Luc Tilly**

Senior Manager, Data Analytics

**Federal Liaisons****Nina Heggs**

Centers for Medicare and Medicaid Services

**Brendan Carr, MD, MA, MS**

HHS Office of the Assistant Secretary for Preparedness and Response

Appendix B: Measure and Measure Concept Inventory

Measures

Measure Title	Measure Description	Measure Type	Developer
PSI 3 Pressure Ulcer Rate	Stage III or IV pressure ulcers or unstageable (secondary diagnosis) per 1,000 discharges among surgical or medical patients ages 18 years and older.		Agency for Healthcare Research and Quality
PSI 5 Retained Surgical Item or Unretrieved Device Fragment Count	The number of hospital discharges with a retained surgical item or unretrieved device fragment (secondary diagnosis) among surgical and medical patients ages 18 years and older or obstetric patients.		Agency for Healthcare Research and Quality
PSI 6 Iatrogenic Pneumothorax Rate	Iatrogenic pneumothorax cases (secondary diagnosis) per 1,000 surgical and medical discharges for patients ages 18 years and older.		Agency for Healthcare Research and Quality
PSI 7 Central Venous Catheter-Related Blood Stream Infection Rate	Central venous catheter-related bloodstream infections (secondary diagnosis) per 1,000 medical and surgical discharges for patients ages 18 years and older or obstetric cases.		Agency for Healthcare Research and Quality
PSI 9 Perioperative Hemorrhage or Hematoma Rate	Perioperative hemorrhage or hematoma cases involving a procedure to treat the hemorrhage or hematoma, following surgery per 1,000 surgical discharges for patients ages 18 years and older.		Agency for Healthcare Research and Quality
PSI 14 Postoperative Wound Dehiscence Rate	Postoperative reclosures of the abdominal wall per 1,000 abdominopelvic surgery discharged for patients aged 17 and older.		Agency for Healthcare Research and Quality
PSI 15 Accidental Puncture or Laceration Rate	Accidental punctures or lacerations (secondary diagnosis) per 1,000 discharges for patients ages 18 years and older who have undergone an abdominopelvic procedure; in which a second abdominopelvic procedure follows one or more days after an index abdominopelvic procedure.		Agency for Healthcare Research and Quality
PSI 16 Transfusion Reaction Count	The number of medical and surgical discharges with a secondary diagnosis of transfusion reaction for patients ages 18 years and older or obstetric patients.		Agency for Healthcare Research and Quality
ACEP 19 Emergency Medicine: Emergency Department Utilization of CT for Minor Blunt Head Trauma for Patients Aged 18 Years and Older	Percentage of emergency department visits for patients aged 18 years and older who presented within 24 hours of a minor blunt head trauma with a Glasgow Coma Scale (GCS) score of 15 and who had a head CT for trauma ordered by an emergency care provider who have an indication for a head CT.	Efficiency	American College of Emergency Physicians
ACEP 20 Emergency Medicine: Emergency Department Utilization of CT for Minor Blunt Head Trauma for Patients Aged 2 Through 17 Years	Percentage of emergency department visits for patients aged 2 through 17 years who presented within 24 hours of a minor blunt head trauma with a Glasgow Coma Scale (GCS) score of 15 and who had a head CT for trauma ordered by an emergency care provider who are classified as low risk according to the Pediatric Emergency Care Applied Research Network (PECARN) prediction rules for traumatic brain injury	Efficiency	American College of Emergency Physicians
ACEP22 Appropriate Emergency Department Utilization of CT for Pulmonary Embolism	Percentage of emergency department visits during which patients aged 18 years and older had a CT pulmonary angiogram (CTPA) ordered by an emergency care provider, regardless of discharge disposition, with either moderate or high pre-test clinical probability for pulmonary embolism OR positive result or elevated D-dimer level	Process	American College of Emergency Physicians
Imaging in adult ED patients with minor head injury	Percent of adult patients who presented within 24 hours of a non-penetrating head injury with a Glasgow coma score (GCS) of 15 and underwent head CT for trauma in the ED who have a documented indication consistent with guidelines prior to imaging	Process	American College of Emergency Physicians
Avoidance of inappropriate use of head CT in ED patients with minor head injury	Percentage of emergency department patients with minor head injury who received inappropriate imaging study (not clinically indicated)	Efficiency	American College of Emergency Physicians
Imaging in pediatric ED patients aged 2 through 17 years with minor head injury	Percent of pediatric patients who presented within 24 hours of a non- penetrating head injury with a Glasgow coma score (GCS) of 14 or 15 and underwent head CT for trauma in the ED who have a documented indication consistent with guidelines (PECARN) prior to imaging	Process	American College of Emergency Physicians
ACSTrauma1 Trauma Initial Assessment Composite	Percentage of blunt multisystem trauma patients having the following components documented upon presentation of the emergency department (ED) within 30 minutes of arrivals: 1. Glasgow Coma Scale score (GCS) 2. Temperature	Process	American College of Surgeons
ACSTrauma2 Mortality Rate Following Blunt Traumatic Injury to the Chest and/or Abdomen	In-hospital mortality rate for patients with severe blunt injury to the abdomen and/or chest (abbreviated injury score AIS >3)	Outcome	American College of Surgeons
ACSTrauma3 Mortality Rate Following Penetrating Traumatic Injury to the Chest and/or Abdomen	In-hospital mortality rate for patients with severe penetrating injury to the abdomen and/or chest (abbreviated injury score AIS >3)	Outcome	American College of Surgeons
ACSTrauma4 Splenic Salvage Rate	Percentage of patients with a spleen injury (spleen AIS >2 and <5) that do not undergo a splenectomy	Outcome	American College of Surgeons
ACSTrauma5 Optimal Timing of Surgical or Procedural Intervention for Hemorrhage in Trauma	Percentage of patients presenting with traumatic hemorrhagic shock who undergo an operative or procedural intervention for hemorrhage control within 4 hours	Process	American College of Surgeons
ACSTrauma6 Optimal Ratio of Blood Product Transfusion	Percentage of patients presenting with traumatic hemorrhagic shock who receive plasma and packed red blood cells (pRBC's) in a ratio or equal to 1 unit of plasma for every 2 units of pRBCs over the first four hours after arrival to the emergency department	Process	American College of Surgeons

Measure Title	Measure Description	Measure Type	Developer
<b>ACSTrauma7 Timely Initiation of VTE Prophylaxis in Trauma Patients</b>	Percentage of seriously injured patients with pharmacologic venous thromboembolism (VTE) prophylaxis initiated within 48 hours of admission.	Process	American College of Surgeons
<b>NQF 0495 Median time from ED arrival to ED departure for admitted ED patients</b>	Median time from emergency department arrival to time of departure from the emergency room for patients admitted to the facility from the emergency department	Process	Centers for Medicare & Medicaid Services
<b>NQF 0496 Median time from ED arrival to ED departure for discharged ED patients</b>	Median time from emergency department arrival to time of departure from the emergency room for patients discharged from the emergency department.	Process	Centers for Medicare & Medicaid Services
<b>NQF 0497 Admit decision time to ED departure time for admitted patients</b>	Median time from admit decision time to time of departure from the emergency department for emergency department patients admitted to inpatient status	Process	Centers for Medicare & Medicaid Services
<b>NQF 0662 Median Time to Pain Management for Long Bone Fracture (no longer endorsed)</b>	Median time from emergency department arrival to time of initial oral, nasal or parenteral pain medication administration for emergency department patients with a principal diagnosis of long bone fracture (LBF)	Process	Centers for Medicare & Medicaid Services
<b>EMSC 01 Submission of NEMSIS compliant data</b>	The degree to which EMS agencies submit NEMSIS compliant version 3.x or higher data to the state EMS office	Structure	Emergency Medical Services for Children
<b>EMSC 02 Pediatric Emergency Care Coordinator</b>	The percentage of EMS agencies in the state or territory that have a designated individual who coordinates pediatric emergency care.	Structure	Emergency Medical Services for Children
<b>EMSC 03 Use of Pediatric Specific Equipment</b>	The percentage of EMS agencies in the state or territory that have a process that requires EMS providers to physically demonstrate the correct use of pediatric specific equipment	Structure	Emergency Medical Services for Children
<b>EMSC 04 Hospital Recognition for Pediatric Medical Emergencies</b>	The percent of hospitals with an Emergency Department (ED) recognized through a statewide, territorial, or regional standardized program that are able to stabilize and/or manage pediatric medical emergencies	Structure	Emergency Medical Services for Children
<b>EMSC 05 Hospital Recognition for Pediatric Trauma</b>	The percent of hospitals with an Emergency Department (ED) recognized through a statewide, territorial, or regional standardized system that are able to stabilize and/or manage pediatric trauma	Structure	Emergency Medical Services for Children
<b>EMSC 06 Interfacility Transfer Guidelines</b>	The percent of hospitals with an Emergency Department (ED) in the state or territory that have written interfacility transfer guidelines that cover pediatric patients and that include 8 components of transfer.	Structure	Emergency Medical Services for Children
<b>EMSC 07 Interfacility Transfer Agreements</b>	The percent of hospitals with an Emergency Department (ED) in the state or territory that have written interfacility transfer agreements that cover pediatric patients	Structure	Emergency Medical Services for Children
<b>EMSC 08 Permanence of EMSC</b>	The degree to which the state or territory has established permanence of EMSC in the state or territory EMS system	Structure	Emergency Medical Services for Children
<b>EMSC 09 Integration of EMSC Priorities into Statutes or Regulations</b>	The degree to which the state or territory has established permanence of EMSC in the state or territory EMS system by integrating EMSC priorities into statutes or regulations	Structure	Emergency Medical Services for Children
<b>Trauma 01 Pain assessment of injured patients</b>	Recognizing that pain is undertreated in injured patients, it is important to assess whether a patient is experiencing pain	Process	EMS Compass Measures
<b>Trauma 02 Pain re-assessment of injured patients</b>	Recognizing that pain is undertreated in injured patients, it is important to assess whether a patient is experiencing pain	Process	EMS Compass Measures
<b>Trauma 03 Effectiveness of pain management for injured patients</b>	Of injured patients, how many had less pain	Outcome	EMS Compass Measures
<b>Trauma 04 Trauma patients transported to trauma center</b>	Trauma patients transported to trauma center	Process	EMS Compass Measures
<b>Alcohol Screening and Brief Intervention (ASBI) in the ER</b>	Percentage of patients aged 15 to 34 seen in the ER for injury who were screened for hazardous alcohol use AND provided a brief intervention within 7 days of the ER visit if screened positive.	Process	Indian Health Service
<b>NQF 0326 Advance Care Plan</b>	Percentage of patients aged 65 years and older who have an advance care plan or surrogate decision maker documented in the medical record or documentation in the medical record that an advance care plan was discussed but the patient did not wish or was not able to name a surrogate decision maker or provide an advance care plan.	Process	National Committee for Quality Assurance
<b>NQF 1626 Patients admitted to ICU who have care preferences documented</b>	Percentage of vulnerable adults admitted to ICU who survive at least 48 hours who have their care preferences documented within 48 hours OR documentation as to why this was not done.	Process	RAND Corporation
<b>NQF 1641 Hospice and Palliative Care – Treatment preferences</b>	Percentage of patients with chart documentation of preferences for life sustaining treatments.	Process	UNC Chapel Hill
<b>NQF 0291 Emergency transfer communication</b>	Percentage of patients transferred to another healthcare facility whose medical record documentation indicated that REQUIRED information was communicated to the receiving facility prior to departure (SUBSECTION 1) OR WITHIN 60 MINUTES OF TRANSFER (SUBSECTION 2-7)	Process	University of Minnesota Rural Health Research Center
<b>NQF 0292 Vital Signs (no longer endorsed)</b>	Percentage of patients transferred to another HEALTHCARE FACILITY whose medical record documentation indicated that the entire vital signs record was communicated to the receiving FACILITY within 60 minutes of departure	Process	University of Minnesota Rural Health Research Center
<b>NQF 0293 Medication Information</b>	Percentage of patients transferred to another HEALTHCARE FACILITY whose medical record documentation indicated that medication information was communicated to the receiving FACILITY within 60 minutes of departure	Process	University of Minnesota Rural Health Research Center
<b>NQF 0294 Patient Information</b>	Percentage of patients transferred to another HEALTHCARE FACILITY whose medical record documentation indicated that patient information was communicated to the receiving FACILITY within 60 minutes of departure	Process	University of Minnesota Rural Health Research Center

Measure Title	Measure Description	Measure Type	Developer
<b>NQF 0295 Physician Information</b>	Percentage of patients transferred to another HEALTHCARE FACILITY whose medical record documentation indicated that physician information was communicated to the receiving FACILITY within 60 minutes of departure	Process	University of Minnesota Rural Health Research Center
<b>NQF 0296 Nursing Information (no longer endorsed)</b>	Percentage of patients transferred to another HEALTHCARE FACILITY whose medical record documentation indicated that nursing information was communicated to the receiving FACILITY within 60 minutes of departure	Process	University of Minnesota Rural Health Research Center
<b>NQF 0297 Procedures and Tests (no longer endorsed)</b>	Percentage of patients transferred to another healthcare facility whose medical record documentation indicated that procedure and test information was communicated to the receiving FACILITY within 60 minutes of departure	Process	University of Minnesota Rural Health Research Center

### Measure Concepts

Title	Description	Type	Source
n/a	Guidelines for blood recovery and reinfusion in surgery and trauma	Structure	American Association of Blood Banks
n/a	Standards for a Patient Blood Management Program (example: There is a process for managing the blood needs of unidentified patients and resolving their identification)	Structure	American Association of Blood Banks
n/a	Standards for a Patient Blood Management Program (example: Adverse events and incidents related to transfusions)	Structure	American Association of Blood Banks
n/a	AABB Standard 5.15.1 Use of low titer O whole blood in patients with life-threatening hemorrhage	Structure	American Association of Blood Banks
n/a	Rapid decision making for anemia/coagulation management	Structure	American Association of Blood Banks
Patient not transferred from a nondesignated facility to a trauma hospital	n/a	Process	American College of Emergency Physicians
Patient transferred from a nondesignated facility to another non designated facility	n/a	Process	American College of Emergency Physicians
Patients transferred between trauma hospitals	n/a	Process	American College of Emergency Physicians
Patients transferred from a nondesignated facility to a trauma hospital	n/a	Process	American College of Emergency Physicians
90-day readmission rates	n/a	Outcome	American College of Surgeons Trauma Quality Improvement Program (ACS TQIP) Geriatric Guidelines
Benzodiazepine use	n/a	Process	ACS TQIP Geriatric Guidelines
Bowel Regimen	n/a	Process	ACS TQIP Geriatric Guidelines
Code Status	n/a	Process	ACS TQIP Geriatric Guidelines
Delirium Diagnosis	n/a	Process	ACS TQIP Geriatric Guidelines
Delirium Screening	n/a	Process	ACS TQIP Geriatric Guidelines
Discharge medications	n/a	Process	ACS TQIP Geriatric Guidelines
Discharge to higher level of care	n/a	Process	ACS TQIP Geriatric Guidelines
Goals of care discussion	n/a	Process	ACS TQIP Geriatric Guidelines
ICU Stay	n/a	Outcome	ACS TQIP Geriatric Guidelines
Initial living situation	n/a	Process	ACS TQIP Geriatric Guidelines
Initial Medication	n/a	Process	ACS TQIP Geriatric Guidelines
Length of stay	n/a	Outcome	ACS TQIP Geriatric Guidelines
Mobilization	n/a	Process	ACS TQIP Geriatric Guidelines
Triage and transfer of orthopaedic injuries	n/a	Process	ACS TQIP Management of Orthopaedic Trauma
Management and treatment of open fractures	n/a	Process	ACS TQIP Management of Orthopaedic Trauma
Damage control orthopaedic surgery	n/a	Process	ACS TQIP Management of Orthopaedic Trauma
Management and treatment for mangled extremities	n/a	Process	ACS TQIP Management of Orthopaedic Trauma
Treatment for compartment syndrome	n/a	Process	ACS TQIP Management of Orthopaedic Trauma
Management of pelvic fractures with associated hemorrhage	n/a	Process	ACS TQIP Management of Orthopaedic Trauma
Rehabilitation of the multisystem trauma patient	n/a	Outcome	ACS TQIP Management of Orthopaedic Trauma
Management of Hip Fractures in the Elderly	n/a	Process	ACS TQIP Management of Orthopaedic Trauma/American American Academy of Orthopaedic Surgeons
Management of pediatric supracondylar humerus fractures	n/a	Process	ACS TQIP Management of Orthopaedic Trauma/American American Academy of Orthopaedic Surgeons
Use of Glasgow Coma Scale with reporting of all three components (eye, verbal and motor response)	n/a	Outcome	ACS TQIP Management of Traumatic Brain Injury (TBI) Guidelines
Proper filed triage and transport for patients with suspected TBI	n/a	Process	ACS TQIP Management of TBI Guidelines
Goals of treatment (all TBI patients)	n/a	Process	ACS TQIP Management of TBI Guidelines



Title	Description	Type	Source
Goals of treatment (ICU patients)	n/a	Process	ACS TQIP Management of TBI Guidelines
Intracranial pressure monitoring	n/a	Process	ACS TQIP Management of TBI Guidelines
Three-tiered management of intracranial hypertension	n/a	Process	ACS TQIP Management of TBI Guidelines
Advanced neuromonitoring in TBI	n/a	Process	ACS TQIP Management of TBI Guidelines
Surgical management of TBI	n/a	Process	ACS TQIP Management of TBI Guidelines
Nutritional support for TBI	n/a	Process	ACS TQIP Management of TBI Guidelines
Tracheostomy for TBI	n/a	Process	ACS TQIP Management of TBI Guidelines
Timing of secondary procedures	n/a	Process	ACS TQIP Management of TBI Guidelines
Timing of pharmacologic VTE prophylaxis in TBI	n/a	Process	ACS TQIP Management of TBI Guidelines
Management of pediatric patients with TBI	n/a	Process	ACS TQIP Management of TBI Guidelines
Management of elderly patients with TBI	n/a	Process	ACS TQIP Management of TBI Guidelines
Prognostic decision making and withdrawal of medical support for severe TBI patients	n/a	Process	ACS TQIP Management of TBI Guidelines
GOS-E at 6 months for TBI patients	n/a	Outcome	ACS TQIP Management of TBI Guidelines
Development of a Massive Transfusion Protocol (MTP)	n/a	Structure	ACS TQIP Massive Transfusion (MT) in Trauma Guidelines
Criteria to trigger the activation of MTP	n/a	Structure	ACS TQIP MT in Trauma Guidelines
Blood product resuscitation	n/a	Process	ACS TQIP MT in Trauma Guidelines
Massive transfusion in the Intensive Care Unit (ICU)	n/a	Process	ACS TQIP MT in Trauma Guidelines
24:7 On-site transfusion service	n/a	Process	ACS TQIP MT in Trauma Guidelines
Transfusion endpoints	n/a	Process	ACS TQIP MT in Trauma Guidelines
Therapeutic adjuncts in massive transfusion	n/a	Process	ACS TQIP MT in Trauma Guidelines
Use of antifibrinolytics to treat blood loss and receipt of transfusion	n/a	Process	ACS TQIP MT in Trauma Guidelines
Interdisciplinary palliative care team	n/a	Structure	ACS TQIP Palliative Care
Palliative care assessment	n/a	Process	ACS TQIP Palliative Care
Goals of care conversation	n/a	Process	ACS TQIP Palliative Care
End of life care	n/a	Process	ACS TQIP Palliative Care
Frailty screen for geriatric patients	n/a	Process	ACS TQIP Palliative Care
Palliative care for pediatric patients	n/a	Process	ACS TQIP Palliative Care
Palliative care for spinal cord injury	n/a	Process	ACS TQIP Palliative Care
Palliative care for traumatic brain injury	n/a	Process	ACS TQIP Palliative Care
Documentation of palliative care activities	n/a	Process	ACS TQIP Palliative Care
Assessment and administration of fluid status	n/a	Process	Eastern Association for the Surgery of Trauma (EAST) Guidelines
Surgical management of pancreatic necrosis	n/a	Process	EAST Guidelines
Prophylaxis against VTE in pediatric trauma	n/a	Process	EAST Guidelines
Operative fixation of rib fractures after blunt trauma	n/a	Process	EAST Guidelines
Damage control resuscitation for severe traumatic hemorrhage	n/a	Process	EAST Guidelines
Surgical management of adult pancreatic injuries	n/a	Process	EAST Guidelines
Pain management for blunt thoracic trauma	n/a	Process	EAST Guidelines
Prevention of fall-related injuries in the elderly	n/a	Process	EAST Guidelines
Surgery or stenting for colonic obstruction	n/a	Process	EAST Guidelines
Management of penetrating extraperitoneal rectal injuries	n/a	Process	EAST Guidelines
Patient selection for emergency department thoracotomy	n/a	Process	EAST Guidelines
Presumptive antibiotics for tube thoracostomy in trauma	n/a	Process	EAST Guidelines
Cervical collar removal	n/a	Process	EAST Guidelines
Identification of cervical spine injuries following trauma	n/a	Process	EAST Guidelines
Cervical spine injury in blunt trauma	n/a	Process	EAST Guidelines
Evaluation and management of blunt traumatic aortic injury (BTAI)	n/a	Process	EAST Guidelines

Title	Description	Type	Source
Optimal timing of femur fracture stabilization in polytrauma patients	n/a	Process	EAST Guidelines
Timing and type of surgical treatment in Clostridium difficile-associated disease	n/a	Process	EAST Guidelines
Evaluation and management of small-bowel obstruction	n/a	Process	EAST Guidelines
Management of pulmonary contusion and flail chest	n/a	Process	EAST Guidelines
Emergency tracheal intubation immediately following traumatic injury	n/a	Process	EAST Guidelines
Screening for thoracolumbar spinal injuries in blunt trauma using MDCT scans with axial collimation	n/a	Process	EAST Guidelines
Prophylactic antibiotic use in penetrating abdominal trauma	n/a	Process	EAST Guidelines
Evaluation and management of penetrating lower extremity arterial trauma	n/a	Process	EAST Guidelines
Nonoperative management of blunt hepatic injury	n/a	Process	EAST Guidelines
Preperitoneal packing for pelvic fracture hemorrhage	n/a	Process	EAST Guidelines
Hemothorax Management	n/a	Process	EAST Guidelines
Prophylactic Antibiotic use in open fractures	n/a	Process	EAST Guidelines
Selective nonoperative management of penetrating abdominal trauma	n/a	Process	EAST Guidelines
Management of the open abdomen in trauma and emergency general surgery	n/a	Process	EAST Guidelines
Open abdomen management	n/a	Process	EAST Guidelines
Diagnosis and Management of injury in pregnant patients	n/a	Process	EAST Guidelines
Red blood cell transfusion in adult trauma and critical care	n/a	Process	EAST Guidelines
Timing of tracheostomy	n/a	Process	EAST Guidelines
Prehospital fluid resuscitation in the injured patient	n/a	Process	EAST Guidelines
Operative versus selective nonoperative management of penetrating zone II neck trauma	n/a	Process	EAST Guidelines
Nutritional support for trauma patients	n/a	Process	EAST Guidelines
Prevention of VTE in trauma patients	n/a	Process	EAST Guidelines
Treatment of traumatic brain injury with beta blockers	n/a	Process	EAST Guidelines
Evaluation and management of geriatric trauma	n/a	Process	EAST Guidelines
Renal trauma	n/a	Process	EAST Guidelines
REBOA use in hemorrhage control	n/a	Process	EAST Guidelines
Duodenal Trauma	n/a	Process	EAST Guidelines
Penetrating colon injury	n/a	Process	EAST Guidelines
Blunt cerebrovascular injury	n/a	Process	EAST Guidelines
Management of pediatric renal trauma	n/a	Process	EAST/Pediatric Trauma Society
Time to computed tomography for severe polytrauma patients	Time to computed tomography (CT) for severe polytrauma patients presenting to a level 1 trauma center (surrogate for trauma team efficacy)	Process	Easton R, Sisak K, Balogh Z. Time to computed tomography scanning for major trauma patients: the Australian reality. <i>ANZ J Surg.</i> 2012; 82:644-647
Use of Trauma Associated Severe Hemorrhage Score (TASH)	n/a	Process	Emergency Nurses Association (ENA) Guideline
Use of Assessment of Blood Consumption (ABC)	n/a	Process	Emergency Nurses Association (ENA) Guideline
All deaths	n/a	Outcome	Florida Department of Health Trauma Center Standards
n/a	Any trauma patient readmitted to ICU, or an unplanned admission to the ICU from a medical/surgical unit	Outcome	Florida Department of Health Trauma Center Standards
n/a	Any trauma patient with an unplanned readmittance to the hospital within thirty days of discharge	Outcome	Florida Department of Health Trauma Center Standards
n/a	Percentage of all traumatic C1, 2, and/or C3 spinal cord injury patients permanently dependent on mechanical ventilator support who were admitted or transferred to the ICU during the quarter or who remained in the ICU from the previous quarter; who received the diaphragm pacer surgery and were discharged to a less restrictive facility, home or home-health.	Outcome	Florida Department of Health Trauma Center Standards
Appropriate activation of massive transfusion protocol	n/a	Process	Gruen RL, Gabbe BJ, Stelfox HT et al. 2011. Indicators of the quality of trauma care and the performance of trauma systems. <i>Br J Surg</i> 2012;99(suppl1):97-104
Deaths due to haemorrhagic shock	n/a	Outcome	Gruen et al. (2011)
Dispatch criteria	n/a	Structure	Gruen et al. (2011)

Title	Description	Type	Source
Injury prevention activities	n/a	Structure	Gruen et al. (2011)
Long-term outcomes	n/a	Outcome	Gruen et al. (2011)
Prehospital deaths	n/a	Outcome	Gruen et al. (2011)
Prehospital triage	n/a	Structure	Gruen et al. (2011)
Proportion of patients returning with new alcohol-or drug-related injuries	n/a	Outcome	Gruen et al. (2011)
Rehabilitation facility in community	n/a	Structure	Gruen et al. (2011)
Return to work, adjusted for severity	n/a	Outcome	Gruen et al. (2011)
Risk adjusted mortality for head injury	n/a	Outcome	Gruen et al. (2011)
Standardized rehabilitation protocols	n/a	Structure	Gruen et al. (2011)
Time to rehabilitation consultations	n/a	Process	Gruen et al. (2011)
Total prehospital time with component parts	n/a	Process	Gruen et al. (2011)
Training	n/a	Structure	Gruen et al. (2011)
Admission to intensive care unit	n/a	Process	Sasser SM, Hunt RC, Faul M et al. Guidelines for field triage of injured patients: recommendations of the National Expert Panel on Field Triage, 2011. <i>Morbidity and Mortality Weekly Report (MMWR)</i> . <a href="https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6101a1.htm">https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6101a1.htm</a> . Published January 13, 2012. Last accessed September 2018.
Interventional radiology procedure	n/a	Process	Sasser SM, Hunt RC, Faul M et al. Guidelines for field triage of injured patients: recommendations of the National Expert Panel on Field Triage, 2011. <i>Morbidity and Mortality Weekly Report (MMWR)</i> . <a href="https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6101a1.htm">https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6101a1.htm</a> . Published January 13, 2012. Last accessed September 2018.
Major nonorthopedic surgery within 24 hours	n/a	Process	Sasser SM, Hunt RC, Faul M et al. Guidelines for field triage of injured patients: recommendations of the National Expert Panel on Field Triage, 2011. <i>Morbidity and Mortality Weekly Report (MMWR)</i> . <a href="https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6101a1.htm">https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6101a1.htm</a> . Published January 13, 2012. Last accessed September 2018.
Complications	n/a	Outcome	Holena, Daniel. “Developing an EGS PIPS Process: what to collect? What to measure?” American College of Surgeons Quality and Safety Conference. July 22 2018, Orlando FL. Conference presentation
Re-operations	n/a	Outcome	Holena, Daniel. “Developing an EGS PIPS Process: what to collect? What to measure?” American College of Surgeons Quality and Safety Conference. July 22 2018, Orlando FL. Conference presentation
Use of higher ratios of plasma to red blood cells in massive transfusions	n/a	Process	Jenkins, Donald. “Trauma Induced Coagulopathy: What is it? What can be done about it? What is the future?” 5th Annual Major John P. Pryor MD FACS Memorial Lecture. October 25, 2013. Conference presentation.
Use of thawed or liquid plasma for treating coagulopathy	n/a	Process	Jenkins, Donald. “Trauma Induced Coagulopathy: What is it? What can be done about it? What is the future?” 5th Annual Major John P. Pryor MD FACS Memorial Lecture. October 25, 2013. Conference presentation.



Title	Description	Type	Source
<b>Pre-hospital plasma transfusion</b>	Any adult injured trauma patient with >2 of the following plus evidence of active hemorrhage or traumatic brain injury: Single reading of systolic blood pressure <90 mm Hg; single reading of heart rate >120; penetrating mechanism (i.e., stabbing, gunshot); positive Focused Assessment with Ultrasound in Trauma (FAST); point of care lactate >5.0 mg/dL; point of care INR >1.5; warfarin use	Process	Jenkins, Donald. “Trauma Induced Coagulopathy: What is it? What can be done about it? What is the future?” 5th Annual Major John P. Pryor MD FACS Memorial Lecture. October 25, 2013. Conference presentation.
<b>Pre-hospital hemorrhage control with hemostatic agents</b>	n/a	Process	Bulletin of the American College of Surgeons. See something, Do Something: Improving Survival. 2015; 100(15). <a href="https://www.facs.org/~media/files/publications/bulletin/hartford%20consensus%20compendium.ashx">https://www.facs.org/~media/files/publications/bulletin/hartford%20consensus%20compendium.ashx</a> . Accessed September 2018.
<b>Pre-hospital hemorrhage control with tourniquets</b>	n/a	Process	Bulletin of the American College of Surgeons. See something, Do Something: Improving Survival. 2015; 100(15). <a href="https://www.facs.org/~media/files/publications/bulletin/hartford%20consensus%20compendium.ashx">https://www.facs.org/~media/files/publications/bulletin/hartford%20consensus%20compendium.ashx</a> . Accessed September 2018.
<b>Compliance with high-risk geriatric protocols</b>	Triages patients based on injury patterns and comorbid conditions for occult hypotension	Process	Bradburn EH, Gross B, Jammula S et al. Improved outcomes in elderly trauma patients with the implementation of two innovative geriatric-specific protocols – Final report. <i>J Trauma Acute Care Surg.</i> 2018; 84(2):301-307.
<b>Compliance with the anticoagulation and trauma (ACT) alert</b>	Streamlines the care of geriatric trauma patients on anticoagulants	Process	Bradburn EH, Gross B, Jammula S et al. Improved outcomes in elderly trauma patients with the implementation of two innovative geriatric-specific protocols – Final report. <i>J Trauma Acute Care Surg.</i> 2018; 84(2):301-307.
<b>Compliance with ACS-COT minimum criterial for full trauma team activation</b>	n/a	Process	Tignanelli CJ, Vander Kolk WE, Mikhail JN et al. Noncompliance with American College of Surgeons Committee on Trauma recommended criteria for full trauma team activation is associated with undertriage deaths. <i>J Trauma Acute Care Surg.</i> 2018;(84)2:287-294
<b>Disability for patients who survive a traumatic brain injury</b>	n/a	Outcome	Washington State Department of Health
<b>Mortality of TBI by Year</b>	n/a	Outcome	Washington State Department of Health
<b>Accessibility of field hemorrhage control equipment for law enforcement, EMS/fire/rescue and the general public</b>	n/a	Access	Hartford Consensus
<b>Documentation of the use of hemorrhage control equipment by law enforcement, EMS/fire/rescue and the general public</b>	n/a	Process	Hartford Consensus
<b>Timeliness and appropriateness of initial hemorrhage control</b>	n/a	Process	Hartford Consensus
<b>Timeliness and effectiveness of rapid extrication</b>	n/a	Process	Hartford Consensus
<b>Readiness of definitive care facilities for control of internal hemorrhage</b>	n/a	Structure	Hartford Consensus
<b>Pediatric patients needing highest-level trauma team activation</b>	Includes: airway management; thoracostomy procedure; receipt of blood; criteria for termination of resuscitation; had surgery; interventional radiology; emergency c-section; received vasopressors; received burr hole; confirmed spinal cord injury; died of injury in the ED		Joint Committee on Surgical Training
<b>HRSA # 325.132(3)( c)(ii)(A) 306.2 injury prevention activities and programs</b>	The RTN is active within the region in the monitoring and evaluation of regional	Structure	Health Resources and Services Administration (HRSA)
<b>HRSA #325.132 (3)(c )(ii)(E)302.6</b>	The region has adopted mandatory regional pre-hospital triage protocols to ensure that trauma patients are transported to an appropriate trauma center based on their injuries.	Structure	HRSA

Title	Description	Type	Source
HRSA #325.132(3)(c )(ii)( C)302.10	There are established procedures for EMS and trauma system communications for major EMS events or multiple jurisdiction incidents that are effectively coordinated with the overall regional response plans	Structure	HRSA
HRSA #325.132(3)(c )(ii)(C)302.9	There is a procedure for communications among medical facilities when arranging for inter-facility transfers including contingences for radio or telephone system failure	Structure	HRSA
HRSA #325.132(3)(c )(ii)(D)302.1	There is well-defined regional trauma system medical oversight integrating the needs of the trauma system with the medical oversight of the overall EMS system.	Structure	HRSA
HRSA #325.132(3)(c )(ii)(F)303.2	The regional trauma network...should develop procedures to insure that trauma patients are transported to an appropriate facility that is prepared to provide care.	Structure	HRSA
HRSA #325.132(3)(c )(ii)(H)303.4	When injured patients arrive at a medical facility that cannot provide the appropriate level of definitive care, there is an organized and regularly monitored system to ensure that the patients are expeditiously transferred to the appropriate, system-defined trauma facility	Structure	HRSA
HRSA #325.132(3)(c )(ii)(J)310.10	As new protocols and treatment approaches are instituted within the regional trauma system, structured processes are in plan to inform or educate all personnel of those changes in a timely manner	Structure	HRSA
HRSA #325.132(3)(c )(ii)(J)310.346	The regional trauma network establishes and ensures that appropriate levels of EMS, nursing and physician trauma training courses are provided on a regular basis	Structure	HRSA
HRSA#325.132(3)(c )(i)(F)308.1	The regional work plan addresses the integration and participation of rehabilitation services within the continuum of care for trauma patients	Structure	HRSA
HRSA#325.132(3)(c )(ii)(G)303.4	There is a regional trauma bypass protocol that provides EMS guidance for bypassing a trauma care facility for another more appropriate trauma care facility	Structure	HRSA
Preventable mortality		Outcome	Military Trauma Care Learning Health System Manual
Total Presumed Ground and Transport Time	The total presumed ground and transport time intervals for the air crews should not exceed that of the time that would been required by ground crews to get the patient to the trauma center	Process	Myers JB, Slovis CM, Eckstein M et al. Evidence-based performance measures for emergency medical services systems: a model for expanded EMS benchmarking. <i>Prehosp Emerg Care.</i> 2008;12(2):141-41
Transport Time	Transporting paramedics should limit on-scene time to less than 10 minutes or document reasons for the exception (e.g., entrapment, scene safety, etc)	Process	Myers JB, Slovis CM, Eckstein M et al. Evidence-based performance measures for emergency medical services systems: a model for expanded EMS benchmarking. <i>Prehosp Emerg Care.</i> 2008;12(2):141-41
Use of TXA in adult trauma patients with severe hemorrhagic shock (SBP <75 mm Hg) with known predictors of fibrinolysis, or with known fibrinolysis by TEG (LY30 >3%)	n/a	Process	Napolitano LM, Cohen MJ, Cotton BA et al. Tranexamic acid in trauma: How should we use it? <i>J Trauma Acute Care Surg.</i> 2013;74(6):1575-1586
Highest GCS Motor	Highest motor GCS within 24 hours of ED/Hospital arrival	Outcome	National Trauma Data Bank
Hospital Complications	Any medical complications that occurred during the patient's stay at your hospital (31 complications)	Outcome	National Trauma Data Bank
Hospital Discharge Disposition	The disposition of the patient when discharged from the hospital. (Field values include Discharged/transferred: to a short-term general hospital for inpatient care/intermediate care facility/organized home health service, hospice, court/law enforcement, inpatient rehab or designated unit, long term care hospital, psychiatric hospital or psychiatric distinct part unit of a hospital, another type of institution not defined elsewhere, to home or self-care, to skilled nursing facility; Left against medical advice or discontinued care; deceased/expired	Outcome	National Trauma Data Bank
Initial ED/Hospital Pupillary Response	Physiological response of the pupil size within 30 minutes or less of ED/hospital arrival	Process	National Trauma Data Bank
Total ICU Length of Stay	The cumulative amount of time spent in the ICU. Each partial or full day should be measured as one calendar day.	Outcome	National Trauma Data Bank
Total Ventilator Days	The cumulative amount of time spent on the ventilator. Each partial or full day should be measured as one calendar day.	Outcome	National Trauma Data Bank
Patient arrival to trauma team leader response time to bedside	The time between when a patient arrives at an LTH and the arrival of the Trauma Team Leader to the patient's bedside.	Process	<i>Regional Trauma Network Development – A guide for Ontario Hospitals.</i> <a href="https://www.criticalcareontario.ca/EN/Toolkits/Regional%20Trauma%20Network%20Development%20Guide.pdf">https://www.criticalcareontario.ca/EN/Toolkits/Regional%20Trauma%20Network%20Development%20Guide.pdf</a> . Accessed September 2018.

Title	Description	Type	Source
Referring hospital time-to-transfer	The time between when a patient arrives at a referring hospital and when the patient departs that hospital to be transferred to a LTH.	Process	<i>Regional Trauma Network Development – A guide for Ontario Hospitals.</i> <a href="https://www.criticalcareontario.ca/EN/Toolkits/Regional%20Trauma%20Network%20Development%20Guide.pdf">https://www.criticalcareontario.ca/EN/Toolkits/Regional%20Trauma%20Network%20Development%20Guide.pdf</a> . Accessed September 2018.
Avoidance of large volume IV fluid crystalloid in pre-hospital resuscitation (#13, p. 13)	n/a	Process	Kotwal RS, Butler Fk, Edgar EP et al. Saving lives on the battlefield - a joint trauma system review of pre-hospital trauma care in combined joint operating area - Afghanistan (CJOA-A) executive summary. <i>J Spec Oper Med.</i> 2013;13(1):77-85
Prevention of hypothermia	n/a	Process	Kotwal RS, Butler Fk, Edgar EP et al. Saving lives on the battlefield - a joint trauma system review of pre-hospital trauma care in combined joint operating area - Afghanistan (CJOA-A) executive summary. <i>J Spec Oper Med.</i> 2013;13(1):77-85
Infection rates after trauma	n/a	Outcome	Kotwal RS, Butler Fk, Edgar EP et al. Saving lives on the battlefield - a joint trauma system review of pre-hospital trauma care in combined joint operating area - Afghanistan (CJOA-A) executive summary. <i>J Spec Oper Med.</i> 2013;13(1):77-85
Cardiovascular reserve index of the arterial pulse as a sign of impending shock	n/a	Process	Kotwal RS, Butler Fk, Edgar EP et al. Saving lives on the battlefield - a joint trauma system review of pre-hospital trauma care in combined joint operating area - Afghanistan (CJOA-A) executive summary. <i>J Spec Oper Med.</i> 2013;13(1):77-85
Preventable death review	n/a	Outcome	Kotwal RS, Butler Fk, Edgar EP et al. Saving lives on the battlefield - a joint trauma system review of pre-hospital trauma care in combined joint operating area - Afghanistan (CJOA-A) executive summary. <i>J Spec Oper Med.</i> 2013;13(1):77-85
Missed Injury	Patients readmitted or treatment changed	Outcome	Beattie E, Mackway-Jones K. A Delphi study to identify performance indicators for emergency medicine. <i>Emerg Med J</i> 2004;21:47–50
Pain management	Pain management with ketamine, fentanyl lozenges	Process	Tactical Combat Casualty Care (TCCC) Guidelines
Tranexamic Acid (TXA) administration	If a casualty is anticipated to need significant blood transfusion (for example: presents with hemorrhagic shock, one or more major amputations, penetrating torso trauma, or evidence of severe bleeding), administer 1 gram of TXA as soon as possible but not later than 3 hours after injury; begin second infusion of 1 gm TXA after Hextend or other fluid treatment)	Process	TCCC Guidelines
Probability of Survival/TRISS	Retrospective measure of patients with same profile on TARN database. Use components of ISS, Age, Gender, GCS, Pre-existing medical conditions	Outcome	Trauma Audit and Research Network (TARN)
ED admit to discharge	Referring hospital ED triage time less than 2 hours	Process	Utah State Trauma System
Inter-hospital transfer time	Measured from time of arrival at referral hospital to time of arrival at definitive care	Process	Victorian State Trauma System
Inter-hospital transfer times with and without retrieval activation	Measured from time of arrival at referral hospital to time of arrival at definitive care	Process	Victorian State Trauma System
In-transit deaths	n/a	Outcome	Victorian State Trauma System
Major trauma ICU mechanical ventilation hours for major trauma patients with an ICU admission	n/a	Outcome	Victorian State Trauma System
Median ambulance response time	Measured from time of ambulance call to arrival at scene	Process	Victorian State Trauma System
Median pre-hospital scene time	Measured from ambulance arrival at scene to depart location time	Process	Victorian State Trauma System
Median pre-hospital time	Measured from ambulance arrival to primary hospital arrival	Process	Victorian State Trauma System
Median time to activation of ARV/PIPER for metropolitan transfers	Measured from time of arrival at a health service to retrieval activation	Process	Victorian State Trauma System

Title	Description	Type	Source
Median time to activation of ARV/PIPER for regional transfers	Measured from time of arrival at a health service to retrieval activation	Process	Victorian State Trauma System
Proportion of ICU admissions at the health service for definitive care	n/a	Process	Victorian State Trauma System
Proportion of MTS trauma team activation for Ambulance Victoria or AAV signal one trauma cases	n/a	Process	Victorian State Trauma System
Total time to an appropriate health service	Measured from time of injury to first presentation)	Process	Victorian State Trauma System

### Instruments, Scales, and Tools

Title	Description
Abbreviated Injury Score >3	
American Spinal Injury Association Impairment Scale	The ASIA scale is based on the Frankel scale, and is a clinician-administered scale used to classify the severity of injury in individuals with SCI. It identifies sensory and motor levels indicative of the highest spinal level demonstrating "unimpaired" function. Preservation of function in the sarcal segments is a key for determining the AIS grade.
Awareness questionnaire	
Beck's Anxiety Inventory	
Beck's Depression Inventory	
Brief Approach/Avoidance Coping Questionnaire	
Centre for Epidemiologic Studies Depression Scale (CES-D)	
Clinician administered PTSD Scale (CAPS)	
Cribari Grid	Measures undertriage and overtriage rates within a trauma system. Undertriage = patients with an ISS >15 for which a major or modified was not activated, and overtriage = patients with an ISS <16 for which a major was activated.
Davidson Trauma Scale	17-item measure with each item corresponding to DSM-IV symptoms of PTSD.
Dissociative Experience Scale (DES)	
EQ-5D	EQ-5D is a standardized instrument as a measure of health related quality of life that can be used in a wide range of health conditions and treatments. The Eq-5D consists of a descriptive system and the EQ VAS. The descriptive system comprises five dimensions: mobility, self-care, usual activities, pain/discomfort and anxiety/depression. The EQ VAS records the patient’s self-rated health on a vertical visual analogue scale. This can be used as a quantitative measure of health outcome that reflects the patient’s own judgement. The scores on these five dimensions can be presented as a health profile or can be converted to a single summary index number (utility) reflecting preferability compared to other health profiles.
Extended Glasgow Outcome Scale	The GOS is a global scale for functional outcome that rates patient status into one of five categories: dead, vegetative state, severe disability, moderate disability or good recovery. The extended GOS provides more detailed categorization into eight categories by subdividing the categoies of severe disability, moderate disability and good recovery into a lower and upper category.
Family Experiences of in-hospital Care Questionnaire for family members of patients with severe Traumatic Brain Injury (FECQ-TBI)	The FECQ-TBI assesses important aspects of in-hospital care in the acute and rehabilitation phases, as seen from a family perspective.
Functional Activities Questionnaire (FAQ)	
Functional Capacity Index	
Functional Independence Measure	The FIM is an 18-item, 7-level functional assessment designed to evaluate the amount of assistance required by a person with a disability to perform basical life activities safely and effectively. The FIM assessments are used clinically to monitor the outcomes of rehabilitative care as required by the Joint Commission on the Accreditation of Health Care Organizations (JCAHO) and the Commission on the Accreditation of Rehabilitative Facilities (CARF). According to VHA Directive 2000-16, medical centers are mandated to measure and track rehabilitation outcomes on all new stroke, lower-extremity amputees, and traumatic brain injury (TBI) patients using the FIM.The FIM assessments are used clinically to monitor the outcomes of rehabilitative care as required by the Joint Commission on the Accreditation of Health Care Organizations (JCAHO) and the Commission on the Accreditation of Rehabilitative Facilities (CARF). According to VHA Directive 2000-16, medical centers are mandated to measure and track rehabilitation outcomes on all new stroke, lower-extremity amputees, and traumatic brain injury (TBI) patients using the FIM.
General Health Questionnaire (GHQ)	
Glasgow Outcome Scale	
Gronigen Activity Restriction Score (GARS)	
Hannover Score for Polytrauma Outcome (HASPOC)	
Health Utilities Index	
Hospital Anxiety and Depression Scale (HADS)	
Identifying Seniors At Risk (ISAR)	Identify patients for likelihood of functional decline or poor long term outcomes
Impact of Events Scale-Revised (IES-R)	
Informant Questionnaire of Cognitive Decline in the Elderly Short Form	
Injury Severity Score	An anatomical score that measures the overall severity of injured patients
Injury Severity Score >15	
Karasek’s 31-item Job Content Questionnaire	
Katz Index of Independence in Activities of Daily Living	



Title	Description
<b>Kings Outcome Scale for Childhood Head Injury (KOSCHI)</b>	The KOSCHI provides a practical scale for paediatric head injury which will enable clinicians to describe rate and extent of recovery, and evaluate the effects of service and research interventions.
<b>Modified Functional Independence Measure</b>	
<b>Modified Physiological Triage Tool-24 (MPTT-24)</b>	A triage tool to predict the need for life saving interventions
<b>Multidimensional Health Locus of Control</b>	
<b>Musculoskeletal Function Assessment (MFA)</b>	
<b>Pediatric Quality of Life Inventory</b>	Evaluates the reported quality of life in children within a healthcare setting as per the child’s self-report or the parent/caregiver’s observations.
<b>Polytrauma Outcome (POLO) chart</b>	Assesses health related QoL, measuring many trauma related aspects of QoL and includes other tools (GOS, EUROQOL, SF-36 and the trauma outcome profile).
<b>Post Traumatic Symptom Scale (PTSS-10)</b>	
<b>Post-Traumatic Stress Disorder Checklist (PCL)</b>	
<b>Quality of Wellbeing Scale</b>	
<b>Quality Rounds Checklist</b>	Checklist to identify compliance with 16 evidence based preventative measures for various ICU complications
<b>RCS-E - Rehabilitation Complexity Scale Extended</b>	The Rehabilitation Complexity Scale Extended (RCS-E) provides a simple overall measure of Care, Nursing, Therapy, Medical and Equipment needs, and is designed to offer crude banding of complexity
<b>Short Form Health 12</b>	
<b>Short Form Health 36</b>	
<b>Sickness Impact Profile</b>	
<b>Social support Questionnaire (Fragebogen zur Sozialen Unterstutzung: F-Sozu-22)</b>	
<b>Social Support Questionnaire (SSQ)</b>	
<b>Spinal Cord Independence Measure</b>	The SCIM captures the ability of a person with spinal cord injury to complete activities of daily living. It assesses independence in 19 key areas including self care (6), respiration and sphincter management (4) and mobility (9 items)
<b>State-Trait Anxiety Inventory (STAI)</b>	
<b>Symptom Checklist 90-Revised (SCL 90R)</b>	
<b>Syndrom-Kurtz Test (SKT)</b>	
<b>T-QoL</b>	A five component, 43-item questionnaire with domains unique to trauma populations
<b>Trauma and Injury Severity Score (TRISS)</b>	
<b>Trauma Early Monitoring Prediction Tool (TEMPT)</b>	A score that can be used for the early identification of those at risk of doing poorly following minor injury.
<b>Trauma Outcomes Profile (TOP)</b>	
<b>World Health Organisations Disability Assessment Schedule II (WHODAS II)</b>	
<b>World Health Organizational Quality of Life Instrument</b>	A general questionnaire for assessment of quality of life (QOL) in both healthy populations and in various diseases subgroups.
<b>Quality of Trauma Care Patient-Reported Experience Measure (QTAC-PREM) - Short Form</b>	
<b>HCAHPS Survey</b>	
<b>(EDPEC Discharge to Community) Emergency Room Patient Survey</b>	Trial version of ED Patient Experience of Care Survey. Intended as an add-on to HCAHPS

## Appendix C: Environmental Scan Strategy

### Purpose

This document details the National Quality Forum (NQF) team's approach to conducting the environmental scan of measures for the Trauma Outcomes project.

#### Research Questions:

The environmental scan will be guided by these research questions. These questions will help to focus the NQF team's research efforts and ensure the information sources collected are relevant to the project objectives.

- What are measures currently in use that can assess trauma care outcomes?
- What are measure concepts that can assess trauma care outcomes?
- What are the measurement gaps in trauma care? What measure concepts can be translated into performance measures to fill existing measurement gaps?
- What frameworks exist related to trauma care?
- What are the key considerations related to shared accountability, attribution, and risk adjustment in developing a trauma outcomes framework?
- Add question on frameworks if possible.

### Scope

The environmental scan will begin with a broad search and gradually decrease in scope as certain settings, types of measures, or concepts are prioritized. The Committee and key informants will provide input on the inclusion of measures and concepts into the environmental scan report. NQF will only collect measures for which there is enough information to understand how the measure should be used (e.g., what is being measured, where does measurement occur, etc.). Therefore, NQF will only collect measures that have the required data elements included in Table C1. The scan will not include measures or concepts related to psychological trauma (e.g., abuse) or secondary trauma.

**Table C1. Data Elements Captured in the Environmental Scan**

Data Element	Description
Title	Name of measure
Description	Measure description
Numerator	Numerator statement*
Denominator	Denominator statement*
Measure Type	Measure type based on NQF taxonomy
Level of Analysis	Entity accountable for improving performance (e.g. state, individual provider, agency, consumer)
Setting	The setting where data are collected (outpatient, inpatient, community, etc.)
Accountability	The extent to which the measure or concept facilitates or discourages accountability

Data Element	Description
Attribution	The extent to which the current attribution approach could enhance delivery system reform
Risk adjustment	Approach to risk adjustment for outcome measures*
Data Source	Data source for measure information (i.e., inventory, database, repository)

\*Only collected for performance measures.

## Sources

The NQF team will conduct the search in a clear and transparent manner. Key informants will be used to identify seminal work, relevant ongoing efforts, as well as measures under development. Sources will be gathered over the life of the project. The search will be an iterative process with constant opportunities for feedback from the project team and the Committee. The environmental scan will include, but not be limited to a review of the peer-reviewed literature and grey literature and:

- NQF's portfolio of endorsed measures;
- Centers for Medicare & Medicaid Services (CMS) Measures Inventory, including measures under development;
- Merit-based Incentive Payment System (MIPS) and Advanced Alternative Payment Models (APM) measures
- Accountable Care Organization (ACO) and Patient Centered Medical Home (PCMH) Primary Care Measures
- Healthcare Effectiveness Data and Information Set (HEDIS)
- Agency for Healthcare Research and Quality (AHRQ) National Quality Measures Clearinghouse and National Guidelines Clearinghouse;
- General and targeted outreach to the NQF membership and the broader public;
- Recommendations from the Trauma Outcomes Committee.

The following publications will also be reviewed:

- CMS Measures Management System Blueprint, including but not limited to Chapter 1: Measure Conceptualization
- A National Trauma Care System: Integrating Military and Civilian Trauma Systems to Achieve Zero Preventable Deaths After Injury
- Resources for Optimal Care of the Injured Patient

Example Sources:

- Databases: Google Scholar, PubMed/Medline (Medicine), Academic Search (multidisciplinary), LexisNexis (News), JSTOR (multidisciplinary), and Web of Science (multidisciplinary)
- Measure Repositories: AHRQ National Quality Measures Clearinghouse (NQMC) and AHRQ National Guidelines Clearinghouse (NGC)
- Grey Literature (i.e., academic or policy literature that is not commercially published):

- Government publications (e.g., Congressional reports, federal or state agency reports, rules and regulations, etc.)
- Reports or publications from foundations, associations, or nonprofit groups (e.g., Commonwealth Fund, Kaiser Family Foundation, AcademyHealth, medical/healthcare associations or specialty societies, etc.)
- Conference papers or proceedings, abstracts

## Search Parameters

The NQF team will use the parameters defined in Table C2. The NQF team will refine the search parameters when appropriate as additional information is gathered.

**Table C2. Search Parameters**

Included	Excluded
<ul style="list-style-type: none"> <li>• Developed or published after 2000 OR originally published prior to 2000 and still current</li> <li>• Measures that include specifications that meet the operational definitions of trauma</li> <li>• Instruments, scales, survey tools, and surveys</li> </ul>	<ul style="list-style-type: none"> <li>• Published before 2000 and not current</li> <li>• Not available in English</li> <li>• Does not include data from the required data elements (Table C1)</li> </ul>

## Search Terms:

The NQF team will use specific “terms” or “strings” to search for information sources. As additional information is gathered, NQF will revisit and refine the list of terms as appropriate. Databases are searched using combinations and variations of the example search terms below. NQF will also use relevant MeSH terms. For the environmental scan of measures, these terms may be combined with terms like “measure,” “survey,” “scale,” “quality,” etc.

## Terms:

- Trauma (-tic)
- Injury (-ies)
- Traumatic injuries
- Intentional injury (-ies)
- Unintentional injury (-ies)
- Advanced trauma
- Polytrauma
- Pediatric trauma
- Geriatric trauma
- Rural trauma
- Shock trauma
- Trauma mortality



- Neurotrauma
- Prehospital
- Critical care
- Emergency care
- Undertriage
- Overtriage
- Burn
- Fracture
- Spinal cord injury
- Traumatic brain injury
- Life support

#### **Operational Definitions**

- *Trauma*: severe blunt, blast, or penetrating injury primarily caused by automobile crashes, gunshots, knife wounds, falls, battery, or burns
- *Traumatic injury*: refers to physical injuries of sudden onset and severity which require immediate medical attention
- *Performance measure*: an assessment tool that aggregates data to assess the structure, processes, and outcomes of care within and between entities (typically specifies a numerator (what/how/when), denominator (who/where/when), and exclusions (not)).
- *Measure concept*: a description of existing or potential assessment tool or instrument that includes planned target and population
- *Instrument*: an assessment tool such as a survey, scale, questions, etc.