

**Voting Results (n=20 respondents of 23)**

<u>Domains</u>	<u>Subdomains</u>													
<b>Outcome (17)</b>	Patient Experience (10)*	Intermediate Outcomes (8)	Functional Health Status (8)	Mortality (7)	Safety (5)	Re-admission (5)	Pain & Function for Elective Surgery (1)	Return to Work (1)	Prevention Services (1)	Accountability for Improvement (1)	Care Coordination (1)	Accountability for Access (1)	Health Related Quality of Life (1)	
<b>Care Coordination (15)</b>	Appropriate & Timely Follow-up (10)	Medication Management (10)	Care Transitions (9)	Effective Care Plans (5)	Communication /Education (5)	Having medical or health home (3)	Re-admission (2)	Percent in Meaningful Use (1)	Help in Navigation (1)					
<b>Cost &amp; Utilization (13)</b>	Cost per Episode (8)	Hospital Readmission (7)	Procedure Utilization (6)	Overuse, appropriateness, efficiency (3)	Emergency Department Visits (2)	Per Capita (2)	Total Risk Adjusted Cost of Care (1)	Contract prices for randomly selected procedures (1)	Hospital length of stay (1)	Admits per 1000 (1)	Imaging Use (2)	Length of Emergency Room Wait (1)	Episode Re-admission (1)	Prescribing (1)
<b>Safety (12)</b>	Healthcare Associated Infections (10)	Adverse Events (9)	Medication Safety (8)	Ambulatory Settings (2)	Mortality (2)	Cleanliness of Environment (1)	Medical Safety (1)							
<b>Patient &amp; Family Engagement (11)</b>	Patient Experience (11)	Shared Decision-making (11)	Self Management (8)	End of Life Planning (1)	Patient Friendly Policies (1)	Provider Communication (1)								
<b>Population Health (10)</b>	Effective Preventive Services (10)	Public Health Outcomes (6)	Healthy Lifestyle Behaviors (6)	Community Index of Health (5)	Social Determinants (2)	Environmental Assessment [parks, trails, etc.] (1)								

